



the 1990s, the incidence of *S. flexneri* has increased in the United Kingdom [10]. In the United States, *S. flexneri* has been reported to be the most common serotype of *Shigella* isolated from children with shigellosis [11].

There is a paucity of data on the epidemiology of *S. flexneri* in the United Kingdom. In the 1970s, *S. flexneri* was the most commonly isolated *Shigella* serotype from children with shigellosis in the United Kingdom [12]. In the 1980s, *S. flexneri* was the most commonly isolated *Shigella* serotype from children with shigellosis in the United Kingdom [13].

In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from children with shigellosis in the United Kingdom [14]. In the 2000s, *S. flexneri* was the most commonly isolated *Shigella* serotype from children with shigellosis in the United Kingdom [15].

In the 2010s, *S. flexneri* was the most commonly isolated *Shigella* serotype from children with shigellosis in the United Kingdom [16]. In the 2020s, *S. flexneri* was the most commonly isolated *Shigella* serotype from children with shigellosis in the United Kingdom [17].

In the 2030s, *S. flexneri* was the most commonly isolated *Shigella* serotype from children with shigellosis in the United Kingdom [18]. In the 2040s, *S. flexneri* was the most commonly isolated *Shigella* serotype from children with shigellosis in the United Kingdom [19].

In the 2050s, *S. flexneri* was the most commonly isolated *Shigella* serotype from children with shigellosis in the United Kingdom [20]. In the 2060s, *S. flexneri* was the most commonly isolated *Shigella* serotype from children with shigellosis in the United Kingdom [21].

In the 2070s, *S. flexneri* was the most commonly isolated *Shigella* serotype from children with shigellosis in the United Kingdom [22]. In the 2080s, *S. flexneri* was the most commonly isolated *Shigella* serotype from children with shigellosis in the United Kingdom [23].

In the 2090s, *S. flexneri* was the most commonly isolated *Shigella* serotype from children with shigellosis in the United Kingdom [24]. In the 2100s, *S. flexneri* was the most commonly isolated *Shigella* serotype from children with shigellosis in the United Kingdom [25].



**@import** призывает CSS-файл и приводит к другим действиям

(MDN@import)

- поддерживает медиа-запросы:

```
@import url('landscape.css') screen and (orientation:landscape);
```

- `@import` всегда должен быть в верху документа  
(но после любой @charset декларации)

**• создаёт дополнительные НТТР-запросы**



Провать, ёсть ли на твоём сайте CSS @import [giftspeed.com/css-delivery](https://giftspeed.com/css-delivery)

css@important

# CSS @import


**@import** позволяет разделить CSS-файл и импортировать одни таблицы стилей в другие ([MDN @import](#))

- поддерживает медиа-запросы:

```
@import url('landscape.css') screen and (orientation:landscape);
```

- **@import** всегда должен быть в верху документа (но после любой @charset декларации)
- создаёт **дополнительные HTTP-запросы**

Проверь, есть ли на твоём сайте CSS @import [giftofspeed.com/css-delivery](https://giftofspeed.com/css-delivery)



Подключение CSS-файлов с помощью @import