

chapter2_1

Nikunj Goel

9/8/2020

Including libraries

```
library(astsa)
```

2.1

In stationarity the statistical properties of a process generating a time series do not change over time. It is important because they are easier to analyze, model and investigate. These processes should be possible to predict, as the way they change is predictable.

2.2

Part a)

Mean of the time series x_t would be

$$\mu_{x,t} = E(x_t) = \beta_0 + \beta_1 t$$

which is not independent of time. It violates the first rule of time series stationarity and is not *Stationary*.

Part b)

$$y_t = x_t - x_{t-1}$$

Replacing the value of x_t in the equation above.

$$y_t = (\beta_0 + \beta_1 t + w_t) - (\beta_0 + \beta_1(t-1) + w_{t-1}) y_t = \beta_1 + w_t - w_{t-1} \mu_{y,t} = E(y_t) = \beta_1$$

The mean is constant and does not depend on time t .

$$\text{cov}(y_{t+h}, y_t) = \text{cov}((w_{t+h} - w_{t+h-1} + \beta_1), (w_t - w_{t-1} + \beta_1))$$

The covariance of (constant, variable) and covariance of (constant, constant) is zero. So the equation reduces to

$$\text{cov}(y_{t+h}, y_t) = \text{cov}(w_{t+h}, w_t) - \text{cov}(w_{t+h}, w_{t-1}) - \text{cov}(w_{t+h-1}, w_t) + \text{cov}(w_{t+h-1}, w_{t-1})$$

For $h=0$

$$2\sigma_w^2$$

For $h=1$ and $h=-1$

$$-\sigma_w^2$$

For other h

$$0$$

Which proves that the process is stationary.

Part c)

$$\mu_{v,t} = 1/3E[(\beta_0 + \beta_1(t-1) + w_{t-1} + \beta_0 + \beta_1(t) + w_t + \beta_0 + \beta_1(t+1) + w_{t+1})] \mu_{v,t} = 1/3E[(3\beta_0 + 3\beta_1 t + w_{t-1} + w_t + w_{t+1})] \mu_{v,t} = \beta_0 + \beta_1 t$$

2.3

The equation is

$$x_t = \frac{1}{4}(w_{t-1} + 2w_t + w_{t+1})$$

AutoCovariance of the equation is

$$\text{cov}(x_{t+h}, x_t) = [\frac{1}{4}(w_{t+h-1} + 2w_{t+h} + w_{t+h+1}), \frac{1}{4}(w_{t-1} + 2w_t + w_{t+1})] = \frac{1}{16}[(w_{t+h-1} + 2w_{t+h} + w_{t+h+1}), (w_{t-1} + 2w_t + w_{t+1})] \text{ for } h$$

As we can see,


$$\mu_{x,t} = \frac{1}{4}E(w_{t-1} + 2w_t + w_{t+1}) = 0$$

As the mean of the time series is time independent and the CoVariance depends on the time difference So this means the the time series x_t is **stationary**.

So AutoCorrelation is

$$\rho_x(h) = \frac{\gamma(h)}{\gamma(0)} \rho_x(h) = \begin{cases} 1, & h=0 \\ \frac{4}{6}, & h=\pm 1 \\ \frac{1}{6}, & h=\pm 2 \\ 0, & \text{otherwise} \end{cases}$$

```
ACF = c(0,0,0,1,4,6,4,1,0,0,0)/6
LAG = -5:5
tsplot(LAG,ACF,type='h',lwd=3,xlab="LAG")
abline(h=0)
points(LAG[-(4:8)],ACF[-(4:8)], pch=20)
axis(1,at=seq(-5,5, by=2))
```



2.4

Part a)

$$x_t = \phi.x_{t-1} + w_t\mu_{x,t} = \phi.E(x_{t-1}) + E(w_t)$$

As x_{t-1} is uncorrelated with w_t and ϕ cannot be one. So only condition that will satisfy is :-

$$\mu_{x,t} = 0$$

Part b)

$$\gamma_x(0) = \text{var}(x_t) = \text{var}(\phi.x_{t-1} + w_t)\gamma_x(0) = \text{var}(x_t) = \text{var}(\phi.x_{t-1}) + \text{var}(w_t) + 2\text{cov}(\phi.x_{t-1}, w_t)\gamma_x(0) = \text{var}(x_t) = \phi^2\text{var}(x_{t-1}) + \text{var}(w_t)$$

Part c)

For the values to make sense $|\phi| < 1$

Part d)

$$\gamma_x(1) = \text{cov}(x_t, x_{t-1}) = \text{cov}(\phi.x_{t-1} + w_t, x_{t-1})\gamma_x(1) = \text{cov}(\phi.x_{t-1}, x_{t-1}) = \phi\gamma_x(0)\rho_x(1) = \frac{\gamma_x(1)}{\gamma_x(0)} = \phi$$

2.5

Part a)

Using the equation

$$x_t = \delta + x_{t-1} + w_t$$

putting $t=1$

$$x_1 = \delta + w_1 x_2 = 2\delta + w_1 + w_2 x_3 = 3\delta + w_1 + w_2 + w_3$$

Assuming it be true for

$$x_{t-1} = (t-1)\delta + \sum_{k=1}^{t-1} w_k$$

$$x_t = \delta + (t-1)\delta + \sum_{k=1}^{t-1} w_k + w_t x_t = (t)\delta + \sum_{k=1}^t w_k x_{t+1} = \delta + t\delta + \sum_{k=1}^{t+1} w_k$$

By mathematical induction, we can prove that the form of the term is as mentioned.

Part b)

Mean of x_t

$$\mu_{x,t} = \delta.t + E\left(\sum_{k=1}^t w_k\right)\mu_{x,t} = \delta.t$$

AutoCovariance of x_t

$$\text{cov}(x_{t+h}, x_t) = E[(x_{t+h} - E(x_{t+h}))(x_t - E(x_t))] \text{cov}(x_{t+h}, x_t) = \text{cov}\left(\sum_{k=1}^{t+h} w_k, \sum_{k=1}^t w_k\right) \text{cov}(x_{t+h}, x_t) = \min(t, t+h) \cdot \sigma_w^2$$

Part c)

Mean of x_t depends on time and doesn't follow the first rule of stationarity.

Part d)

$$\rho_x(t-1, t) = \frac{\gamma_x(t-1, t)}{\sqrt{\gamma_x(t-1, t-1)\gamma_x(t, t)}} \rho_x(t-1, t) = \frac{\text{cov}(x_t, x_{t-1})}{\sqrt{\text{cov}(x_{t-1}, x_{t-1})\text{cov}(x_t, x_t)}} \rho_x(t-1, t) = \frac{(t-1)\sigma_w^2}{\sqrt{(t-1)\sigma_w^2(t)\sigma_w^2}} \rho_x(t-1, t) = \sqrt{\frac{t-1}{t}}$$

This implies that as time goes on ∞ the ACF loses its power.

Part e)

To make the series **Stationary** we might try differencing:

$$y_t = x_t - x_{t-1}y_t = \delta + x_{t-1} + w_t - x_{t-1}y_t = \delta + w_t$$

Check Stationary

$$E(y_t) = E(\delta + w_t)E(y_t) = \delta$$

Not dependent on t.

ACV:

$$\gamma(t+h, t) = cov(x_{t+h}, x_t)\gamma(t+h, t) = cov(\delta + w_{t+h}, \delta + w_t) \text{ when } h = 0 \gamma(t+h, t) = \sigma^2 \text{ when } h \text{ is not } 0 \gamma(t+h, t) = 0$$

This Satisfies both stationary conditions. So it is stationary.