01/20/2022 MED 14:20 FMA



## 3620 East Tremont Ave., Suite 104 Bronx, New York 10465

Patient Name:

ABIBI, THIERRY

AGE: 040Y (M)

RDI#:

3153 TA

DOB: 12/17/1981

Study Date:

07/19/2022

Exam:

MRI RT KNEE

Physician:

DR IMELDA CRUZ-BANTING

FINAL REPORT

HISTORY: 040Y withright knee pain

PRIORS: none

**PROTOCOL:** Sagittal proton density and T2 weighted images and coronal T1 and STIR images were obtained. Additional axial fat suppressed T2 weighted images were obtained. Images were obtained on a high field MRI.

## FINDINGS:

Cruciate Ligaments:Intrasubstance tear of the anterior cruciate ligament is identified. The posterior cruciate ligament is intact

Colateral Ligaments: Medial collateral ligament is intact. Intrasubstance tear of the biceps femoris tendon as well as partial tear and sprain of the iliotibial band is felt present. The fibular collateral ligament is intact.

Menisci:Tear within the posterior horn of the lateral meniscus as well as intrasubstance grade 2 signal in the anterior horn of the lateral meniscus is identified. There is also grade 2 signal posterior horn of the medial meniscus.

Osseous Structures: There is a fracture of the fibular head. The remainder the osseous structures are intact

Extensor Mechanism: The patella tendon is intact. The visualized portion of the quadriceps tendon is likewise intact. The patella refinaculae are intact.

Joint Effusion: There is a joint effusion present as well as medial and lateral soft tissue edema with a hematoma identified in the medial soft tissues adjacent to the proximal tibia

## IMPRESSION

FRACTURE OF THE FIBULAR HEAD.

TEAR WITHIN THE POSTERIOR HORN OF THE LATERAL MENISCUS AS WELL AS INTRASUBSTANCE GRADE 2 SIGNAL IN THE ANTERIOR HORN OF THE LATERAL MENISCUS AND POSTERIOR HORN OF THE MEDIAL MENISCUS.

Phone: 718,484,3989 / Fax: 718,484,8757



## 3620 East Tremont Ave., Suite 104 Bronx, New York 10465

Patient Name: ABIBL/THIERRY

AGE: 040Y (M)

RDI#:

3153 TA

DOB: 12/17/1981

Study Date:

07/19/2022

MRI RT KNEE

Exam: Physician:

DR IMELDA CRUZ-BANTING

INTRASUBSTANCE TEAR OF THE ANTERIOR CRUCIATE LIGAMENT.

INTRASUBSTANCE TEAR OF THE BICEPS FEMORIS TENDON AT THE FIBULAR ATTACHMENT AS WELL AS SPRAIN/PARTIAL TEAR OF THE ILIOTIBIAL BAND. NO DISRUPTION IS NOTED.

JOINT EFFUSION AS WELL AS MEDIAL AND LATERAL SOFT TISSUE EDEMA AND PRESUMED HEMATOMA IN THE MEDIAL SOFT TISSUES ADJACENT TO THE PROXIMAL TIBIA.

THESE RESULTS WERE DISCUSSED WITH DR. CRUZ-BANTING ON JULY 19, 2022

Thank you for referring this patient to us for evaluation.

Very truly yours,

MICHAEL CARLIN MD

Electronically signed: 07/19/2022 17:59

Phone: 718.484.3989 / Fax: 718.484.8757