

Nassau Open MRI

1570 Old Country Road · Westbury, New York 11590
tel: 516-OPEN-MRI · fax: 516-385-6488

November 23, 2021

Dr. Nasar Shahid
95 Clinton Street
Hempstead, NY 11550

Re: Sarmiento, Manuel
DOB: 07/22/1993
Date of Exam: 11/22/2021

MAGNETIC RESONANCE IMAGING OF THE RIGHT HIP

TECHNIQUE: MRI of the right hip is performed utilizing coronal STIR and T1 axial sequence as well as fast spin echo, coronal, sagittal, and axial oblique sequence.

HISTORY: A 28-year-old male patient with hip pain after injury.

INTERPRETATION: Pubic symphysis is intact. Sacroiliac joints are intact. No fracture of the bony pelvis.

No hip fracture. No avascular necrosis. Loss of contour at the femoral head-neck junction on non-oblique axial imaging.

Anterior superior labral tear. Joint effusion.

Iliopsoas is intact. No bursitis.

Gluteal tendons are intact. No bursitis.

Fat planes around the sciatic nerve are intact. Hamstrings are intact. No muscle atrophy or tear.

No free fluid in the pelvis. Pelvic lipomatosis.

No enlarged lymphadenopathy. No inguinal hernia.

IMPRESSION:

1. Cam deformity with mild hip joint narrowing, anterior superior labral tear and joint effusion. No fracture.

Thank you for the courtesy of your referral; it has been a pleasure working with you.

Sincerely,



Mark Decker, M.D.
Musculoskeletal and Spine Specialist

MD/prs/NA