



30-80 31st Street, Astoria, NY 11102
Tel: 718-335-7100 | Fax: 718-709-4136

PATIENT:	OVALLS-BARRIO, MARIA	EXAM DATE:	04/06/2022 3:19 PM
STUDY DESCRIPTION:	MRI SHOULDER WITHOUT CONTRAST	MRN:	OVAM60840
DOB:	01/21/1983	REFERRING PHYSICIAN:	Qureshi, Adnan
CLINICAL HISTORY	C/O OF LEFT SHOULDER PAIN DUE TO WORK ACCIDENT	GENDER	F

MAGNETIC RESONANCE IMAGING OF LEFT SHOULDER WITHOUT CONTRAST

HISTORY: Complains of left shoulder pain due to work accident.

TECHNIQUE: Multiplanar, multi-sequence MRI of the left shoulder was performed without intravenous contrast.

COMPARISON: None available.

OSSEOUS STRUCTURES/MARROW: Normal marrow signal.

ROTATOR CUFF:

SUPRASPINATUS: There is a partial-thickness undersurface tear of the supraspinatus tendon.
INFRASPINATUS: The infraspinatus tendon maintains intact tendon fibers. No tendon retraction is found. No skeletal muscle atrophy is seen.
TERES MINOR: The teres minor tendon maintains intact tendon fibers. No tendon retraction is found. No skeletal muscle atrophy is seen.
SUBSCAPULARIS: The subscapularis tendon maintains intact tendon fibers. No tendon retraction is found. No skeletal muscle atrophy is seen.

SUBACROMIAL/SUBDELTOID BURSA: Mild fluid in subacromial-subdeltoid bursa compatible with bursitis or may be seen with full thickness rotator cuff tear.

MUSCLES: No muscle edema or fatty muscle atrophy.



30-80 31st Street, Astoria, NY 11102
Tel: 718-335-7100 | Fax: 718-709-4136

PATIENT:	OVALLES-BARRIO, MARIA	EXAM DATE:	04/06/2022 3:19 PM
STUDY DESCRIPTION:	MRI SHOULDER WITHOUT CONTRAST	MRN:	OVAM60840
DOB:	01/21/1983	REFERRING PHYSICIAN:	Qureshi, Adnan
CLINICAL HISTORY	C/O OF LEFT SHOULDER PAIN DUE TO WORK ACCIDENT	GENDER	F

AC JOINT: Intact.

BICEPS TENDON: Intact long-head of the biceps tendon.

LABRUM/LIGAMENTS: No labral tear or ligament abnormalities.

CORACOACROMIAL LIGAMENT/ROTATOR INTERVAL: Rotator interval is normal.

GLENOHUMERAL CARTILAGE: Intact articular cartilage.

SYNOVIUM/JOINT FLUID: No joint effusion or synovial thickening.

NEUROVASCULAR STRUCTURES: Normal in course and caliber.

PERIPHERAL SOFT TISSUES: Normal.

IMPRESSION:

1. Partial-thickness undersurface tear of the supraspinatus tendon.
2. Mild fluid in subacromial-subdeltoid bursa compatible with bursitis or may be seen with full thickness rotator cuff tear.

Digitally Signed By: Imam, Nalayer
Digitally Signed Date: 04/07/2022 7:53 AM.