UK Sinha Physician, P.C.

102-31 Jamaica Ave. Richmond Hill, NY 11418 Ph: 718-480-1130 Fax: 718-480-1132

October 05, 2022

Office seen at: P.R. Medical, P.C. 79-09B Northern Boulevard Jackson Heights, NY 11372 Phone# (718) 507-1438

Re: Allotey, Emmanuel

DOB: 12/18/1963 DOA: 06/28/2022

FOLLOW-UP NOTE

CHIEF COMPLAINT: Follow up of left shoulder and neck pain.

HISTORY OF PRESENT ILLNESS: The patient presents today in followup with continued pain in the left shoulder and neck.

ADL CAPABILITIES: As a direct result of the injuries sustained in this accident, the patient states that he is unable to do the following activities: driving, lifting heavy objects, carrying, reaching overhead, laundry, shopping, running errands, and exercising.

PRESENT COMPLAINTS: Left shoulder: Left shoulder pain is 7/10, described as intermittent, sharp, stabbing, dull, achy pain. The patient is able to reach overhead, able to reach behind the back, and unable to sleep at night due to pain. Worse with range of motion and improves with rest, medication, and physical therapy.

PHYSICAL EXAMINATION: The left shoulder reveals tenderness to palpation over supraspinatus tendon region and AC joint. There is crepitus appreciated. There is no heat, swelling, erythema or deformity appreciated. Negative drop arm test. Negative cross-over test. Negative empty can test. Negative Yergason test. Negative deltoid atrophy. Positive O'Brien test. Positive impingement sign. Negative Lift-off test. Positive Hawkins test. Range of motion, as per goniometer, abduction 160/180 degrees, adduction 40/45 degrees, forward flexion 120/180 degrees, extension 50/60 degrees, internal rotation 80/90 degrees, and external rotation 80/90 degrees. Internal rotation to the sacrum. The patient has no motor or sensory deficit of the left upper extremity.

DIAGNOSTIC TESTING: MRI of the left shoulder, done on 08/07/2022, shows a midsubstance and bursal surface tear is seen anteriorly at the infraspinatus tendon, as noted. A

Allotey, Emmanuel October 05, 2022 Page 2 of 3

prominent insertional tendinopathy of the supraspinatus is seen. There is no impingement. No incidental findings are seen..

ASSESSMENT:

- 1. S46.012A Partial rotator cuff tear, left shoulder.
- 2. M24.812 Internal derangement, left shoulder.
- 3. M75.42 Impingement, left shoulder.
- 4. M25.512 Pain, left shoulder.
- 5. S49.92XA Injury, left shoulder.
- 6. M25.412 Joint effusion, left shoulder.

PLAN:

- 1. Imaging studies and clinical examinations were reviewed with the patient.
- 2. All treatment options discussed with the patient.
- 3. Cold compresses for left shoulder.
- 4. Continue anti-inflammatory and muscle relaxant medications p.r.n.
- 5. Continue physical therapy for left shoulder 3 days/week.
- 6. Discussed left shoulder arthroscopy versus conservative management with the patient. The patient states that due to continual pain and lack of relief with physical therapy and the inability to perform day-to-day activities due to pain, the patient would like to proceed with surgery.
- 7. The patient needs medical clearance prior to surgery.
- 8. It is medically necessary to perform the suggested surgery to properly diagnose the patient's condition and to objectively verify presence and severity of internal derangement and other left shoulder pathology in quantitative and qualitative terms and achieve better prognosis. This surgery is crucial to provide most specific and maximally effective treatment to this patient.
- 9. Discussed the length of the arthroscopy, the postoperative instructions, and the option of continuing with conservative management alternatives to surgery, including no surgery.
- 10. All the benefits and risks of the left shoulder arthroscopy have been discussed with the patient. The risks include, but not limited to bleeding, infection, pain, stiffness, muscle injury, nerve injury, DVT, and recurrence.
- 11. All the questions in regard to the procedure were answered.
- 12. The patient verbally consents for the arthroscopy of left shoulder and the patient will be scheduled for left shoulder surgery. On the day of the surgery, before the procedure, the surgeon will explain all related questions of the procedure. The patient will sign a written consent and be witnessed by the surgeon.
- 13. The patient will follow up 1-2 weeks postop or in 2 months if surgery has not been performed yet.

<u>CAUSALITY</u>: It is within a certain degree of medical certainty, that the history presented by the patient, the objective physical findings as well as the diagnosis rendered is causally related to the injury the patient incurred on the specified date. These current symptoms were nonexistent prior to the accident. Findings were discussed with the patient.

AFFIRMATION: Being duly licensed to practice medicine in the state of New York, pursuant to the applicable provisions of the Civil Practice Laws and Rules, I hereby affirm under penalty of perjury that the statements contained herein are true and accurate.

U.K. Sinha, MD, MS (Ortho), FAAOS Board Certified Orthopedic Surgeon

apendo & wink

UKS/AEI