



119-137 Clifford Street • Newark, New Jersey, 07105

Tel: (973) 508-1400 • Fax: (973) 522-2009

PATIENT: DOUMBIA, HOUSSENI
FILE#: 042264
DOB: 04/04/1984
DOS: 06/15/2022
PHYSICIAN: DR. CLARKE, COLIN

MRI OF THE RIGHT KNEE

Clinical History: Pain; s/p injury.

Protocol: T1 and T2 axial and sagittal images

Findings: The anterior and posterior horns of the medial and lateral menisci are normal in configuration. There is linear signal in the posterior horn of the medial meniscus not seen to definitely involve the articular cartilaginous surfaces.

There is normal cartilaginous thickness present in the weight bearing aspects of the knee.

There is partial tear of the anterior cruciate ligament present with intraarticular joint fluid. The posterior cruciate ligament is normal. There is a sprain or partial tear of the medial collateral ligament. The lateral collateral ligamentous complex is intact.

The quadriceps tendon and patellar tendons are normal. Fluid is present in the suprapatellar bursa. The medial and lateral patellar facets demonstrate normal cartilaginous thickness and retinacular attachments.

There are no acute femoral, tibial, or fibular fractures or acute osteochondral lesion.

The popliteus tendon is intact. The gastrocnemius muscles are normal in signal intensity. The popliteal fossa is unremarkable.

IMPRESSION:

1. PARTIAL TEAR OF THE ANTERIOR CRUCIATE LIGAMENT WITH INTRAARTICULAR JOINT EFFUSION.
2. SPRAIN OR PARTIAL TEAR OF THE MEDIAL COLLATERAL LIGAMENT.
3. SUPRAPATELLAR EFFUSION.

Thank you for the courtesy of this referral.

**** Electronically Signed ****

Charles DeMarco, M.D.

Diplomate, American Board of Radiology