

1575 Hillside Ave, Suite 100 New Hyde Park, NY 11040 Tel: 516 962 9544, 516 962 9599

Cell: 516 549 6963 Fax: 516 467 3130

Email: qri.newhydepark@yahoo.com

PATIENT:	ROCHER, DORIS	EXAM DATE:	08/13/2022 2:30 PM
STUDY DESCRIPTION:	MRI SHOULDER WITHOUT CONTRAST	MRN:	ROCD72154
DOB:	01/24/1949	REFERRING PHYSICIAN:	Opam, Osafradu, Md
		•	
CLINICAL HISTORY:	N/FI left shoulder PAIN AFTER MVA.	GENDER:	F

## LEFT SHOULDER MRI WITHOUT CONTRAST

HISTORY: Left shoulder pain status post motor vehicle accident

**TECHNIQUE**: Multiplanar, multi-sequence MRI of the left shoulder was performed without intravenous contrast.

**COMPARISON:** None

FINDINGS:

**OSSEOUS STRUCTURES/MARROW:** There is increased signal of the glenoid humeral joint AC joint indicating microtrabecular fracture from trauma sequela. No dislocation. There is calcification/cyst of the humeral head

## ROTATOR CUFF

SUPRASPINATUS: There is increased signal at the anterior leading edge and fluid seen inferiorly in the supraspinatus tendon indicating partial tear at the articular surface. There is no muscle tendon tear or retraction.

INFRASPINATUS: Tendinopathy of the supraspinatus tendon but no muscle tendon retraction

TERES MINOR: Normal and intact

SUBSCAPLUARIS: . Intact

1575 Hillside Ave, Suite 100 New Hyde Park, NY 11040 Tel: 516 962 9544, 516 962 9599

Cell: 516 549 6963 Fax: 516 467 3130

Email: qri.newhydepark@yahoo.com

PATIENT:	ROCHER, DORIS	EXAM DATE:	08/13/2022 2:30 PM
STUDY DESCRIPTION:	MRI SHOULDER WITHOUT CONTRAST	MRN:	ROCD72154
DOB:	01/24/1949	REFERRING PHYSICIAN:	Opam, Osafradu, Md
CLINICAL HISTORY:	N/FI left shoulder PAIN AFTER MVA.	GENDER:	F

MUSCLES: No muscle edema or fat atrophy

**SUBACROMIAL/SUBDELTOID:** Moderate fluid seen in the subacromial and subdeltoid bursa indicating bursitis

AC JOINT: AC hypertrophy with type III acromion contributing to supraspinous outlet obstruction

BICEPS TENDON: Longitudinal split of the biceps tendon and within the supra groove.

LABRUM/LIGAMENTS: No labral or ligament abnormality

CORCOACROMIAL/ROTATOR: Normal rotator interval SYNOVIUM/JOINT FLUID: No synovial hypertrophy.

Glenohumeral joint: Narrowing of the glenohumeral joint with moderate joint effusion

NEUROVASCULAR STRUCTURES: Normal in caliber and course

PERIPHERAL SOFT TISSUES: Unremarkable

## IMPRESSION:

- 1. increased signal of the glenoid humeral joint AC joint indicating microtrabecular fracture from trauma sequela.
- 2.increased signal at the anterior leading edge and fluid seen inferiorly in the supraspinatus tendon indicating partial tear at the articular surface. There is no muscle tendon tear or retraction.
- 3. Tendinopathy of the supraspinatus tendon but no muscle tendon retraction
- 4. Moderate fluid seen in the subacromial and subdeltoid bursa indicating bursitis

1575 Hillside Ave, Suite 100 New Hyde Park, NY 11040 Tel: 516 962 9544, 516 962 9599

Cell: 516 549 6963 Fax: 516 467 3130

Email: qri.newhydepark@yahoo.com

PATIENT:	ROCHER, DORIS	EXAM DATE:	08/13/2022 2:30 PM
STUDY DESCRIPTION:	MRI SHOULDER WITHOUT CONTRAST	MRN:	ROCD72154
DOB:	01/24/1949	REFERRING PHYSICIAN:	Opam, Osafradu, Md
CLINICAL HISTORY:	N/FI left shoulder PAIN AFTER MVA.	GENDER:	F

- 5. AC hypertrophy with type III acromion contributing to supraspinous outlet obstruction 6Longitudinal split of the biceps tendon and within the supra groove.
- 7.. Narrowing of the glenohumeral joint with moderate joint effusion

Digitally Signed By: Izzo, Joseph

Digitally Signed Date: 08/16/2022 11:30 AM