

Community Medical Imaging

ACR Accredited Facility
159-16 Union Tpke • Fresh Meadows, NY 11366
Tel: 718-275-1010 • Fax: 718-591-3300

ISAAC KREIZMAN, M.D. 79-09 B NORTHERN BLVD. JACKSON HEIGHTS, NY 11372

PATIENT: CRISTIAN HERNAN AYALA

DOB: 07/28/1987 DOS: 08/07/2022 CHART #: 25491

EXAM: MRI OF THE LEFT SHOULDER WITHOUT CONTRAST

HISTORY: Superior lateral pain.

TECHNIQUE: Multiplanar MR imaging of the left shoulder was performed

without contrast on Hitachi open MRI unit.

Coronal T1, T2 and STIR; Sagittal T2; Axial PD and gradient echo images of the shoulder were obtained.

COMPARISON: None.

Additional sanitizing / safety protocols recommended by the CDC were performed.

FINDINGS: There is trace fluid seen at the subacromial — subdeltoid bursa. Small subchondral cysts are seen posteriorly at the humeral head measuring 2 to 3 mm. There is slight increased signal at the mid substance posteriorly at the supraspinatus tendon. The subscapularis and teres minor tendons are intact. The anterior and posterior labra are unremarkable. There is no tear of the superior labrum. Biceps anchor is unremarkable. There is no biceps tendon tear. There is type I acromion. There is no AC separation. There is no hematoma or seroma.

IMPRESSION:

- 1. MILD TENDINITIS CHANGES ARE SEEN AT THE SUPRASPINATUS AND INFRASPINATUS TENDONS.
- 2. A MIDSUBSTANCE INTERSTITIAL TEAR IS NOTED POSTERIORLY AT THE SUPRASPINATUS TENDON.
- 3. MINIMAL FLUID IS SEEN WITHOUT EVIDENCE OF A LOOSE BODY.

Thank you for referring this patient to us.

Andrew McDonnell, MD

Neuroradiologist

Diplomate, American Board of Radiology

AM/man/pr D: 08/07/2022

when I the Downelle

E-Sig By A. McDonnell, MD on 08/08/2022 06:51:12