

UK Sinha Physician, P.C.

102-31 Jamaica Ave.
Richmond Hill, NY 11418
Ph: 718-480-1130 Fax: 718-480-1132

September 12, 2022

Office seen at:
Graham Wellness Medical P.C.
150 Graham Avenue Suite A
Brooklyn NY 11206
Phone# (718) 218-6616

Re: Rodriguez, Hector
DOB: 11/14/1954
DOA: 06/12/2022

INITIAL ORTHOPEDIC CONSULT EXAMINATION

CHIEF COMPLAINT: Left knee pain.

HISTORY OF PRESENT ILLNESS: A 69-year-old right-hand dominant male involved in a motor vehicle accident on 06/12/2022. The patient was a driver and was wearing a seatbelt. The vehicle was struck on the front driver's side. The airbags deployed. The EMS did not arrive on the scene. The police were called to the scene of the accident. The patient did not go to any hospital that same day. The patient presents today complaining of left knee pain sustained in the motor vehicle accident. The patient was attending physical therapy for the last 3 months with little relief.

WORK HISTORY: The patient is currently not working.

PAST MEDICAL HISTORY: Diabetes and asthma. There is no previous history of trauma.

PAST SURGICAL HISTORY: Right hand surgery on 09/11/2001, status post trauma.

DRUG ALLERGIES: NO KNOWN DRUG ALLERGIES.

MEDICATIONS: The patient is taking metformin 1000 mg.

SOCIAL HISTORY: The patient is a nonsmoker. The patient does not drink alcohol. The patient does not use recreational drugs.

ADL CAPABILITIES: The patient states that he can walk for 2 blocks. He can stand for 10 minutes before he has to sit. He can sit for 5 minutes before needing to change positions

secondary to pain. As a direct result of the injuries sustained in this accident, the patient states that he is unable to do the following activities: play sports, running errands, kneeling, squatting, negotiating stairs, jogging and exercising.

PRESENT COMPLAINTS: Left knee: Left knee pain is 8/10, described as constant, sharp, stabbing, dull, achy pain. The patient has difficulty rising from a chair and has difficulty going up and down stairs. The patient also notes clicking and popping. Worse with range of motion and improves with medication and physical therapy.

REVIEW OF SYSTEMS: Temperature taken at the time of exam was 98.6 degrees Fahrenheit.

General: No fever, chills, night sweats, weight gain, or weight loss.

HEENT: No double vision, eye pain, eye redness, decreased hearing, earache, ear ringing, nosebleeds, sore throat or hoarseness.

Endocrine: No cold intolerance, appetite changes or hair changes.

Skin: Clear, no rashes or lesions.

Neuro: No headaches, dizziness, vertigo or tremor.

Respiratory: No wheezing, coughing, shortness of breath or difficulty breathing. The patient has asthma.

Cardiovascular: No chest pain, murmurs, irregular heart rate or hypertension.

GI: No nausea, vomiting, diarrhea, constipation, jaundice or changes in bowel habits.

GU: No blood in urine, painful urination, loss of bladder control or urinary retention.

Hematology: No active bleeding, bruising, anemia or blood clotting disorders.

Psychiatric: No anxiety, change in sleep pattern, depression or suicidal thoughts.

PHYSICAL EXAMINATION: The patient's height is 5 feet 8 inches, weight is 162 pounds, and BMI is 24.6. The left knee reveals tenderness along the medial joint line. There is crepitus appreciated. Positive McMurray test. Negative Lachman test. Positive patellofemoral grinding test. Negative anterior drawer. Negative posterior drawer. Range of motion reveals flexion 85/130 degrees and extension 4/5 degrees. Knee is stable with varus and valgus stress test. The patient has no motor or sensory deficit of the left lower extremity.

DIAGNOSTIC TESTING: MRI of the left knee, done on 07/21/2022, shows complex tear at the posterior horn and body of medial meniscus with a displaced meniscal flap interposed between the posterior horn/root attachment and the medial femoral condyle. Edema within the posterior medial meniscocapsular junction consistent with meniscocapsular junction sprain/separation. Focal reactive subchondral marrow edema signal at the medial corner of the medial tibial plateau. Joint effusion with a small popliteal cyst. Loculated fluid/ganglion cyst formation within the popliteus tendon sheath. Edema/inflammatory change within Hoffa fat.

ASSESSMENT:

1. S83.242A Medial meniscus tear, left knee.
2. M23.92 Internal derangement, left knee.
3. M25.462 Joint effusion, left knee.
4. S80.912A Injury, left knee.
5. M25.562 Pain, left knee.

PLAN:

1. Imaging studies and clinical examinations were reviewed with the patient.
2. All treatment options discussed with the patient.
3. Cold compresses for left knee.
4. Continue anti-inflammatory and muscle relaxant medications p.r.n.
5. Continue physical therapy for left knee 3 days/week.
6. Recommend steroid injections with pain management for left knee. The patient refuses due to side effects.
7. Discussed left knee arthroscopy versus conservative management with the patient. The patient states that due to continual pain and lack of relief with physical therapy and the inability to perform day-to-day activities due to pain, the patient would like to proceed with surgery.
8. The patient needs medical clearance prior to surgery.
9. It is medically necessary to perform the suggested surgery to properly diagnose the patient's condition and to objectively verify presence and severity of internal derangement and other left knee pathology in quantitative and qualitative terms and achieve better prognosis. This surgery is crucial to provide most specific and maximally effective treatment to this patient.
10. Discussed the length of the arthroscopy, the postoperative instructions, and the option of continuing with conservative management alternatives to surgery, including no surgery.
11. All the benefits and risks of the left knee arthroscopy have been discussed with the patient. The risks include, but not limited to bleeding, infection, pain, stiffness, muscle injury, nerve injury, DVT, and recurrence.
12. All the questions in regard to the procedure were answered.
13. The patient verbally consents for the arthroscopy of left knee and the patient will be scheduled for left knee surgery. On the day of the surgery, before the procedure, the surgeon will explain all related questions of the procedure. The patient will sign a written consent and be witnessed by the surgeon.
14. The patient will follow up 1-2 weeks postop or in 2 months if surgery has not been performed yet.

CAUSALITY: It is within a certain degree of medical certainty, that the history presented by the patient, the objective physical findings as well as the diagnosis rendered is causally related to the injury the patient incurred on the specified date. These current symptoms were nonexistent prior to the accident. Findings were discussed with the patient.

AFFIRMATION: Being duly licensed to practice medicine in the state of New York, pursuant to the applicable provisions of the Civil Practice Laws and Rules, I hereby affirm under penalty of perjury that the statements contained herein are true and accurate.

A handwritten signature in black ink, consisting of a large, stylized 'S' shape with a horizontal line extending to the right.

Mellita Shakhmurov, PA-C

MS/AEI

A handwritten signature in black ink, featuring a cursive 'U.K. Sinha' with a large, stylized 'S' at the end.

U.K. Sinha, MD, MS (Ortho), FAAOS
Board Certified Orthopedic Surgeon