

1575 Hillside Ave, Suite 100 New Hyde Park, NY 11040 Tel: 516 962 9544, 516 962 9599

Cell: 516 549 6963 Fax: 516 467 3130

Email: gri.newhydepark@yahoo.com

PATIENT:	RODRIGUEZ,RAFAEL	EXAM DATE:	09/12/2022 10:41 AM
STUDY DESCRIPTION:	MRI SHOULDER WITHOUT CONTRAST	MRN:	RODR75715
DOB:	10/24/1968	REFERRING PHYSICIAN:	Jurkowich, Michael
CLINICAL HISTORY:	NF left shoulder PAIN DUE TO MVA	GENDER:	М

## X LEFT SHOULDER MRI WITHOUT CONTRAST

HISTORY: Left shoulder pain status post motor vehicle accident

**TECHNIQUE**: Multiplanar, multi-sequence MRI of the left shoulder was performed without intravenous contrast.

COMPARISON: None

FINDINGS:

**OSSEOUS STRUCTURES/MARROW** there is no fracture or dislocation. There is no abnormal bone marrow signal.

## ROTATOR CUFF

SUPRASPINATUS: There is increased signal at the anterior leading edge of the supraspinatus tendon at the greater tuberosity with fluid above and below the tendon representing partial tear.

There is no muscle or tendon retraction

INFRASPINATUS: Intact TERES MINOR: Intact SUBSCAPLUARIS: Intact.

MUSCLES: No muscle edema fatty atrophy

SUBACROMIAL/SUBDELTOID: No subacromial subdeltoid bursitis



1575 Hillside Ave, Suite 100 New Hyde Park, NY 11040 Tel: 516 962 9544, 516 962 9599

Cell: 516 549 6963 Fax: 516 467 3130

Email: gri.newhydepark@yahoo.com

RODRIGUEZ, RAFAEL	EXAM DATE:	09/12/2022 10:41 AM
MRI SHOULDER WITHOUT CONTRAST	MRN:	RODR75715
10/24/1968	REFERRING	Jurkowich, Michael
	PHYSICIAN:	
NF left shoulder PAIN DUE TO MVA	GENDER:	M
	MRI SHOULDER WITHOUT CONTRAST  10/24/1968  NF left shoulder PAIN	MRI SHOULDER WITHOUT CONTRAST  10/24/1968  REFERRING PHYSICIAN:  NF left shoulder PAIN GENDER:

AC JOINT: AC hypertrophy causing supraspinatus outlet obstruction.

BICEPS TENDON: Long head biceps tendon is normal in the bicipital groove with tenosynovitis

LABRUM/LIGAMENTS: Labrum and ligaments intact

CORCOACROMIAL/ROTATOR: Rotator interval intact GLENOHUMERAL CARTILAGE: Normal cartilage

SYNOVIUM/JOINT FLUID: No joint effusion

NEUROVASCULAR STRUCTURES: Intact PERIPHERAL SOFT TISSUES: Intact

## IMPRESSION:

1.increased signal at the anterior leading edge of the supraspinatus tendon at the greater tuberosity with fluid above and below the tendon representing partial tear. There is no muscle or tendon retraction

2.Long head biceps tendon is normal in the bicipital groove with tenosynovitis

Digitally Signed By: Izzo, Joseph

Digitally Signed Date: 09/15/2022 8:29 AM