



170-10 Cedarcroft Road, Jamaica, NY 11432
Tel: 718-206-1000 | Fax: 718-532-0633

PATIENT:	MANE DE LA ROSA, ALBERTO	EXAM DATE:	10/01/2022 12:42 PM
STUDY DESCRIPTION:	MRI ELBOW WITHOUT CONTRAST	MRN:	MAND78017
DOB:	01/07/1982	REFERRING PHYSICIAN:	Martone, Joseph
CLINICAL HISTORY:	NF PAIN DUE TO ACCIDENT	GENDER	M

MAGNETIC RESONANCE IMAGING OF RIGHT ELBOW WITHOUT CONTRAST

HISTORY: Pain due to accident.

TECHNIQUE: Multiplanar, multi-sequence MRI of right elbow was performed without intravenous contrast.

COMPARISON: None available.

FINDINGS:

OSSEOUS STRUCTURES/MARROW: No fractures or osteonecrosis.

OSSEOUS ALIGNMENT: Anatomic alignment.

ANTERIOR MUSCLES/TENDONS:

BICEPS MUSCLE/TENDON: The distal biceps muscle and tendon are intact.

BRACHIALIS MUSCLE/TENDON: The distal brachialis muscle and tendon are intact.

POSTERIOR MUSCLES/TENDON:

TRICEPS MUSCLE/TENDON: Linear interstitial tearing of the triceps tendon noted at the level of its insertion.

LATERAL ELBOW LIGAMENTS AND TENDONS:

RADIAL COLLATERAL LIGAMENT: The radial collateral ligament is intact.

LATERAL ULNAR COLLATERAL LIGAMENT: The lateral ulnar collateral ligament is intact.

COMMON EXTENSOR TENDON: The common extensor ligament is intact.

MEDIAL ELBOW LIGAMENTS AND TENDONS:

ULNAR COLLATERAL LIGAMENT: The ulnar collateral ligament is intact.



170-10 Cedarcroft Road, Jamaica, NY 11432
Tel: 718-206-1000 | Fax: 718-532-0633

PATIENT:	MANE DE LA ROSA, ALBERTO	EXAM DATE:	10/01/2022 12:42 PM
STUDY DESCRIPTION:	MRI ELBOW WITHOUT CONTRAST	MRN:	MAND78017
DOB:	01/07/1982	REFERRING PHYSICIAN:	Martone, Joseph
CLINICAL HISTORY:	NF PAIN DUE TO ACCIDENT	GENDER	M

COMMON FLEXOR TENDON: Hyperintense signal noted at the origin of the common flexor tendon compatible with partial tear.

CARTILAGE/SUBCHONDRAL BONE: Cartilage and subchondral bone intact.

SYNOVIUM /JOINT FLUID: No synovial thickening or joint effusion.

NEUROVASCULAR STRUCTURES: Normal in course and caliber.

PERIPHERAL SOFT TISSUES: Normal.

IMPRESSION:

1. Linear interstitial tearing of the triceps tendon at the level of its insertion.
2. Hyperintense signal at the origin of the common flexor tendon compatible with partial tear.

Digitally Signed By: Imam, Naiyer
Digitally Signed Date: 10/03/2022 9:15 AM.