



30-80 31st Street, Astoria, NY 11102
Tel: 718-335-7100 | Fax: 718-709-4136

PATIENT:	BODRE LINAREZ, JULIAN JR	EXAM DATE:	07/15/2022 3:02 PM
STUDY DESCRIPTION:	MRI SHOULDER WITHOUT CONTRAST	MRN:	BODJ70867
DOB:	04/03/1996	REFERRING PHYSICIAN:	Qureshi, Adnan
CLINICAL HISTORY	C/O RT SHOULDER PAIN DUE TO MVA	GENDER	M

MAGNETIC RESONANCE IMAGING OF RIGHT SHOULDER WITHOUT CONTRAST

HISTORY: Complains of right shoulder pain due to motor vehicle accident.

TECHNIQUE: Multiplanar, multi-sequence MRI of the right shoulder was performed without intravenous contrast.

COMPARISON: None available.

OSSEOUS STRUCTURES/MARROW: Normal marrow signal.

ROTATOR CUFF:

SUPRASPINATUS: There is an interstitial tear of length 5 mm noted at the attachment of the supraspinatus tendon. No skeletal muscle atrophy is seen.

INFRASPINATUS: The infraspinatus tendon maintains intact tendon fibers. No tendon retraction is found. No skeletal muscle atrophy is seen.

TERES MINOR: The teres minor tendon maintains intact tendon fibers. No tendon retraction is found. No skeletal muscle atrophy is seen.


SUBSCAPULARIS: The subscapularis tendon maintains intact tendon fibers. No tendon retraction is found. No skeletal muscle atrophy is seen.

SUBACROMIAL/SUBDELTOID BURSA: Mild fluid in subcoracoid bursa noted.

MUSCLES: No muscle edema or fatty muscle atrophy.

AC JOINT: Intact.

BICEPS TENDON: Intact long-head of the biceps tendon.



30-80 31st Street, Astoria, NY 11102
Tel: 718-335-7100 | Fax: 718-709-4136

PATIENT:	BODRE LINAREZ, JULIAN JR	EXAM DATE:	07/15/2022 3:02 PM
STUDY DESCRIPTION:	MRI SHOULDER WITHOUT CONTRAST	MRN:	BODJ70867
DOB:	04/03/1996	REFERRING PHYSICIAN:	Qureshi, Adnan
CLINICAL HISTORY	C/O RT SHOULDER PAIN DUE TO MVA	GENDER	M

LABRUM/LIGAMENTS: No labral tear or ligament abnormalities.

CORACOACROMIAL LIGAMENT/ROTATOR INTERVAL: Rotator interval is normal.

GLENOHUMERAL CARTILAGE: Intact articular cartilage.

SYNOVIUM/JOINT FLUID: No joint effusion or synovial thickening.

NEUROVASCULAR STRUCTURES: Normal in course and caliber.

PERIPHERAL SOFT TISSUES: Normal.

IMPRESSION:

1. Interstitial tear of length 5 mm at the attachment of the supraspinatus tendon.
2. Mild fluid in subcoracoid bursa.

Digitally Signed By: Imam, Naiyer
Digitally Signed Date: 07/16/2022 5:43 PM

