

with And the formula we the fixed 120 to differ the Figure New party g and g are g and g and g and g are g and g and g are g and

PATIENT NAME: SCOTT, MAKAYLYNN

PATIENT DOB: 01/07/2008 CHART NUMBER: 045094 EXAM DATE: MAY 25, 2022

REFERRING PHYSICIAN: DR.KOPACH, ALEKSANDR

MRI OF THE LEFT SHOULDER

Clinical History: Pain; s/p injury.

Protocol: T1 and T2 axial and sagittal images

Findings: The acromioclavicular joint is intact, but slightly widened. There is no fracture or edema of the distal clavicle.

The supraspinatus, infraspinatus, and subscapularis tendons are intact. There is a somewhat frayed appearance to the bursal surface of the distal supraspinatus tendon, a finding suspicious for partial tear.

The humeral head is normal in position and appearance. There is no fracture or bone marrow edema. The long head of the biceps tendon is normally situated in the bicipital groove.

The anterior and posterior labrum are intact. The inferior glenohumeral ligamentous complex is intact.

The visualized musculature is normal in signal intensity. There is no subcutaneous mass or abnormal fluid collection.

IMPRESSION:

- 1. WIDENING OF THE ACROMIOCLAVICULAR JOINT. NO FRACTURES OR EDEMA OF THE DISTAL CLAVICLE.
- 2. FRAYED BURSAL SURFACE OF THE DISTAL SUPRASPINATUS TENDON, A FINDING CONSISTENT WITH PARTIAL TEAR.

Thank you for the courtesy of this referral.

**** Electronically Signed ****
Charles DeMarco, M.D.
Diplomate, American Board of Radiology