UK Sinha Physician, P.C.

102-31 Jamaica Ave. Richmond Hill, NY 11418 Ph: 718-480-1130 Fax: 718-480-1132

October 11, 2022

Office seen at: JL Medical PC 172-17 Jamaica Avenue Jamaica, NY 11432 Phone# (929) 499-3003

Re: Bravo, Kim DOB: 08/24/1966 DOA: 07/07/2022

FOLLOW-UP NOTE

CHIEF COMPLAINT: Follow up of left knee pain.

HISTORY OF PRESENT ILLNESS: The patient presents today in followup with continued pain in the left knee.

ADL CAPABILITIES: The patient states that she can walk for 1/2 block. She can stand for 5 minutes before she has to sit. She can sit for 5 minutes before needing to change positions secondary to pain. As a direct result of the injuries sustained in this accident, the patient states that she is unable to do the following activities: carrying heavy objects, laundry, shopping, running errands, kneeling, squatting, negotiating stairs, jogging, and exercising.

PRESENT COMPLAINTS: Left knee: Left knee pain is 9-10/10, described as constant, sharp, stabbing, dull, achy pain. The patient has stiffness and weakness. The patient has difficulty rising from a chair and has difficulty going up and down stairs. The patient also notes popping, buckling, and intermittent locking. Worse with range of motion and improves with rest.

PHYSICAL EXAMINATION: The left knee reveals tenderness along the medial joint line, superior pole of patella, and inferior pole of the patella. There is swelling and crepitus appreciated. There is no heat, erythema or deformity appreciated. Positive McMurray test. Negative Lachman test. Negative patellofemoral grinding test. Negative anterior drawer. Negative posterior drawer. Range of motion reveals flexion 110/130 degrees and extension - 5/5 degrees. Knee is stable with varus and valgus stress test. The patient has no motor or sensory deficit of the left lower extremity.

DIAGNOSTIC TESTING: MRI of the left knee, done on 07/26/2022, shows joint effusion. Partial LCL sprain. Partial ACL tear. Tear of the posterior horn of the medial meniscus.

Bravo, Kim October 11, 2022 Page 2 of 3

Quadriceps and patellar tendinosis/tendonitis. Prepatellar edema and/or bursitis. Lateral patellar tilt and luxation. High grade chondromalacia involving multiple patellar facets.

ASSESSMENT:

- 1. M23.92 Internal derangement, left knee.
- 2. M94.262 Chondromalacia, left knee.
- 3. M25.462 Joint effusion, left knee.
- 4. M12.569 Traumatic arthropathy, left knee.
- 5. S80.912A Injury, left knee.
- 6. M25.562 Pain, left knee.

PLAN:

- 1. Imaging studies and clinical examinations were reviewed with the patient.
- 2. All treatment options discussed with the patient.
- 3. Cold compresses for left knee.
- 4. Continue anti-inflammatory and muscle relaxant medications p.r.n.
- 5. Continue physical therapy for left knee 3 days/week.
- 6. Discussed left knee arthroscopy versus conservative management with the patient. The patient states that due to continual pain and lack of relief with physical therapy and the inability to perform day-to-day activities due to pain, the patient would like to proceed with surgery.
- 7. The patient needs medical clearance prior to surgery.
- 8. It is medically necessary to perform the suggested surgery to properly diagnose the patient's condition and to objectively verify presence and severity of internal derangement and other left knee pathology in quantitative and qualitative terms and achieve better prognosis. This surgery is crucial to provide most specific and maximally effective treatment to this patient.
- 9. Discussed the length of the arthroscopy, the postoperative instructions, and the option of continuing with conservative management alternatives to surgery, including no surgery.
- 10. All the benefits and risks of the left knee arthroscopy have been discussed with the patient. The risks include, but not limited to bleeding, infection, pain, stiffness, muscle injury, nerve injury, DVT, and recurrence.
- 11. All the questions in regard to the procedure were answered.
- 12. The patient verbally consents for the arthroscopy of left knee and the patient will be scheduled for left knee surgery. On the day of the surgery, before the procedure, the surgeon will explain all related questions of the procedure. The patient will sign a written consent and be witnessed by the surgeon.
- 13. The patient will follow up 1-2 weeks postop or in 2 months if surgery has not been performed yet.

<u>CAUSALITY</u>: It is within a certain degree of medical certainty, that the history presented by the patient, the objective physical findings as well as the diagnosis rendered is causally related to the injury the patient incurred on the specified date. These current symptoms were nonexistent prior to the accident. Findings were discussed with the patient.

Bravo, Kim October 11, 2022 Page 3 of 3

AFFIRMATION: Being duly licensed to practice medicine in the state of New York, pursuant to the applicable provisions of the Civil Practice Laws and Rules, I hereby affirm under penalty of perjury that the statements contained herein are true and accurate.

U.K. Sinha, MD, MS (Ortho), FAAOS Board Certified Orthopedic Surgeon

apenas & Jink

UKS/AEI