UK Sinha Physician, P.C.

102-31 Jamaica Ave. Richmond Hill, NY 11418 Ph: 718-480-1130 Fax: 718-480-1132

July 5, 2022

Office seen at: Rehab Time PT PC 2088B Flatbush Avenue Brooklyn, NY 11234 Phone # (718) 975-8179

Re: Martino, Emil DOB: 02/13/1972 DOA: 04/23/2022

FOLLOW-UP NOTE

CHIEF COMPLAINT: Follow up of right shoulder and left shoulder pain.

HISTORY OF PRESENT ILLNESS: The patient presents today in followup with continued pain in the right shoulder and left shoulder.

ADL CAPABILITIES: As a direct result of the injuries sustained in this accident, the patient states that he is unable to do the following activities: driving, lifting heavy objects, carrying, reaching overhead, laundry, shopping, running errands, and exercising.

PRESENT COMPLAINTS: Right shoulder: Right shoulder pain is 8/10, described as constant, sharp, stabbing, dull, achy pain. The patient has stiffness, weakness, popping, and clicking. The patient is unable to reach overhead or behind the back and unable to sleep at night due to pain. Worse with range of motion and improves with rest, medication, physical therapy, and ice.

Left shoulder: Left shoulder pain is 9/10, described as constant, sharp, stabbing, dull, achy pain. The patient has stiffness, weakness, popping, and clicking. The patient is unable to reach overhead or behind the back and unable to sleep at night due to pain. Worse with range of motion and improves with rest, medication, physical therapy, and ice.

PHYSICAL EXAMINATION: The right shoulder reveals tenderness to palpation over supraspinatus tendon region, coracoid and deltoid. There is no heat, swelling, erythema, crepitus or deformity appreciated. Negative drop arm test. Negative cross-over test. Negative empty can test. Negative Yergason test. Negative deltoid atrophy. Positive O'Brien test. Positive impingement sign. Positive Lift-off test. Positive Hawkins test. Range of motion, as per goniometer, abduction 110/180 degrees, adduction 40/45 degrees, forward flexion 110/180 degrees, extension 45/60 degrees, internal rotation 50/90 degrees, and external

rotation 65/90 degrees. Internal rotation to the sacrum. The patient has no motor or sensory deficit of the right upper extremity.

The left shoulder reveals tenderness to palpation over supraspinatus tendon region and deltoid. There is no heat, swelling, erythema, crepitus or deformity appreciated. Negative drop arm test. Negative cross-over test. Negative empty can test. Negative Yergason test. Negative deltoid atrophy. Positive O'Brien test. Positive impingement sign. Positive Lift-off test. Positive Hawkins test. Range of motion, as per goniometer, abduction 110/180 degrees, adduction 35/45 degrees, forward flexion 100/180 degrees, extension 45/60 degrees, internal rotation 50/90 degrees, and external rotation 65/90 degrees. Internal rotation to the sacrum. The patient has no motor or sensory deficit of the left upper extremity.

DIAGNOSTIC TESTING: MRI of the right shoulder, done on 05/17/2022, shows mild subluxation of the acromioclavicular joint with significant hypertrophy of the joint capsule. Postsurgical changes in the humeral head.

ASSESSMENT:

- 1. M24.811 Internal derangement, right shoulder.
- 2. M75.01 Adhesive capsulitis, right shoulder.
- 3. M75.81 Shoulder tendinitis, right shoulder.
- 4. S43.431A Labral tear, right shoulder.
- 5. M75.41 Impingement, right shoulder.
- 6. M65.811 Tenosynovitis, right shoulder.
- 7. M75.51 Bursitis, right shoulder.
- 8. M75.21 Bicipital tendinitis, right shoulder.
- 9. M25.511 Pain, right shoulder.
- 10. S49.91XA Injury, right shoulder.
- 11. M67.211 Hypertrophic synovitis, right shoulder.
- 12. M89.311 Acromioclavicular joint hypertrophy, right shoulder.
- 13. M25.411 Joint effusion, right shoulder.
- 14. M24.812 Internal derangement, left shoulder.
- 15. M75.02 Adhesive Capsulitis, left shoulder.
- 16. M75.82 Shoulder tendinitis, left shoulder.
- 17. S43.432A Labral tear, left shoulder.
- 18. M75.42 Impingement, left shoulder.
- 19. M65.812 Tenosynovitis, left shoulder.
- 20. M75.52 Bursitis, left shoulder.
- 21. M75.22 Bicipital Tendinitis, left shoulder.
- 22. M25.512 Pain, left shoulder.
- 23. S49.92XA Injury, left shoulder.
- 24. M67.212 Hypertrophic synovitis, left shoulder.
- 25. M25.412 Joint effusion, left shoulder.

PLAN:

- 1. Imaging studies and clinical examinations were reviewed with the patient.
- 2. All treatment options discussed with the patient.

- 3. Cold compresses for right shoulder and left shoulder.
- 4. Continue anti-inflammatory and muscle relaxant medications p.r.n.
- 5. Continue physical therapy for right shoulder and left shoulder 3 days/week.
- **6.** Discussed right shoulder and left shoulder arthroscopy versus conservative management with the patient. The patient states that due to continual pain and lack of relief with physical therapy and the inability to perform day-to-day activities due to pain, the patient would like to proceed with surgery.
- 7. The patient needs medical clearance prior to surgery.
- 8. It is medically necessary to perform the suggested surgery to properly diagnose the patient's condition and to objectively verify presence and severity of internal derangement and other right shoulder and left shoulder pathology in quantitative and qualitative terms and achieve better prognosis. This surgery is crucial to provide most specific and maximally effective treatment to this patient.
- 9. Discussed the length of the arthroscopy, the postoperative instructions, and the option of continuing with conservative management alternatives to surgery, including no surgery.
- 10. All the benefits and risks of the right shoulder and left shoulder arthroscopy have been discussed with the patient. The risks include, but not limited to bleeding, infection, pain, stiffness, muscle injury, nerve injury, DVT, and recurrence.
- 11. All the questions in regard to the procedure were answered.
- 12. The patient verbally consents for the arthroscopy of right shoulder and the patient will be scheduled for right shoulder surgery. On the day of the surgery, before the procedure, the surgeon will explain all related questions of the procedure. The patient will sign a written consent and be witnessed by the surgeon.
- 13. The patient will follow up 1-2 weeks postop or in 2 months if surgery has not been performed yet.

<u>CAUSALITY</u>: It is within a certain degree of medical certainty, that the history presented by the patient, the objective physical findings as well as the diagnosis rendered is causally related to the injury the patient incurred on the specified date. These current symptoms were nonexistent prior to the accident. Findings were discussed with the patient.

AFFIRMATION: Being duly licensed to practice medicine in the state of New York, pursuant to the applicable provisions of the Civil Practice Laws and Rules, I hereby affirm under penalty of perjury that the statements contained herein are true and accurate.

U.K. Sinha, MD, MS (Ortho), FAAOS

Board Certified Orthopedic Surgeon

Upenon k Dink

Martino, Emil July 5, 2022 Page 2 of 2

MS/AEI