damadian mri in canarsie, p.c.

2035 Ralph Avenue, Suite A-5, Brooklyn, NY 11234 t 718.209.1070 f 718.209.1138

ROMARIO POWELL

N10114570-CA Report Date:

07/26/2022

DOB:

02/02/2002

Exam Date: 07/23/2022

GORDON DAVIS, DO 1611 EAST NEW YORK AVE BROOKLYN, NY 11212-

MAGNETIC RESONANCE IMAGING OF THE RIGHT KNEE

TECHNIQUE: Multiplanar, multisequential MRI was performed in the recumbent position.

HISTORY: The patient complains of right knee pain, status post MVA.

INTERPRETATION: Synovial fluid is identified in the knee joint.

There is lateral patellar tilt and lateral patellar subluxation with narrowing of the lateral patellofemoral joint compartment with patellofemoral chondromalacia.

Patellar tendinosis/tendinopathy is also identified. Fluid posterior to the distal lateral patellar tendon is compatible with localized bursitis/synovitis.

Osseous signal and morphology are otherwise unremarkable. The medial and lateral menisci, the medial and lateral collateral ligaments, the anterior and posterior cruciate ligaments, quadriceps tendon are otherwise unremarkable.

IMPRESSION:

- Synovial fluid in the knee joint.
- Lateral patellar tilt and lateral patellar subluxation with narrowing of the lateral patellofemoral joint compartment with patellofemoral chondromalacia.
- Patellar tendinosis/tendinopathy.

ROMARIO POWELL

N10114570

Exam Date:

07/23/2022

Page 2 of 2 KNEE RIGHTMRI 73721

• Fluid posterior to the distal lateral patellar tendon is compatible with localized bursitis/synovitis.

Thank you for referring your patient to us for evaluation.

Sincerely,

Robert Diamond, M.D.

Diplomate of the American Board of Radiology

RD/BC