


## Nova Medical Diagnostic, PC

6317 Ave N • Brooklyn, NY 11234 Tel: 718-676-7828 • Fax: 718-676-7829



DENNY X. RODRIGUEZ, M.D. 4720 AVENUE N BROOKLYN, NY 11234

PATIENT: JENNIFER LAGUERRE

DOB: 10/26/1976 DOS: 06/01/2022 CHART #: 4099 EXAM: MRI OF THE LEFT SHOULDER WITHOUT CONTRAST

HISTORY: Shoulder pain, radiating down the arm.

TECHNIQUE: Multiplanar MR imaging of the left shoulder was performed without contrast on Hitachi open MRI unit.

Coronal T1, T2 and STIR; Sagittal T2; Axial PD and gradient echo images of the shoulder were obtained.

COMPARISON: None.

FINDINGS: Bone Marrow: No bone marrow edema, bony lesion or fracture identified.

There is fluid surrounding the acromioclavicular joint.

The glenohumeral joint is preserved.

There is small effusion in the glenohumeral joint.

Increased signal is noted in the distal aspect of the supraspinatus tendon along the articular surface suggesting a low-grade non-retracted tear.

The infraspinatus, long head of the biceps and subscapularis tendons are normal in signal and appearance.

The glenoid labrum demonstrates no abnormality.

The rotator cuff muscles are normal in signal and appearance.

## IMPRESSION:


1. PARTIAL ARTICULAR SURFACE TEAR OF THE DISTAL ASPECT OF THE SUPRASPINATUS TENDON.

Movareres about 160 to beautiful the second of the second



## Nova Medical Diagnostic, PC

6317 Ave N • Brooklyn, NY 11234 Tel: 718-676-7828 • Fax: 718-676-7829



PATIENT: JENNIFER LAGUERRE

DOB: 10/26/1976 DOS: 06/01/2022 CHART #: 4099 EXAM: MRI OF THE LEFT SHOULDER WITHOUT CONTRAST

PAGE 2

2. FLUID SURROUNDING THE ACROMIOCLAVICULAR JOINT.

3. SMALL EFFUSION IN THE GLENOHUMERAL JOINT.

Thank you for referring this patient to us.

Guenadi Amoachi, MD

6 Amachi

Diagnostic Radiologist

Diplomate, American Board of Radiology

E-Sig By G. Amoachi, MD on 06/02/2022 07:50:26