



## ***Next Generation Diagnostic Imaging P.C.***

**MERSHAD HAGIGI, MD, PHD**  
**DIPLOMATE, AMERICAN BOARD OF RADIOLOGY**

1664 East 14<sup>th</sup> Street, Suite LL • Brooklyn, NY 11229  
TEL: (718) 336-1865 • FAX: (718) 336-1275

DATE OF STUDY: 9/20/2022  
PATIENT NAME: BRADY ANDRE  
DATE OF BIRTH: 9/14/1997  
PATIENT NUMBER: SR3046  
REFERRING PHYSICIAN: DR.CARLOTTA ROSS-DISTINE

### **MRI OF THE LEFT SHOULDER WITHOUT CONTRAST**

**HISTORY:** Patient was involved in a motor vehicle accident and now complains of pain.

**COMPARISON:** None

**TECHNIQUE:** MRI of the left shoulder was performed using T1 and T2 weighted sequences in multiple planes.

### **FINDINGS:**

Minimal fluid is seen in subacromial – subdeltoid bursa and along the biceps tendon.

There is mild synovial effusion in the gleno-humeral joint. Mild lateral downsloping of the acromion is seen.

The alignment of the shoulder joint is normal. The muscles and their attachments also appear normal. The rotator cuff is well visualized. There is no evidence of oedema / tear. The bones around the shoulder joint reveal normal intensity.

The gleno-humeral joint is normal. There is no evidence of erosion or destruction of articular cartilage. The articular margins are intact.

The glenoid labrum is normal. The acromio-clavicular joint also appears normal. Major neurovascular bundles are normal.

(Continued on Page Two)

DATE OF STUDY: 9/20/2022  
PATIENT NAME: BRADY ANDRE  
DATE OF BIRTH: 9/14/1997  
PATIENT NUMBER: SR3046  
REFERRING PHYSICIAN: DR.CARLOTTA ROSS-DISTINE

**IMPRESSION:**

1. Minimal fluid in subacromial – subdeltoid bursa and along the biceps tendon.
2. Mild synovial effusion in the gleno-humeral joint.
3. Mild lateral downsloping of the acromion.

Thank you for the courtesy of this referral.

Electronically Signed  
Merhad Hagigi, MD, PHD  
Board Certified Radiologist  
Date: 9/21/2022