

40-34 74th Street, Elmhurst, NY 11373 Tel: 718-335-7100 | Fax: 718-709-4136

PATIENT:

LYTON, SAMANTHA

EXAM DATE:

06/25/2022 11:17 AM

STUDY

MRI SHOULDER WITHOUT

MRN:

SAML69619

DESCRIPTION:

CONTRAST

DOB:

08/31/2001

REFERRING PHYSICIAN: Zilberman, Igor DC

CLINICAL HISTORY: NF: PAIN DUE TO ACCIDENT

MAGNETIC RESONANCE IMAGING OF RIGHT SHOULDER WITHOUT CONTRAST

HISTORY: Pain due to accident.

TECHNIQUE: Multiplanar, multi-sequence MRI of the right shoulder was performed without

intravenous contrast.

COMPARISON: None available.

OSSEOUS STRUCTURES/MARROW: Normal marrow signal.

ROTATOR CUFF:

SUPRASPINATUS: The supraspinatus tendon maintains intact tendon fibers. No tendon retraction is found. No skeletal muscle atrophy is seen.

INFRASPINATUS: The infraspinatus tendon maintains intact tendon fibers. No tendon retraction is found. No skeletal muscle atrophy is seen.

TERES MINOR: The teres minor tendon maintains intact tendon fibers. No tendon retraction is found. No skeletal muscle atrophy is seen.

SUBSCAPULARIS: The subscapularis tendon maintains intact tendon fibers. No tendon retraction is found. No skeletal muscle atrophy is seen.

SUBACROMIAL/SUBDELTOID BURSA: No fluid in subacromial-subdeltoid bursa to suggest bursitis.

MUSCLES: No muscle edema or fatty muscle atrophy.

AC JOINT: AC joint hypertrophy may contribute to rotator cuff impingement.



40-34 74th Street, Elmhurst, NY 11373 Tel: 718-335-7100 | Fax: 718-709-4136

PATIENT:

LYTON, SAMANTHA

EXAM DATE:

06/25/2022 11:17 AM

STUDY

MRI SHOULDER WITHOUT

MRN:

SAML69619

DESCRIPTION:

CONTRAST

DOB:

08/31/2001

REFERRING PHYSICIAN: Zilberman, Igor DC

CLINICAL HISTORY: NF: PAIN DUE TO ACCIDENT

BICEPS TENDON: Intact long-head of the biceps tendon.

LABRUM/LIGAMENTS: There is a tear of the superior glenoid labrum.

CORACOACROMIAL LIGAMENT/ROTATOR INTERVAL: Rotator interval is normal.

GLENOHUMERAL CARTILAGE: Intact articular cartilage.

SYNOVIUM/JOINT FLUID: No joint effusion or synovial thickening.

NEUROVASCULAR STRUCTURES: Normal in course and caliber.

PERIPHERAL SOFT TISSUES: Normal.

IMPRESSION:

- 1. Tear of the superior glenoid labrum.
- 2. AC joint hypertrophy may contribute to rotator cuff impingement.

Digitally Signed By: Imam, Naiyer

Digitally Signed Date: 06/27/2022 8:14 PM