

1575 Hiliside Ave, Suite 100 New Hyde Park, NY,11040 Tel: 516 962 9544, 516 962 9599 Cett: 516 549 6963 Fax: 516 467 3130 Email: qrl.newhydepark@yahoo.com

PATIENT:	ISSAINT, WILSON	EXAM DATE:	08/05/2022 6:29 PM
STUDY DESCRIPTION:	MRI SHOULDER WITHOUT CONTRAST	MRN:	ISSW71713
DOB:	06/16/1951	REFERRING PHYSICIAN:	Jurkowich, Michael
CLINICAL HISTORY:	Pain in both shoulders after mva	GENDER:	M

MAGNETIC RESONANCE IMAGING OF RIGHT SHOULDER WITHOUT CONTRAST

HISTORY: Pain in both shoulders after motor vehicle accident.

TECHNIQUE: Multiplanar, multi-sequence MRI of the right shoulder was performed without intravenous contrast.

COMPARISON: None available.

OSSEOUS STRUCTURES/MARROW: Normal marrow signal.

ROTATOR CUFF:

SUPRASPINATUS: The supraspinatus tendon maintains intact tendon fibers. No tendon retraction is found. No skeletal muscle atrophy is seen.

INFRASPINATUS: The infraspinatus tendon maintains intact tendon fibers. No tendon retraction is found. No skeletal muscle atrophy is seen.

TERES MINOR: The teres minor tendon maintains intact tendon fibers. No tendon retraction is found. No skeletal muscle atrophy is seen.

SUBSCAPULARIS: The subscapularis tendon maintains intact tendon fibers. No tendon retraction is found. No skeletal muscle atrophy is seen.

SUBACROMIAL/SUBDELTOID BURSA: Mild fluid in subacromial-subdeltoid bursa compatible with bursitis or may be seen with full thickness rotator cuff tear.

MUSCLES: No muscle edema or fatty muscle atrophy.

AC JOINT: Mild subluxation of the acromioclavicular joint noted with significant hypertrophy of the joint capsule.

BICEPS TENDON: Intact long-head of the biceps tendon.

LABRUM/LIGAMENTS: No labral tear or ligament abnormalities.



1575 Hillside Ave, Suite 100 New Hyde Park, NY 11040 Tel: 516 962 9544, 516 962 9599 Cell: 516 549 6963 Fax: 516 467 3130 Email: qrl.newhydepark@yahoo.com

PATIENT:	ISSAINT, WILSON	EXAM DATE:	08/05/2022 6:29 PM
STUDY DESCRIPTION:	MRI SHOULDER WITHOUT CONTRAST	MRN:	ISSW71713
DOB:	06/16/1951	REFERRING PHYSICIAN:	Jurkowich, Michael
CLINICAL HISTORY:	Pain in both shoulders after mva	GENDER:	М

CORACOACROMIAL LIGAMENT/ROTATOR INTERVAL: Rotator interval is normal.

GLENOHUMERAL CARTILAGE: Intact articular cartilage.

SYNOVIUM/JOINT FLUID: No joint effusion or synovial thickening.

NEUROVASCULAR STRUCTURES: Normal in course and caliber.

PERIPHERAL SOFT TISSUES: Normal.

IMPRESSION:

- 1. Mild subluxation of the acromioclavicular joint with significant hypertrophy of the joint capsule.
- 2. Mild fluid in subacromial-subdeltoid bursa compatible with bursitis or may be seen with full thickness rotator cuff tear.

Digitally Signed By: Imam, Naiyer

Digitally Signed Date: 08/07/2022 12:24 AM