

40-34 74th Street, Elmhurst, NY 11373
Tel: 718-335-7100 | Fax: 718-709-4136

PATIENT:	XINASTLE, REYES	EXAM DATE:	11/08/2022 1:53 PM
STUDY DESCRIPTION:	MRI SHOULDER WITHOUT CONTRAST	MRN:	REYX80874
DOB:	11/01/1981	REFERRING PHYSICIAN:	Levtsenko, Inna Np
CLINICAL HISTORY:	NF: PAIN DUE TO ACCIDENT		

MAGNETIC RESONANCE IMAGING OF LEFT SHOULDER WITHOUT CONTRAST

HISTORY: Pain due to accident.

TECHNIQUE: Multiplanar, multi-sequence MRI of the left shoulder was performed without intravenous contrast.

COMPARISON: None available.

FINDINGS:

OSSEOUS STRUCTURES/MARROW: Normal marrow signal.

ROTATOR CUFF:

SUPRASPINATUS: There is a partial-thickness undersurface tear of the supraspinatus tendon (image 8, series 3).

INFRASPINATUS: The infraspinatus tendon maintains intact tendon fibers. No tendon retraction is found. No skeletal muscle atrophy is seen.


TERES MINOR: The teres minor tendon maintains intact tendon fibers. No tendon retraction is found. No skeletal muscle atrophy is seen.

SUBSCAPULARIS: The subscapularis tendon maintains intact tendon fibers. No tendon retraction is found. No skeletal muscle atrophy is seen.

SUBACROMIAL/SUBDELTOID BURSA: Mild fluid in subacromial-subdeltoid bursa compatible with bursitis or may be seen with full thickness rotator cuff tear.

MUSCLES: No muscle edema or fatty muscle atrophy.

AC JOINT: Intact.



40-34 74th Street, Elmhurst, NY 11373
Tel: 718-335-7100 | Fax: 718-709-4136

PATIENT:	XINASTLE, REYES	EXAM DATE:	11/08/2022 1:53 PM
STUDY DESCRIPTION:	MRI SHOULDER WITHOUT CONTRAST	MRN:	REYX80874
DOB:	11/01/1981	REFERRING PHYSICIAN:	Levtsenko, Inna Np
CLINICAL HISTORY:	NF: PAIN DUE TO ACCIDENT		

BICEPS TENDON: There is tenosynovitis of the extra articular long head of the biceps tendon.
LABRUM/LIGAMENTS: No labral tear or ligament abnormalities.

CORACOACROMIAL LIGAMENT/ROTATOR INTERVAL: Rotator interval is normal.

GLENOHUMERAL CARTILAGE: Intact articular cartilage.

SYNOVIUM/JOINT FLUID: There is no joint effusion

NEUROVASCULAR STRUCTURES: Normal in course and caliber.

PERIPHERAL SOFT TISSUES: Normal.

IMPRESSION:

1. Partial-thickness undersurface tear of the supraspinatus tendon.
2. Mild fluid in subacromial-subdeltoid bursa compatible with bursitis or may be seen with full thickness rotator cuff tear.
3. Tenosynovitis of the extra articular long head of the biceps tendon.

Digitally Signed By: Imam, Naiyer
Digitally Signed Date: 11/09/2022 7:23 AM