



QUEENS RADIOLOGY IMAGING PC

DIAGNOSTIC RADIOLOGY

1575 Hillside Ave, Suite 100
New Hyde Park, NY 11040
Tel: 516 962 9544, 516 962 9599
Cell: 516 549 6963
Fax: 516 467 3130
Email: qrl.newhydepark@yahoo.com

PATIENT:	BARRINGTON, HANNAM	EXAM DATE:	07/26/2022 5:00 PM
STUDY DESCRIPTION:	MRI SHOULDER WITHOUT CONTRAST	MRN:	BARH71617
DOB:	10/26/1967	REFERRING PHYSICIAN:	Jurkovich, Michael
CLINICAL HISTORY:	pain in both shoulders after mva	GENDER:	F

MAGNETIC RESONANCE IMAGING OF LEFT SHOULDER WITHOUT CONTRAST

HISTORY: Pain in both shoulders after motor vehicle accident.

TECHNIQUE: Multiplanar, multi-sequence MRI of the left shoulder was performed without intravenous contrast.

COMPARISON: None available.

FINDINGS:

OSSEOUS STRUCTURES/MARROW: Normal marrow signal.

ROTATOR CUFF:

SUPRASPINATUS: The supraspinatus tendon maintains intact tendon fibers. No tendon retraction is found. No skeletal muscle atrophy is seen.

INFRASPINATUS: The infraspinatus tendon maintains intact tendon fibers. No tendon retraction is found. No skeletal muscle atrophy is seen.


TERES MINOR: The teres minor tendon maintains intact tendon fibers. No tendon retraction is found. No skeletal muscle atrophy is seen.

SUBSCAPULARIS: The subscapularis tendon maintains intact tendon fibers. No tendon retraction is found. No skeletal muscle atrophy is seen.

SUBACROMIAL/SUBDELTOID BURSA: Mild fluid in subacromial-subdeltoid bursa compatible with bursitis or may be seen with full thickness rotator cuff tear.

MUSCLES: No muscle edema or fatty muscle atrophy.

AC JOINT: Mild acromioclavicular joint disease with capsular thickening and small marginal osteophytes. Lateral downsloping of the acromion with subacromial spurring. These factors contribute to rotator cuff impingement.



**QUEENS RADIOLOGY
IMAGING PC**
DIAGNOSTIC RADIOLOGY

1575 Hillside Ave, Suite 100
New Hyde Park, NY 11040
Tel: 516 962 9544, 516 962 9599
Cell: 516 549 6963
Fax: 516 467 3130
Email: qri.newhydepark@yahoo.com

PATIENT:	BARRINGTON, HANNAM	EXAM DATE:	07/26/2022 5:00 PM
STUDY DESCRIPTION:	MRI SHOULDER WITHOUT CONTRAST	MRN:	BARH71617
DOB:	10/26/1967	REFERRING PHYSICIAN:	Jurkovich, Michael
CLINICAL HISTORY:	pain in both shoulders after mva	GENDER:	F

BICEPS TENDON: There is tenosynovitis of the extra articular long head of the biceps tendon.

LABRUM/LIGAMENTS: No labral tear or ligament abnormalities.

CORACOACROMIAL LIGAMENT/ROTATOR INTERVAL: Rotator interval is normal.

GLENOHUMERAL CARTILAGE: Intact articular cartilage.

SYNOVIUM/JOINT FLUID: No joint effusion or synovial thickening.

NEUROVASCULAR STRUCTURES: Normal in course and caliber.

PERIPHERAL SOFT TISSUES: Normal.

IMPRESSION:

1. Mild fluid in subacromial-subdeltoid bursa compatible with bursitis or may be seen with full thickness rotator cuff tear.
2. Mild acromioclavicular joint disease contributing to rotator cuff impingement.
3. Tenosynovitis of the extra articular long head of the biceps tendon.

Digitally Signed By: Imam, Naiyer

Digitally Signed Date: 07/28/2022 10:38 AM