

**AMI****American Medical Initiatives**

30-80 31st Street, Astoria, NY 11102  
Tel: 718-335-7100 | Fax: 718-709-4136

<b>PATIENT:</b>	DERBY, ISSA	<b>EXAM DATE:</b>	05/04/2022 4:22 PM
<b>STUDY DESCRIPTION:</b>	MRI SHOULDER WITHOUT CONTRAST	<b>MRN:</b>	DERI64895
<b>DOB:</b>	11/06/1961	<b>REFERRING PHYSICIAN:</b>	Qureshi, Adnan
<b>CLINICAL HISTORY</b>	C/O RT SHOULDER PAIN DUE TO MVA	<b>GENDER</b>	F

**MAGNETIC RESONANCE IMAGING OF RIGHT SHOULDER WITHOUT CONTRAST**

**HISTORY:** Complains of right shoulder pain due to motor vehicle accident.

**TECHNIQUE:** Multiplanar, multi-sequence MRI of the right shoulder was performed without intravenous contrast.

**COMPARISON:** None available.

**OSSEOUS STRUCTURES/MARROW:** Normal marrow signal.

**ROTATOR CUFF:**

**SUPRASPINATUS:** There is a focal non-retracted interstitial tear at the attachment of the supraspinatus, well appreciated on image 11 of the coronal series. No tendon retraction is found. No skeletal muscle atrophy is seen.

**INFRASPINATUS:** The infraspinatus tendon maintains intact tendon fibers. No tendon retraction is found. No skeletal muscle atrophy is seen.

**TERES MINOR:** The teres minor tendon maintains intact tendon fibers. No tendon retraction is found. No skeletal muscle atrophy is seen.

**SUBSCAPULARIS:** The subscapularis tendon maintains intact tendon fibers. No tendon retraction is found. No skeletal muscle atrophy is seen.

**SUBACROMIAL/SUBDELTOID BURSA:** Mild fluid in subacromial-subdeltoid bursa compatible with bursitis or may be seen with full thickness rotator cuff tear.

**AMI****American Medical Initiatives**

30-80 31st Street, Astoria, NY 11102  
Tel: 718-335-7100 | Fax: 718-709-4136

<b>PATIENT:</b>	DERBY, ISSA	<b>EXAM DATE:</b>	05/04/2022 4:22 PM
<b>STUDY DESCRIPTION:</b>	MRI SHOULDER WITHOUT CONTRAST	<b>MRN:</b>	DERI64895
<b>DOB:</b>	11/06/1961	<b>REFERRING PHYSICIAN:</b>	Qureshi, Adnan
<b>CLINICAL HISTORY</b>	C/O RT SHOULDER PAIN DUE TO MVA	<b>GENDER</b>	F

**MUSCLES:** No muscle edema or fatty muscle atrophy.

**AC JOINT:** Intact.

**BICEPS TENDON:** Intact long-head of the biceps tendon.

**LABRUM/LIGAMENTS:** There is a tear of the superior glenoid labrum anterior to posterior (SLAP tear) with a partial tear of the biceps labral anchor complex.

**CORACOACROMIAL LIGAMENT/ROTATOR INTERVAL:** Rotator interval is normal.

**GLENOHUMERAL CARTILAGE:** Intact articular cartilage.

**SYNOVIUM/JOINT FLUID:** No joint effusion or synovial thickening.

**NEUROVASCULAR STRUCTURES:** Normal in course and caliber.

**PERIPHERAL SOFT TISSUES:** Normal.

**IMPRESSION:**

1. Focal non-retracted interstitial tear at the attachment of the supraspinatus.
2. Type II SLAP tear.
3. Mild fluid in subacromial-subdeltoid bursa compatible with bursitis or may be seen with full thickness rotator cuff tear.

**AMI****American Medical Initiatives**

30-80 31st Street, Astoria, NY 11102  
Tel: 718-335-7100 | Fax: 718-709-4136

<b>PATIENT:</b>	DERBY, ISSA	<b>EXAM DATE:</b>	05/04/2022 4:22 PM
<b>STUDY DESCRIPTION:</b>	MRI SHOULDER WITHOUT CONTRAST	<b>MRN:</b>	DERI64895
<b>DOB:</b>	11/06/1961	<b>REFERRING PHYSICIAN:</b>	Qureshi, Adnan
<b>CLINICAL HISTORY</b>	C/O RT SHOULDER PAIN DUE TO MVA	<b>GENDER</b>	F

Digitally Signed By: Imam, Naiyer  
Digitally Signed Date: 05/05/2022 4:33 AM