eMedicalOfficeSolutions

Procedure Reading

Bronx Medical Diagnostic, P.C.

3117 Buhre Avenue

Phone; (718) 764-5000

Fax: (718) 239-6308

Bronx, NY 10461 PATIENT NAME:

Francisco De La Rosa Marte

DOB:

3/9/1973

DATE OF SERVICE:

1/7/2021

REFERRING DOCTOR: Adnan A. Qureshi, M.D.

MRI Left Shoulder:

TECHNIQUE: Magnetic Resonance Imaging is Performed in Multiple Projections Utilizing T1/T2 Pulse Sequences.

The supraspinatus tendon hyperintensity at its greater tuberosity insertion suggests tendinitis. Possible partial tear component cannot be excluded. Frank rupture with muscular retraction has not developed.

The visualized glenoid labrum outlines anatomically. Biceps tendon is free from interruption. Some fluid extends along joint capsule and subdettold bursa. Osseous structures maintain unremarkable alignment and signal characteristics. Suspicion is not raised for fracture, subluxation or replacement process. Smooth humeral head. Acromicolavicular joint remains intact. Coraccacromial arch is without appreciable encroachment. No penarticular mass is declared. Axilla appears clear of lymphadenopathy.

<u>IMPRESSION:</u>

- 1. Evidence for retator cuff tendinopathy.
- 2. Associated bursitis.
- 3. No discernible bony deformity.

Thank you for the courtesy of this consultation.

Mark Lodespoto, M.D.

Diplomat, American Board of Radiology

C.A.Q. Neuroradiology