

30-80 31st Street, Astoria, NY 11102 Tel: 718-335-7100 | Fax: 718-709-4135

PATIENT:	FRAGOSA, JUDITH	EXAM DATE: 09/21/2022 2:52 PM
SYUDY DESCRIPTION:	MRI SHOULDER WITHOUT CONTRAST	MEUS: 1. FRA.7/1543
DOB:	11/20/1987	REFERRING Koutsospyros, Demetilios A
CLINICAL BISTORY	C/O LT SHOULDER PAIN QUE TO MVA	GENDER F

MAGNETIC RESONANCE IMAGING OF LEFT SHOULDER WITHOUT CONTRAST

HISTORY: Complains of left shoulder pain due to motor vehicle accident.

TECHNIQUE: Multiplanar, multi-sequence MRI of the left shoulder was performed without intravenous contrast.

COMPARISON: None available.

FINDINGS:

Frem ASTORIA- American Medical Initia

OSSEOUS STRUCTURES/MARROW: Normal marrow signal.

ROTATOR CUFF:

SUPRASPINATUS: Interstitial tear of length 10mm at the attachment of the supraspinatus tendon. INFRASPINATUS: The infraspinatus tendon maintains intact tendon fibers. No tendon retraction is found. No skeletal muscle atrophy is seen.

TERES MINOR: The teres minor tendon maintains intact tendon fibers. No tendon retraction is found. No skeletal muscle atrophy is seen.

SUBSCAPULARIS: The subscapularis tendon maintains intact tendon fit. 16. No tendon retraction is found. No skeletal muscle atrophy is seen.

SUBACROMIAL/SUBDELTOID BURSA: There is a minimal subcoracoid huid.

MUSCLES: No muscle edema or fatty muscle atrophy.

ACJOINT: ACjoint arthroisis.

BICEPS TENDON: Intact long-head of the bice ps tendon.



30-80 31st Street, Astoria, NY 11102 Tel: 718-335-7100 | Fax: 718-709-4136

MATENT	FRAGOSA, JUDITH	EXAM DATE:	0 %21/ 2022 2:52 PM
STUDY DESCRIPTION:	MRI SHOULDER WITHOUT CONTRAST	The state of the s	FRAJ71543
DOa:	11/20/1987	REFERRING PHYSICIAN	Koutsospyros, Demetrios A
CLINICAL HISTORY	C/O LT SHOULDER PAIN DUE TO MVA	GENDER	F

LABRUM/LIGAMENTS: No labral tear or ligament abnormalities.

CORACGACROMIAL LIGAMENT/ROTATOR INTERVAL: Rotator interval is normal.

GLENOHUMERAL CARTILAGE: Intact articular cartilage.

SYNOVIUM/JOINT FLUID: No joint effusion or synovial thickening.

NEUROVASCULAR STRUCTURES: Normal in course and caliber

PERIPHERAL SOFT TISSUES: Normal.

IMPRESSION:

- 1. Interstitial tear of length 10mm at the attachment of the supraspinatus tendon,
- 2. There is a minimal subcoracoid fluid.
- 3. AC joint arthroisis.

Digitally Signed By: Imam, Noiyar

Digitally Signed Date: 09/22/2022 10:28 AM

