



30-80 31st Street, Astoria, NY 11102
Tel: 718-335-7100 | Fax: 718-709-4136

PATIENT:	VICTOR, JORGE	EXAM DATE:	08/06/2022 10:38 AM
STUDY DESCRIPTION:	MRI ELBOW WITHOUT CONTRAST	MRN:	VICJ70995
DOB:	01/03/1979	REFERRING PHYSICIAN:	Qureshi, Adnan
CLINICAL HISTORY	C/O LT ELBOW PAIN DUE TO MVA	GENDER	M

MAGNETIC RESONANCE IMAGING OF LEFT ELBOW WITHOUT CONTRAST

HISTORY: Complains of left elbow pain due to motor vehicle accident.

TECHNIQUE: Multiplanar, multi-sequence MRI of left elbow was performed without intravenous contrast.

COMPARISON: None available.

FINDINGS:

OSSEOUS STRUCTURES/MARROW: No fractures or osteonecrosis.

OSSEOUS ALIGNMENT: Anatomic alignment.

ANTERIOR MUSCLES/TENDONS:

BICEPS MUSCLE/TENDON: The distal biceps muscle and tendon are intact.

BRACHIALIS MUSCLE/TENDON: The distal brachialis muscle and tendon are intact.

POSTERIOR MUSCLES/TENDON:

TRICEPS MUSCLE/TENDON: Linear interstitial tearing of the triceps tendon noted at the level of its insertion.

LATERAL ELBOW LIGAMENTS AND TENDONS:

RADIAL COLLATERAL LIGAMENT: The radial collateral ligament is intact.

LATERAL ULNAR COLLATERAL LIGAMENT: The lateral ulnar collateral ligament is intact.

COMMON EXTENSOR TENDON: The common extensor ligament is intact.

MEDIAL ELBOW LIGAMENTS AND TENDONS:



30-80 31st Street, Astoria, NY 11102
Tel: 718-335-7100 | Fax: 718-709-4136

PATIENT:	VICTOR, JORGE	EXAM DATE:	08/06/2022 10:38 AM
STUDY DESCRIPTION:	MRI ELBOW WITHOUT CONTRAST	MRN:	VICJ70995
DOB:	01/03/1979	REFERRING PHYSICIAN:	Qureshi, Adnan
CLINICAL HISTORY	C/O LT ELBOW PAIN DUE TO MVA	GENDER	M

ULNAR COLLATERAL LIGAMENT: The ulnar collateral ligament is intact.

COMMON FLEXOR TENDON: Hyperintense signal noted at the origin of the common flexor tendon compatible with partial tear.

CARTILAGE/SUBCHONDRAL BONE: Cartilage and subchondral bone intact.

SYNOVIUM /JOINT FLUID: No synovial thickening or joint effusion.

NEUROVASCULAR STRUCTURES: Normal in course and caliber.

PERIPHERAL SOFT TISSUES: Normal.

IMPRESSION:

1. Linear interstitial tearing of the triceps tendon at the level of its insertion.
2. Hyperintense signal at the origin of the common flexor tendon compatible with partial tear.

Digitally Signed By: Imam, Naiyer
Digitally Signed Date: 08/09/2022 4:26 AM

