



39

Next Generation Diagnostic Imaging P.C.

MERSHAD HAGIGI, MD, PHD
DIPLOMATE, AMERICAN BOARD OF RADIOLOGY
1664 East 14th Street, Suite LL • Brooklyn, NY 11229
TEL: (718) 336-1865 • FAX: (718) 336-1275

DATE OF STUDY: 09-SEP-2022 01:31:48 PM
PATIENT NAME: BARTLEY, WAYNE
DATE OF BIRTH: 25-JAN-1964
PATIENT NUMBER: SR3138
REFERRING PHYSICIAN: DR.CARLOTTE ROSS

MRI OF THE LEFT SHOULDER WITHOUT CONTRAST

CLINICAL HISTORY: Status post MVA.

TECHNIQUE: Multisequential multiplanar imaging of the left shoulder was performed. No contrast was administered for this exam.

FINDINGS:

Bones/Joints: There is mild fibrocystic change within the greater tuberosity. There is no fracture or contusion pattern. The glenohumeral joint space is preserved. The acromioclavicular joint space is maintained. There is a type II acromion. There is mild impingement at the level of the acromial tip. This is seen on image 25 of the sagittal T2 series.

Rotator Cuff: There is mild diffuse rotator cuff tendinosis. There is a linear undersurface tearing within the distal fibers of the mid to anterior supraspinatus tendon. This is seen on images 6 and 7 of the coronal STIR series. There is no rotator cuff muscle atrophy.

Labrum and Biceps Anchor: The glenoid labrum is intact. The long head of the biceps tendon and biceps anchor are intact.

IMPRESSION:

1. Mild diffuse rotator cuff tendinosis. Linear undersurface tearing within the distal fibers of the mid to anterior supraspinatus tendon.
2. A type II acromion. Mild impingement at the level of the acromial tip.

Thank you for the courtesy of this referral.

Electronically Signed by LERER, DANIEL BRIAN at 9/11/2022 4:41:47 PM GMT-04:00