

170-10 Cedarcroft Road, Jamaica, NY 11432
Tel: 718-206-1000 | Fax: 718-532-0633

PATIENT:	BASTOS, LUIS	EXAM DATE:	06/02/2022 3:15 PM
STUDY DESCRIPTION:	MRI KNEE WITHOUT CONTRAST (JOINT)	MRN:	BASL69576
DOB:	12/23/1952	REFERRING PHYSICIAN:	Qureshi, Adnan
CLINICAL HISTORY:	C/O OF LEFT KNEE PAIN DUE TO MVA	GENDER	M

MAGNETIC RESONANCE IMAGING OF THE LEFT KNEE WITHOUT IV CONTRAST

HISTORY: Complains of left knee pain due to motor vehicle accident.

TECHNIQUE: Multiplanar, multi-sequence MRI of the left knee was performed without intravenous contrast.

COMPARISON: None available.

FINDINGS:

OSSEOUS STRUCTURES/MARROW: A bone contusion of the lateral femoral condyle noted. No fractures or osteonecrosis.

LIGAMENTS:

ANTERIOR CRUCIATE: The anterior cruciate ligament is intact.

POSTERIOR CRUCIATE: The posterior cruciate ligament is intact.

MEDIAL COLLATERAL LIGAMENT: The medial collateral ligament is intact.

LATERAL COLLATERAL LIGAMENT: The lateral collateral ligament is intact.


JOINT SPACES:

MEDIAL COMPARTMENT: A horizontal tear of the body and posterior horn of the medial meniscus noted.

LATERAL COMPARTMENT: Intact lateral meniscus and articular cartilage.

PATELLOFEMORAL COMPARTMENT: Articular cartilage intact.

SYNOVIUM/ JOINT FLUID: There is no joint effusion.



170-10 Cedarcroft Road, Jamaica, NY 11432
Tel: 718-206-1000 | Fax: 718-532-0633

PATIENT:	BASTOS, LUIS	EXAM DATE:	06/02/2022 3:15 PM
STUDY DESCRIPTION:	MRI KNEE WITHOUT CONTRAST (JOINT)	MRN:	BASL69576
DOB:	12/23/1952	REFERRING PHYSICIAN:	Qureshi, Adnan
CLINICAL HISTORY:	C/O OF LEFT KNEE PAIN DUE TO MVA	GENDER:	M

MUSCLES: No muscle edema or fatty muscle atrophy.

NEUROVASCULAR STRUCTURES: Normal in course and caliber.

EXTENSOR MECHANISM: The quadriceps tendon is intact. The patellar tendon is intact.

PERIPHERAL SOFT TISSUES: Significant edema noted in the prepatellar region compatible with trauma sequelae.

PLICAE: No plicae demonstrated.

IMPRESSION:

1. Horizontal tear of the body and posterior horn of the medial meniscus.
2. Bone contusion of the lateral femoral condyle.
3. Significant edema in the prepatellar region compatible with trauma sequelae.

Digitally Signed By: Imam, Naiyer
Digitally Signed Date: 06/03/2022 7:45 PM

