

Westchester Rad ology & Imaging, PC

933 Saw Nick River Road Ardsley NY 10502

Phone: 914-740-1: 38 Fax: 914-478-0303

PATIENT:

HAYNES TIARA

DOB:

02/17/1993

PHYSICIAN:

DR. FERSEL

EXAM DATE:

04/07/2022

MRI OF THE LEFT SHOULDER

INDICATION: Pain.

TECHNIQUE: Multiple T1 and T2 weighted MRI images of the left shoulder were obtained in the axial, sagittal and coronal planes without intravenous or intraacticular contrast.

FINDINGS: There are no acute displaced fractures, dislocations, destructive bony lesions or marrow infiltration in the proximal humerus and glenoid.

The rotator cuff musculature including the supraspinatus, subscapularis, infraspinatus and teres minor are normal in bulk without atrophy, edema or fatty infiltration. The rotator cuff tendons including the subscapularis, infraspinatus and teres minor are intact without MRI evidence of a tear or tendinosis/tendinopathy. The glenoid labrum is grossly intact. There are no masses associated with the glenohumeral joint.

There is a partial tear of the distal supraspinatus tendon. Fluid in the long head of the biceps tendon sheath consistent with tenosynovitis. Low lying acromion with impingement of rotator cuff, in an appropriate clinical setting. Edema in the distal clavick and adjacent acromion with fluid in the acromioclavicular joint, consistent with recent trauma. Fluid in the subacromial/subdeltoid bursa suggestive of underlying rotator cuff tears and/or subacromial/subdeltoid bursitis, in an appropriate clinical setting. Mild joint effusion consistent with recent trauma or synovitis, in an appropriate clinical setting.

IMPRESSION:

- 1. Partial tear of the distal supraspinatus tendon.
- 2. Fluid in the long head of the biceps tendon sheath consistent with tenosynovitis.
- 3. Low lying acromion with impingement of rotator cuff, in an appropriate clinical setting.
- Edema in the distal clavicle and adjacent acromion with fluid in the acromioclavicular joint, consistent with recent trauma.

PATIENT:

HAYNES TIARA

DOB:

02/17/1993

PHYSICIAN:

DR. FERSEL

EXAM DATE:

04/07/2022

Fluid in the subacromial/subdeltoid bursa suggestive of underlying rotator cuff tears and/or subacromial/subdeltoid bursitis, in an appropriate clinical setting.

 Mild joint effusion consistent with recent trauma or synovitis, in an appropriate clinical setting.

Steve B. Losik M.D.

Steve B. Losik, M.D. Board Certified Radiologist Electronically Signed