UK Sinha Physician, P.C.

102-31 Jamaica Ave. Richmond Hill, NY 11418 Ph: 718-480-1130 Fax: 718-480-1132

October 26, 2022

Office seen at: Chiro 4226 4226-A 3rd Ave Bronx, NY 10457 Phone# (718) 684-7676

Re: Moise, Lenny DOB: 09/04/1982 DOA: 12/31/2021

FOLLOW-UP NOTE

CHIEF COMPLAINT: Follow up of right shoulder and left shoulder pain.

HISTORY OF PRESENT ILLNESS: The patient presents today in followup with continued pain in the right shoulder and to evaluate pain in the left shoulder.

ADL CAPABILITIES: As a direct result of the injuries sustained in this accident, the patient states that he is unable to do the following activities: driving, lifting heavy objects, carrying, reaching overhead, laundry, shopping, running errands, and exercising.

PRESENT COMPLAINTS: Right shoulder: Right shoulder pain is 10/10, described as constant, sharp, stabbing, dull, achy pain. The patient has stiffness, weakness, popping, and clicking. The patient is able to reach overhead and able to reach behind the back but is unable to sleep at night due to pain. Worse with range of motion and improves with physical therapy.

Left shoulder: Left shoulder pain is 6/10, described as intermittent, sharp, stabbing, dull, achy pain. The patient has stiffness, weakness, popping, and clicking. The patient is able to reach overhead and able to reach behind the back, but is unable to sleep at night due to pain. Worse with range of motion and improves with rest.

PHYSICAL EXAMINATION: The right shoulder reveals tenderness to palpation over supraspinatus tendon region and AC joint. There is crepitus appreciated. There is no heat, swelling, erythema or deformity appreciated. Negative drop arm test. Positive cross-over test. Positive empty can test. Positive Yergason test. Negative deltoid atrophy. Negative O'Brien test. Positive impingement sign. Positive Lift-off test. Positive Hawkins test. Range of motion, as per goniometer, abduction 110/180 degrees, adduction 40/45 degrees, forward flexion 125/180 degrees, extension 45/60 degrees, internal rotation 45/90 degrees, and

Moise, Lenny October 26, 2022 Page 2 of 3

external rotation 55/90 degrees. Internal rotation to the sacrum. The patient has no motor or sensory deficit of the right upper extremity.

The left shoulder reveals tenderness to palpation over supraspinatus tendon region and proximal biceps tendon. There is crepitus appreciated. There is no heat, swelling, erythema or deformity appreciated. Negative drop arm test. Positive cross-over test. Positive empty can test. Positive Yergason test. Negative deltoid atrophy. Negative O'Brien test. Positive impingement sign. Positive Lift-off test. Positive Hawkins test. Range of motion, as per goniometer, abduction 115/180 degrees, adduction 40/45 degrees, forward flexion 130/180 degrees, extension 45/60 degrees, internal rotation 45/90 degrees, and external rotation 60/90 degrees. Internal rotation to the sacrum. The patient has no motor or sensory deficit of the left upper extremity.

DIAGNOSTIC TESTING: MRI of the right shoulder, done on 04/29/2022, shows intact right shoulder; osteophytes at the acromioclavicular joint. Tendinopathy and partial tear of the supraspinatus tendon. Tendinopathy of the infraspinatus tendon. Intact glenohumeral joint with effusion. Biceps tendinopathy. MRI of the left shoulder, done on 04/22/2022, shows effusion at the glenohumeral joint. Partial tear of the anterior leading edge of the supraspinatus tendon with tendinopathy.

ASSESSMENT:

- 1. S46.011A Partial rotator cuff tear, right shoulder.
- 2. M24.811 Internal derangement, right shoulder.
- 3. M75.81 Shoulder tendinitis, right shoulder.
- 4. M75.41 Impingement, right shoulder.
- 5. M75.21 Bicipital tendinitis, right shoulder.
- 6. M25.511 Pain, right shoulder.
- 7. S49.91XA Injury, right shoulder.
- 8. M25.411 Joint effusion, right shoulder.
- 9. S46.012A Partial rotator cuff tear, left shoulder.
- 10. M24.812 Internal derangement, left shoulder.
- 11. M75.82 Shoulder tendinitis, left shoulder.
- 12. M75.42 Impingement, left shoulder.
- 13. M25.512 Pain, left shoulder.
- 14. S49.92XA Injury, left shoulder.
- 15. M25.412 Joint effusion, left shoulder.

PLAN:

- 1. Imaging studies and clinical examinations were reviewed with the patient.
- 2. All treatment options discussed with the patient.
- 3. Cold compresses for right shoulder and left shoulder.
- 4. Continue anti-inflammatory and muscle relaxant medications p.r.n.
- 5. Continue physical therapy for right shoulder and left shoulder 3 days/week.
- 6. Recommend steroid injections with pain management for right shoulder and left shoulder. The patient refuses due to side effects.

- 7. Discussed right shoulder and left shoulder arthroscopy versus conservative management with the patient. The patient states that due to continual pain and lack of relief with physical therapy and the inability to perform day-to-day activities due to pain, the patient would like to proceed with surgery.
- 8. The patient needs medical clearance prior to surgery.
- 9. It is medically necessary to perform the suggested surgery to properly diagnose the patient's condition and to objectively verify presence and severity of internal derangement and other right shoulder and left shoulder pathology in quantitative and qualitative terms and achieve better prognosis. This surgery is crucial to provide most specific and maximally effective treatment to this patient.
- 10. Discussed the length of the arthroscopy, the postoperative instructions, and the option of continuing with conservative management alternatives to surgery, including no surgery.
- 11. All the benefits and risks of the right shoulder and left shoulder arthroscopy have been discussed with the patient. The risks include, but not limited to bleeding, infection, pain, stiffness, muscle injury, nerve injury, DVT, and recurrence.
- 12. All the questions in regard to the procedure were answered.
- 13. The patient verbally consents for the arthroscopy of right shoulder and the patient will be scheduled for right shoulder surgery. On the day of the surgery, before the procedure, the surgeon will explain all related questions of the procedure. The patient will sign a written consent and be witnessed by the surgeon.
- 14. The patient will follow up 1-2 weeks postop or in 2 months if surgery has not been performed yet.

<u>CAUSALITY</u>: It is within a certain degree of medical certainty, that the history presented by the patient, the objective physical findings as well as the diagnosis rendered is causally related to the injury the patient incurred on the specified date. These current symptoms were nonexistent prior to the accident. Findings were discussed with the patient.

AFFIRMATION: Being duly licensed to practice medicine in the state of New York, pursuant to the applicable provisions of the Civil Practice Laws and Rules, I hereby affirm under penalty of perjury that the statements contained herein are true and accurate.

Mellita Shakhmurov, PA-C

U.K. Sinha, MD, MS (Ortho), FAAOS Board Certified Orthopedic Surgeon

MS/AEI