

SURGICARE OF BROOKLYN

313 43rd St.

Brooklyn, NY 11232

Phone: (201) 549-9998 – Fax: (646) 585-4468

Right Wrist Arthroscopy Operative Report

Patient Name: Givens, Jeanine

Medical Record Number: 10022

Date of Birth: 05/23/1964

Date of Procedure: 10/13/2022

Surgeon: Upendra K. Sinha, MD.

Assistant: Gennadiy Shamalov, P.A.

Preoperative Diagnoses: de Quervain's disease, right wrist.
Post-traumatic synovitis, rule out chondral damage, right wrist.

Postoperative Diagnoses: M65.4 de Quervain's disease, right wrist.
M65.849 Hypertrophic synovitis, right wrist.
M24.139 Chondral damage, scaphoid, radius.
S63.392A Partial tear scapholunate ligament, right wrist.
S63.591A Triangular fibrocartilage partial tear, right wrist.

Operative Procedure: 29846 Arthroscopic debridement.
25118 Complete synovectomy.
25332 Arthroplasty, radius.
29999 Coblation arthroplasty, scaphoid.

Anesthesia: Regional with IV sedation.

Estimated Blood Loss: Minimal.

Complications: None.

Instrumentation: None.

Intraoperative Findings:

Post traumatic synovitis.
Chondral damage, scaphoid and radius.
Partial tear of scapholunate ligament.
Partial tear of triangular fibrocartilage.

Indications for Surgery:

Indications: The patient has been complaining of chronic pain in the right wrist. The patient failed conservative treatment. All the treatment options were discussed with the patient. The advantages and disadvantages, complications and recovery from surgery was discussed. The possibility of infection, complication from anesthesia, possibility of neurovascular structure injury, blood clots after surgery, pulmonary embolism and need for further surgery were discussed. No guarantees were given.

The patient expressed an understanding of the risks and possible benefits of the procedure and was also made aware of the alternatives to surgery. An informed consent was obtained, and was checked immediately preop.

Description of Procedure:

The patient was brought to the operating room and placed supine on the table. Under regional anesthesia and IV sedation, the hand was suspended vertically with a traction device. Adequate traction was applied. Then after usual aseptic draping and prepping, a standard dorsal 3/4 viewing portal was created and a 4/5 working portal was established. Around 10 mL of saline was used to distend the joint. An 18-gauge needle was used to confirm the outflow of fluid.

The 3/4 portal was made in line with the radial border of the third metacarpal and the 4/5 portal was created in line with the mid axis of the fourth metacarpal. The 6R portal was not used. A 4/5 portal was made just proximal to the 3/4 portal because of the natural radial slope of the distal radius.


A small incision was made in the skin with a #11 blade at the intended portal site. A blunt dissection was carried out with a small curved hemostat. The joint capsule was perforated. Then the arthroscope with a blunt trocar was introduced into the wrist joint. A 4/5 portal was made as usual under direct visualization. the joint was irrigated.

Diagnostic arthroscopic examination was done. Examination revealed extensive synovitis of the wrist. The scapholunate ligament had a tear. The patient had chondral damage of the radius and the scaphoid. The mid carpal joint was normal. Total synovectomy and debridement was performed up to a stable border. Copious irrigation was done followed by closure of the wound. Sterile dressing was applied.

The patient tolerated the procedure well. A post op volar splint was applied. The patient was awakened from anesthesia and brought to the recovery room in a satisfactory condition.

Physician Assistant:

Throughout the procedure, I was assisted by a physician assistant, licensed in the State of New York. He assisted in positioning the patient on the operating room table as well as transferring the patient from the operating room table to the recovery room stretcher. He assisted me during the actual procedure with positioning of the patient's extremity to allow for ease of arthroscopic access to all areas of the joint. The presence of physician assistant as my operating assistant was medically necessary to ensure the utmost safety of the patient in the operative, interim and postoperative period.



U.K. Sinha, MD, MS (Ortho), FAAOS
Board Certified Orthopedic Surgeon