MI

COMMUNITY MEDICAL IMAGING

OF BROOKLYN

ACR Accredited Facility

2102 Ave Z . Brooklyn, NY 11235 . Tel: 718-455-4444 . Fax: 718-615-2121

UPENDRA SINHA, M.D. 1314 CONEY ISLAND AVENUE BROOKLYN, NY 11230

PATIENT: RICARDO FELISME

DOB: 09/30/1980 DOS: 01/20/2022 CHART #: 24405

EXAM: MRI OF THE RIGHT SHOULDER WITHOUT CONTRAST

HISTORY: Pain.

TECHNIQUE: Multiplanar MR imaging of the right shoulder was performed without contrast on Hitachi open MRI unit.

Coronal T1, T2 and STIR; Sagittal T2; Axial PD and gradient echo images of the shoulder were obtained.

COMPARISON: None.

Additional sanitizing / safety protocols recommended by the CDC were performed.

FINDINGS: There is mild enlargement of the supraspinatus and infraspinatus tendons. Generalized heterogeneous T2 signal is seen at the anterior margin of the supraspinatus tendon. Interstitial tearing is noted seen on image #9 of series #5.

The subscapularis and teres minor tendons are intact.

There is a type I acromion. There is no impingement. The musculotendinous junction is unremarkable. There is no muscular injury.


There is no fracture. There is no bone bruise. There is no osteochondral defect.

The glenoid is unremarkable. There is no subluxation.

The anterior and posterior labra are intact. The superior labrum and biceps anchor are unremarkable.

There is no biceps tendon tear or tenosynovitis. The transverse humeral ligament is unremarkable.

There is no muscular injury. There is no hematoma or seroma.



COMMUNITY MEDICAL IMAGING

OF BROOKLYN

ACR Accredited Facility

2102 Ave Z • Brooklyn, NY 11235 • Tel: 718-455-4444 • Fax: 718-615-2121

PATIENT: RICARDO FELISME

DOB: 09/30/1980 DOS: 01/20/2022 CHART #: 24405

EXAM: MRI OF THE RIGHT SHOULDER WITHOUT CONTRAST

PAGE 2

The upper humerus is unremarkable.

There is no AC separation.

Limited visualization of the chest wall is unremarkable.

IMPRESSION:

1. THERE IS GENERALIZED INCREASED T2 SIGNAL SEEN AT THE MID SUBSTANCE OF THE ANTERIOR MARGIN OF THE SUPRASPINATUS TENDON. AN INTERSTITIAL TEAR IS NOTED.

2. TENDINITIS CHANGES ARE SEEN AT THE SUPRASPINATUS AND INFRASPINATUS TENDONS.

Thank you for referring this patient to us.

Andrew McDonnell, MD

Neuroradiologist

Diplomate, American Board of Radiology

AM/man/pr D: 01/20/2022

Edan & Mad similar

E-Sig By A. McDonnell, MD on 01/21/2022 06:54:04