



30-80 31st Street, Astoria, NY 11102
Tel: 718-335-7100 | Fax: 718-709-4136

PATIENT:	QUEZADA, GUSTAVO	EXAM DATE:	03/05/2022 9:56 AM
STUDY DESCRIPTION:	MRI SHOULDER WITHOUT CONTRAST	MRN:	QUEG58251
DOB:	05/11/1979	REFERRING PHYSICIAN:	Qureshi, Adnan
CLINICAL HISTORY	LT SHOULDER PAIN DUE TO MVA	GENDER	M

MAGNETIC RESONANCE IMAGING OF LEFT SHOULDER WITHOUT CONTRAST

HISTORY: Left shoulder pain due to motor vehicle accident.

TECHNIQUE: Multiplanar, multi-sequence MRI of the left shoulder was performed without intravenous contrast.

COMPARISON: None available.

OSSEOUS STRUCTURES/MARROW: Normal marrow signal.

ROTATOR CUFF:

SUPRASPINATUS: The supraspinatus tendon maintains intact tendon fibers. No tendon retraction is found. No skeletal muscle atrophy is seen.

INFRASPINATUS: The infraspinatus tendon maintains intact tendon fibers. No tendon retraction is found. No skeletal muscle atrophy is seen.

TERES MINOR: The teres minor tendon maintains intact tendon fibers. No tendon retraction is found. No skeletal muscle atrophy is seen.

SUBSCAPULARIS: The subscapularis tendon maintains intact tendon fibers. No tendon retraction is found. No skeletal muscle atrophy is seen.

SUBACROMIAL/SUBDELTOID BURSA: No fluid in subacromial-subdeltoid bursa to suggest bursitis.

MUSCLES: No muscle edema or fatty muscle atrophy.

AC JOINT: Severe acromioclavicular joint disease with capsular thickening, small marginal osteophytes and hyperintense articular margins.

BICEPS TENDON: Intact long-head of the biceps tendon.



30-80 31st Street, Astoria, NY 11102
 Tel: 718-335-7100 | Fax: 718-709-4136

PATIENT:	QUEZADA, GUSTAVO	EXAM DATE:	03/05/2022 9:56 AM
STUDY DESCRIPTION:	MRI SHOULDER WITHOUT CONTRAST	MRN:	QUEG58251
DOB:	05/11/1979	REFERRING PHYSICIAN:	Qureshi, Adnan
CLINICAL HISTORY	LT SHOULDER PAIN DUE TO MVA	GENDER	M

LABRUM/LIGAMENTS: There is a tear of the superior glenoid labrum.

CORACOACROMIAL LIGAMENT/ROTATOR INTERVAL: Rotator Interval is normal.

GLENOHUMERAL CARTILAGE: Intact articular cartilage.

SYNOVIUM/JOINT FLUID: No joint effusion or synovial thickening.

NEUROVASCULAR STRUCTURES: Normal in course and caliber.

PERIPHERAL SOFT TISSUES: Normal.

IMPRESSION:

1. Tear of the superior glenoid labrum.
2. Severe acromioclavicular joint disease may contribute to rotator cuff impingement.

Digitally Signed By: Imam, Naiyer
 Digitally Signed Date: 03/08/2022 4:27 PM