



## Community Medical Imaging

ACR Accredited Facility  
159-16 Union Tpke • Fresh Meadows, NY 11366  
Tel: 718-275-1010 • Fax: 718-591-3300

PERVAIZ QURESHI, M.D.  
79-09 B NORTHERN BLVD  
JACKSON HEIGHTS, NY 11372

PATIENT: TANZILA CHOWDHURY  
DOB: 02/03/1995  
DOS: 04/05/2022  
CHART #: 24809  
EXAM: MRI OF THE LEFT SHOULDER WITHOUT CONTRAST

HISTORY: Posterior pain.

TECHNIQUE: Multiplanar MR imaging of the left shoulder was performed without contrast on Hitachi open MRI unit.

Coronal T1, T2 and STIR; Sagittal T2; Axial PD and gradient echo images of the shoulder were obtained.

COMPARISON: None.

Additional sanitizing / safety protocols recommended by the CDC were performed.

FINDINGS: Mild tendinosis changes are seen at the supraspinatus and infraspinatus tendons. There is no impingement. The supraspinatus and infraspinatus tendons are intact. The teres minor tendon is unremarkable. There is slight disorganization of the fibers of the subscapularis tendon at the upper aspect.

There is type I acromion. There is no impingement or lateral downsloping. There is no muscular injury.

There is no fracture or bone bruise.


The glenoid is unremarkable. There is no subluxation. Morphology of the humeral head is unremarkable.

The anterior and posterior labra are intact. There is no attenuation. The superior labrum and biceps anchor are unremarkable.

There is no biceps tendon tear or tenosynovitis. There is no laxity or tear of the transverse humeral ligament.

The upper humerus is unremarkable.

There is no muscular injury. There is no hematoma or seroma.



## Community Medical Imaging

ACR Accredited Facility  
159-16 Union Tpke • Fresh Meadows, NY 11366  
Tel: 718-275-1010 • Fax: 718-591-3300

PATIENT: TANZILA CHOWDHURY  
DOB: 02/03/1995  
DOS: 04/05/2022  
CHART #: 24809  
EXAM: MRI OF THE LEFT SHOULDER WITHOUT CONTRAST  
PAGE 2

There is no AC separation or fracture. Chest wall is unremarkable.

### IMPRESSION:

1. TENDINOSIS CHANGES ARE SEEN AT THE SUPRASPINATUS AND INFRASPINATUS TENDONS, AS NOTED.
2. A PARTIAL-THICKNESS TEAR IS SEEN AT THE UPPER ASPECT OF THE SUBSCAPULARIS TENDON, AS NOTED. THERE IS NO ATTENUATION.

Thank you for referring this patient to us.

Andrew McDonnell, MD  
Neuroradiologist  
Diplomate, American Board of Radiology  
AM/man/pr D: 04/05/2022

E-Sig By A. McDonnell, MD on 04/06/2022 06:55:01