

# CHARACTERISTIC CLASSES II

NILAY KUMAR

## 1. STIEFEL-WHITNEY CLASSES

Recall from last time the axiomatic definition of the Stiefel-Whitney classes of a vector bundle  $\xi : E \rightarrow B$ .<sup>1</sup>

**Theorem 1.** *Let  $\xi : E \rightarrow B$  be a real vector bundle. Then there exists a unique sequence of cohomology classes*

$$w_i(\xi) \in H^i(B, \mathbb{F}_2)$$

for  $i = 0, 1, 2, \dots$  called the Stiefel-Whitney classes of  $\xi$  satisfying the following properties:

- (I) the class  $w_0(\xi)$  is equal to the unit element  $1 \in H^0(B, \mathbb{F}_2)$  and  $w_i(\xi) = 0$  for  $i > \text{rk } \xi$ ;
- (II) if  $f^*\xi$  is the pullback of  $\xi$  along  $f : A \rightarrow B$  then  $w_i(f^*\xi) = f^*w_i(\xi)$ ;
- (III) if  $\eta : E' \rightarrow B$  is another real vector bundle then

$$w_k(\xi \oplus \eta) = \sum_{i=0}^k w_i(\xi) \smile w_{k-i}(\eta);$$

- (IV) if  $\gamma_1^1$  is the tautological line bundle over  $\mathbb{R}P^1$  then  $w_1(\gamma_1^1) \in H^1(\mathbb{R}P^1, \mathbb{F}_2) = \mathbb{F}_2$  is the unique nonzero element.

make a remark about bundle map diagrams just being pullback diagrams, as Dylan pointed out last time. so we don't lose anything on II

Thus given, the Stiefel-Whitney classes allowed us to make some strong statements about parallelizability and cobordisms. The goal of the first half of this talk is to sketch a proof of the above theorem, i.e. show that Stiefel-Whitney classes do indeed exist. We will first prove the existence of certain analogous classes:

**Theorem 2.** *Let  $\iota_n : O(n-1) \rightarrow O(n)$  and  $p_{ij} : O(i) \times O(j) \rightarrow O(i+j)$  be the obvious inclusions. Denote by the same symbols the induced maps on classifying spaces. Then there are unique classes  $w_i \in H^i(\mathcal{B}O(n), \mathbb{F}_2)$  satisfying:*

- (I)  $w_0 = 1$  and  $w_i = 0$  if  $i > n$ ;
- (II)  $\iota_n^* w_i = w_i$  (and hence  $\iota_n^* w_n = 0$ );
- (III)  $p_{ij}^* w_k = \sum_{a+b=k} w_a \otimes w_b$ ;
- (IV)  $w_1 \in H^1(\mathcal{B}O(1), \mathbb{F}_2) = H^1(\mathbb{R}P^\infty, \mathbb{F}_2)$  is the unique nonzero element.

---

Date: January 20, 2016.

<sup>1</sup>Throughout, all base spaces are assumed to be connected and paracompact.

Here, as usual,

$$\mathcal{B} : \text{GRP} \rightarrow \text{TOP}$$

denotes the classifying space functor. Almost all the work lies in computing the cohomology of  $\mathcal{B}\text{O}(n)$ . In an earlier talk, Guchuan mentioned that – via a number of computations using spectral sequences – one can show the following:

**Lemma 3.** *Let  $\Sigma_n$  be the symmetric group on  $n$  letters. Then there is a map  $\Psi_n : (\mathbb{R}P^\infty)^n \rightarrow \mathcal{B}\text{O}(n)$  inducing an isomorphism*

$$\Psi_n^* : H^\bullet(\mathcal{B}\text{O}(n), \mathbb{F}_2) \cong H^\bullet((\mathbb{R}P^\infty)^n, \mathbb{F}_2)^{\Sigma_n} \cong \mathbb{F}_2[\sigma_1, \dots, \sigma_n],$$

where  $\Sigma_n$  acts on  $(\mathbb{R}P^\infty)^n$  by permutation and hence  $\sigma_i$  are the symmetric polynomials in  $n$  variables with  $\deg \sigma_i = i$ .

*Proof.* This is rather technical and requires more wizardry with spectral sequences than I am familiar with. Reference [unfinished May](#).  $\square$

With this in hand, the proof of Theorem 2 is now a straightforward diagram chase with symmetric polynomials.

*Proof of Theorem 2.* We begin by proving existence of the classes  $w_i$ . Define the Stiefel-Whitney classes as

$$w_i \equiv (\Psi_n^*)^{-1} \sigma_i$$

where  $\Psi_n^*$  is the isomorphism from Lemma 3 above, and where

$$\begin{aligned} \sigma_1 &= x_1 + \dots + x_n \\ \sigma_2 &= x_1 x_2 + x_1 x_3 + \dots + x_{n-1} x_n \\ &\vdots \\ \sigma_n &= x_1 \dots x_n, \end{aligned}$$

are the symmetric polynomials on the generators of  $H^\bullet((\mathbb{R}P^\infty)^n, \mathbb{F}_2) \cong \otimes_i^n \mathbb{F}_2[x_i]$ . Set  $\sigma_0 = 1$  and  $\sigma_i = 0$  for  $i > n$ . Under these definitions, axiom I is immediate.

Next, denoting  $h_n : \mathcal{B}\text{O}(1)^{n-1} \rightarrow \mathcal{B}\text{O}(1)^n$  the map induced by the inclusion, we have a commutative diagram of inclusions

$$\begin{array}{ccc} \mathcal{B}\text{O}(1)^{n-1} & \xrightarrow{\Psi_{n-1}} & \mathcal{B}\text{O}(n-1) \\ \downarrow h_n & & \downarrow \iota_n \\ \mathcal{B}\text{O}(1)^n & \xrightarrow{\Psi_n} & \mathcal{B}\text{O}(n) \end{array}$$

and taking cohomology,

$$\begin{array}{ccc} H^\bullet(\mathcal{B}\text{O}(n)) & \xrightarrow{\Psi_n^*} & H^\bullet(\mathcal{B}\text{O}(1)^n) \\ \downarrow \iota_n^* & & \downarrow h_n^* \\ H^\bullet(\mathcal{B}\text{O}(n-1)) & \xrightarrow{\Psi_{n-1}^*} & H^\bullet(\mathcal{B}\text{O}(1)^{n-1}) \end{array}$$

Clearly  $h_n^* x_i = x_i$  for  $i < n$  and  $h_n^* x_n = 0$ . Therefore  $h_n^* \sigma_i = \sigma_i$ , which implies – by the diagram above – that  $\iota_n^* w_i = w_i$ . So much for axiom II.

Consider now the commutative diagram of inclusions

$$\begin{array}{ccc} \mathcal{B}O(1)^i \times \mathcal{B}O(1)^j & \xrightarrow{\Psi_i \times \Psi_j} & \mathcal{B}O(i) \times \mathcal{B}O(j) \\ \parallel & & \downarrow p_{ij} \\ \mathcal{B}O(1)^{i+j} & \xrightarrow{\Psi_{i+j}} & \mathcal{B}O(i+j) \end{array}$$

which after taking cohomology and applying Künneth becomes

$$\begin{array}{ccc} H^\bullet(\mathcal{B}O(i+j)) & \xrightarrow{\Psi_{i+j}^*} & H^\bullet(\mathcal{B}O(1)^{i+j}) \\ \downarrow p_{ij}^* & & \downarrow \\ H^\bullet(\mathcal{B}O(i)) \otimes H^\bullet(\mathcal{B}O(j)) & \xrightarrow{\Psi_i^* \otimes \Psi_j^*} & H^\bullet(\mathcal{B}O(1)^i) \otimes H^\bullet(\mathcal{B}O(1)^j) \end{array}$$

In this diagram, all the arrows are injective except for  $p_{ij}^*$  and hence  $p_{ij}^*$  is injective. Moreover,

$$(\Psi_i^* \otimes \Psi_j^*)p_{ij}^* w_k = \Psi_{i+j}^* w_k = \sigma_k(x_1, \dots, x_{i+j}).$$

Some algebra with symmetric polynomials reveals that

$$\sigma_k(x_1, \dots, x_{i+j}) = \sum_{a+b=k} \sigma_a(x_1, \dots, x_i) \sigma_b(x_{i+1}, \dots, x_{i+j}),$$

whence

$$(\Psi_i \otimes \Psi_j^*)p_{ij}^* w_k = \sum_{a+b=k} \Psi_i^* w_a \otimes \Psi_j^* w_b = (\Psi_i^* \otimes \Psi_j^*) \sum_{a+b=k} w_a \otimes w_b,$$

proving axiom III by injectivity of  $\Psi_i \otimes \Psi_j$ .

Axiom IV is clear:  $w_1 = (\Psi_1^*)^{-1} \sigma_1(x_1) = x_1$ , the nonzero element in  $H^1(\mathbb{R}P^\infty, \mathbb{F}_2)$ . Finally, we prove uniqueness by induction on  $n$ . The base case  $n = 1$  is trivial. Assume uniqueness of the  $w_i$  in  $H^\bullet(\mathcal{B}O(m), \mathbb{F}_2)$  for  $m < n$ . Then for  $i < n$  the  $w_i \in H^\bullet(\mathcal{B}O(n), \mathbb{F}_2)$  are uniquely determined by axiom II and the fact that  $\iota_n$  is an isomorphism in degrees smaller than  $n$ . For  $i = n$  we note that  $p_{1,n-1}^* w_n \in H^\bullet(\mathcal{B}O(1), \mathbb{F}_2) \otimes H^\bullet(\mathcal{B}O(n-1), \mathbb{F}_2)$  and hence  $w_n$  is determined by the induction hypothesis since  $p_{1,n-1}^*$  is injective. This completes the proof.  $\square$

To relate these classes sitting in the cohomology of  $\mathcal{B}O(n)$  to the previous axiomatic definition of classes sitting in the cohomology of the base  $B$ , we need a classification theorem for vector bundles on  $B$ . Before we start, we note that

$$\text{Vect}_{\mathbb{R}} : \text{TOP}^{\text{op}} \rightarrow \text{SET}$$

will denote the contravariant functor taking  $B$  to the set (isomorphism classes) of real vector bundles over  $B$  and taking  $f : B \rightarrow B'$  to the pullback  $f^* : \text{Vect}_{\mathbb{R}} B' \rightarrow \text{Vect}_{\mathbb{R}} B$ . Recall that two vector bundles over  $B$  are isomorphic if there is a map lifting  $\text{id}_B$  that is a fiberwise linear isomorphism.

**Theorem 4.** *The space  $\mathcal{B}O(n) \cong \text{Gr}_n \mathbb{R}^\infty$  classifies vector bundles, i.e. the natural transformation*

$$\Phi : [-, \mathcal{B}O(n)] \dashrightarrow \text{Vect}_{\mathbb{R}} -,$$

*given by pullback  $[f] \mapsto f^* \gamma_\infty^n$  of the tautological bundle, is a natural isomorphism.*

We first check that  $\Phi$  is well-defined:

**Lemma 5.** *The pullbacks of a vector bundle along homotopic maps are isomorphic, i.e. the functor  $\text{Vect}_{\mathbb{R}}$  factors through the homotopy category.*

*Proof sketch.* Let  $\xi : E \rightarrow B$  be a rank  $n$  vector bundle and let  $f, g : A \rightarrow B$  be two maps homotopic via  $h : A \times I \rightarrow B$ . Note first that  $h^*E|_{A \times \{0\}} = f^*E$  and  $h^*E|_{A \times \{1\}} = g^*E$ . Thus it suffices to prove that for a vector bundle  $\eta : F \rightarrow A \times I$ , there is an isomorphism  $F|_{A \times \{0\}} \cong F|_{A \times \{1\}}$ . The idea, roughly, is to find countably many local trivializations over  $U_i \subset B$  for  $E$  and to then locally push  $F|_{A \times \{0\}}$  to the right along  $U_i \times I$ . For details, see [Hatcher](#), [VBKT](#).  $\square$

*Proof of Theorem 4.* Naturality of  $\Phi$  follows immediately from the fact that if  $\alpha : A \rightarrow B$  and  $f \in [B, \mathcal{B}\mathcal{O}(n)]$  then  $(f \circ \alpha)^*\gamma_n^\infty = \alpha^*f^*\gamma_n^\infty$ . We now prove that  $\Phi_B : [B, \mathcal{O}(n)] \rightarrow \text{Vect}_{\mathbb{R}} B$  is a bijection.

The key observation is as follows. Let  $\xi : E \rightarrow B$  be a rank  $n$  vector bundle. Then an isomorphism  $E \cong f^*\gamma_n^\infty$  (for some map  $f : B \rightarrow \text{Gr}_n \mathbb{R}^\infty$ ) is equivalent to a map  $g : E \rightarrow \mathbb{R}^\infty$  that is a linear injection on each fiber. To see this, suppose first that we have such an isomorphism. Then we have a commutative diagram

$$\begin{array}{ccccc} E \cong f^*\gamma_n^\infty & \xrightarrow{\tilde{f}} & \gamma_n^\infty & \xrightarrow{\pi} & \mathbb{R}^\infty \\ \downarrow \xi & & \downarrow & & \\ B & \xrightarrow{f} & \text{Gr}_n \mathbb{R}^\infty & & \end{array}$$

where  $\pi$  is the obvious projection. Now  $\pi \circ \tilde{f} : E \rightarrow \mathbb{R}^\infty$  is a fiberwise linear injection as both  $f$  and  $\pi$  are. Conversely, given  $g : E \rightarrow \mathbb{R}^\infty$  a fiberwise linear injection, we can define a map  $f : B \rightarrow \text{Gr}_n \mathbb{R}^\infty$  given by  $x \mapsto [g(\xi^{-1}(x))]$ . Then  $E \cong f^*\gamma_n^\infty$  because we have fiberwise linear isomorphisms

$$f^*\gamma_n^\infty|_b \cong \gamma_n^\infty|_{f(b)} \cong E|_b.$$

Now, for surjectivity of  $\Phi_B$ , given  $\xi : E \rightarrow B$  it suffices by the previous paragraph to construct a map  $E \rightarrow \mathbb{R}^\infty$  a linear injection on each fiber. To do this, we fix countably many local trivializations over  $U_i \subset B$  of  $E$  together with partitions of unity  $\phi_i$  subordinate to the  $U_i$ . Then for each  $i$  we obtain a map  $g_i : E \rightarrow \mathbb{R}^n$  that is zero outside  $\xi^{-1}U_i$  and the composition  $E \rightarrow U_i \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  otherwise. Summing  $g = \sum g_i$ , we obtain a map  $g : E \rightarrow (\mathbb{R}^n)^\infty \cong \mathbb{R}^\infty$  that is obviously a linear injection on fibers.

For injectivity, suppose we have isomorphisms  $E \cong f_0^*\gamma_n^\infty$  and  $E \cong f_1^*\gamma_n^\infty$  for two maps  $f_0, f_1 : B \rightarrow \text{Gr}_n \mathbb{R}^\infty$ . By arguments above, we obtain maps  $g_0, g_1 : E \rightarrow \mathbb{R}^\infty$  that are fiberwise linear injections. We claim that  $g_0$  and  $g_1$  are homotopic through maps  $g_t$  that are fiberwise linear injections; this implies that  $f_0$  and  $f_1$  are homotopic via  $f_t(x) = g_t(\xi^{-1}x)$ . To do this, we first homotope  $g_0$  so that it takes values only in odd coordinates via

$$(x_1, x_2, \dots) \mapsto (1-t)(x_1, x_2, \dots) + t(x_1, 0, x_2, 0, \dots)$$

and homotope  $g_1$  so that it takes values only in even coordinates similarly. Now  $g_t = (1-t)g_0 + tg_1$  provides the necessary homotopy and it is clearly linear and injective on fibers.  $\square$

We can finally prove the existence of Stiefel-Whitney classes for vector bundles.

*Proof of Theorem 8.* Let  $\xi : E \rightarrow B$  be a real vector bundle of rank  $n$ . By Theorem 4 there exists a unique map  $\Phi_B : B \rightarrow \mathcal{BO}(n)$  such that  $E \cong f^*\gamma_n^\infty$ . Define

$$w_i(\xi) \equiv \Phi_B^* w_i.$$

Axiom I now follows immediately from Theorem 2. Now suppose we have a pullback diagram

$$\begin{array}{ccc} f^*E & \longrightarrow & E \\ \downarrow & & \downarrow \xi \\ A & \xrightarrow{f} & B \end{array}$$

Then

$$w_i(f^*\xi) = \Phi_A^* w_i = f^* \Phi_B^* w_i = f^* w_i(\xi),$$

which proves axiom II.

To prove axiom III we recall that the direct sum of vector bundles can be described as a pullback

$$\begin{array}{ccc} E \oplus E' \cong \Delta^*(E \times E') & \longrightarrow & E \times E' \\ \downarrow & & \downarrow \\ B & \xrightarrow{\Delta} & B \times B \end{array}$$

where  $\Delta$  is the diagonal map. Then

$$w_k(\xi \oplus \eta) = w_k(\Delta^*(\xi \times \eta)) = \Phi_{B \times B}^* w_k$$

where  $\Phi_{\xi \times \eta} : B \times B \rightarrow \mathcal{BO}(n+m)$  is the unique classifying map. **Figure out how to prove axiom III.**

The tautological line bundle on  $\mathbb{R}P^1$  is given by the pullback of  $\gamma_1^\infty$  along the inclusion  $j : \mathbb{R}P^1 \hookrightarrow \mathbb{R}P^\infty$  so  $w_1(\gamma_1^1) = j^* w_1$  is the unique nonzero element in  $H^1(\mathbb{R}P^1, \mathbb{Z}/2)$  by axiom IV of Theorem 2 and because  $j^*$  is an isomorphism in degrees  $\leq 1$ .

**Uniqueness blah blah blah.** □

**Application:  $w_1(\xi) = 0$  if and only if  $E$  is orientable? Why?**

## 2. CHERN CLASSES

Chern classes are the complex analog of Stiefel-Whitney classes, but with integral coefficients.

**Theorem 6.** Let  $\iota_n : \mathcal{U}(n-1) \rightarrow \mathcal{U}(n)$  and  $p_{ij} : \mathcal{U}(i) \times \mathcal{U}(j) \rightarrow \mathcal{U}(i+j)$  be the obvious inclusions. Denote by the same symbols the induced maps on classifying spaces. Then there are unique Chern classes  $c_i \in H^{2i}(\mathcal{BU}(n), \mathbb{Z})$  satisfying:

- (I)  $c_0 = 1$  and  $c_i = 0$  if  $i > n$ ;
- (II)  $\iota_n^* c_i = c_i$  (and hence  $\iota_n^* c_n = 0$ );
- (III)  $p_{ij}^* c_k = \sum_{a+b=k} c_a \otimes c_b$ ;
- (IV)  $c_1 \in H^2(\mathcal{BU}(1), \mathbb{Z}) = H^2(\mathbb{C}P^\infty, \mathbb{Z})$  is the canonical generator.

*Proof.* The proof is exactly the same as in the case of Stiefel-Whitney classes. We note, in particular, that  $H^\bullet(\mathcal{BU}(n), \mathbb{Z}) = \mathbb{Z}[c_1, \dots, c_n]$ . □

Chern classes can be related to complex vector bundles via the following analog of Theorem 4. Notice that here  $\text{Gr}_n V$  denotes the space of complex  $n$ -planes in  $V$ .

**Theorem 7.** *The space  $\mathcal{BU}(n) \cong \text{Gr}_n \mathbb{C}^\infty$  classifies complex vector bundles, i.e. the natural transformation*

$$\Phi : [-, \mathcal{BU}(n)] \longrightarrow \text{Vect}_{\mathbb{C}} -,$$

*given by pullback  $[f] \mapsto f^* \gamma_\infty^n$  of the tautological bundle, is a natural isomorphism.*

We can now define the Chern class of a complex vector bundle.

**Theorem 8.** *Let  $\xi : E \rightarrow B$  be a complex vector bundle. Then there exists a unique sequence of cohomology classes*

$$c_i(\xi) \in H^{2i}(B, \mathbb{Z})$$

*for  $i = 0, 1, 2, \dots$  called the Chern classes of  $\xi$  satisfying the following properties:*

- (I) *the class  $c_0(\xi)$  is equal to the generator  $1 \in H^0(B, \mathbb{Z})$  and  $c_i(\xi) = 0$  for  $i > \text{rk } \xi$ ;*
- (II) *if  $f^* \xi$  is the pullback of  $E$  along  $f : A \rightarrow B$  then  $c_i(f^* \xi) = f^* c_i(\xi)$ ;*
- (III) *if  $\eta : E' \rightarrow B$  is another complex vector bundle then*

$$c_k(\xi \oplus \eta) = \sum_{i=0}^k c_i(\xi) \smile c_{k-i}(\eta);$$

- (IV) *if  $\gamma_1^1$  is the tautological complex line bundle over  $\mathbb{CP}^1$  then  $c_1(\gamma_1^1) \in H^2(\mathbb{CP}^1, \mathbb{Z}) = \mathbb{Z}$  is the canonical generator.*

How does the Chern class restrict to the Stiefel-Whitney class? See p.47 of May unfinished

What are some applications of this machinery different than the ones Yajit gave?