



Introduction to Statistical Methods

BITS Pilani

Pilani Campus

ISM TEAM



BITS Pilani
Pilani Campus

Course No: AIML ZC418
Webinar 1 : 11.11.2025

Team ISM

Topics - Webinar



- Descriptive Statistics
 - ❖ Measures of Central Tendency
 - ❖ Measures of Variability
- Probability
 - ❖ Introduction and Basics
 - ❖ Conditional probability

Measures of Central Tendency

1) A psychologist wrote a computer program to simulate the way a person responds to a standard IQ test. To test the program, he gave the computer 15 different forms of a popular IQ test and computed its IQ from each form.

IQ Values:

134	136	137	138	138	143	144	144	145	146	146	146
147	148	153									

Find the following Statistical measures:

- i. Mean, median, and mode
- ii. Range, Variance and standard deviation
- iii. The interquartile range.
- v. Identify potential outliers, if any.
- vi. Construct and interpret a boxplot

Measures of Central Tendency

Arranging the data in ascending order:

134, 136, 137, 138, 138, 143, 144, 144, 145, 146, 146, 146, 147, 148, 153

N=15

Measure	Formula	Solution
Mean	$\mu = \frac{\sum_{i=1}^n x_i}{n}$	$\mu = 143$
Median	$p = \frac{n+1}{2}$	Median = 144
Mode	The mode is the value or values that occur most frequently in the data set. A data set can have more than one mode, and it can also have no mode	Mode = 146

Mode, Bimodal, and Multimodal



A given set of data may have one or more than one Mode. A set of numbers with one Mode is unimodal, a set of numbers having two Modes is bimodal, a set of numbers having three Modes is trimodal, and any set of numbers having four or more than four Modes is known as multimodal.

Bimodal Mode – A set of data including two modes is identified as a bimodal model. This indicates that there are two data values that possess the highest frequencies. For example, the mode of data set B = { 8, 12, 12, 14, 15, 19, 17, 19} is 12 and 19 as both 12 and 19 are repeated twice in the given set.

No mode:

If no number in a set of numbers occurs more than once, that set has no mode: 3, 6, 9, 16, 27, 37, 48.

A unimodal mode is a set of data with only one mode.

A bimodal mode is a set of data that has two modes.

A trimodal mode is a set of data that has three modes.

Measures of Variability

Range	Range = $x_n - x_1$	Minimum = 134 Maximum = 153 Range R = 19
Variance	For a Population $\sigma^2 = \frac{\sum_{i=1}^n (x_i - \mu)^2}{n}$ For a Sample $s^2 = \frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n - 1}$	Variance = 26
Standard deviation	For a Population $\sigma = \sqrt{\frac{\sum_{i=1}^n (x_i - \mu)^2}{n}}$ For a Sample $s = \sqrt{\frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n - 1}}$	Standard Deviation = 5.09901951

$$\begin{aligned}
 \sigma^2 &= \frac{\sum (X - \mu)^2}{N} \\
 &= \frac{\sum (X^2 - 2\mu X + \mu^2)}{N} \\
 &= \frac{\sum X^2}{N} - \frac{2\mu \sum X}{N} + \frac{N\mu^2}{N} \\
 &= \frac{\sum X^2}{N} - 2\mu^2 + \mu^2 \\
 &= \frac{\sum X^2}{N} - \mu^2
 \end{aligned}$$

Measures of Central Tendency




- ❖ **Minimum.**
- ❖ **Q1** (the first quartile, or the 25% mark).
- ❖ **Median.**
- ❖ **Q3** (the third quartile, or the 75% mark).
- ❖ **Maximum.**

Measures of Central Tendency

Quartiles	Quartiles separate a data set into four sections. The median is the second quartile Q_2. It divides the ordered data set into higher and lower halves. The first quartile, Q_1, is the median of the lower half not including Q_2. The third quartile, Q_3, is the median of the higher half not including Q_2.	Quartiles: <table> <tr> <td>Q_1</td> <td>--></td> <td>138</td> </tr> <tr> <td>Q_2</td> <td>--></td> <td>144</td> </tr> <tr> <td>Q_3</td> <td>--></td> <td>146</td> </tr> </table>	Q_1	-->	138	Q_2	-->	144	Q_3	-->	146
Q_1	-->	138									
Q_2	-->	144									
Q_3	-->	146									
Interquartile range	$IQR = Q_3 - Q_1$	Interquartile Range $IQR = 8$									
Potential outliers, if any.	Upper Fence = $Q_3 + 1.5 \times IQR$ Lower Fence = $Q_1 - 1.5 \times IQR$	none									


Graphical Representation:Box plot




Box-plot for the given data:




Population size: 15
Median: 144
Minimum: 134
Maximum: 153
First quartile: 138
Third quartile: 146
Interquartile Range: 8
Outliers: none



Real-time Example:



Q1 , Median (Q2) and Q3 (n is even)



Example:



Data: 5, 7, 8, 9, 10, 12, 13, 15, 18, 30

Five- point summary:

- **Q1 = 8**
- **Q2 (Median) = 11**
- **Q3 = 15**
- **IQR = 15 - 8 = 7**
- **Lower bound = 8 - (1.5 × 7) = -2.5**
- **Upper bound = 15 + (1.5 × 7) = 25.5**
- ✓ Any value **> 25.5** or **< -2.5** is an **outlier** → here, **30** is an outlier.



innovate

achieve

lead

2) Consider the following statistical summary of a dataset. Write at least three useful observations as a part of data pre – processing. **[Midsem Sep' 2023]**

	Nr	Cells	QValue	Fat	Protein
count	969.000000	969.000000	969.000000	969.000000	969.000000
mean	7.074303	358.284830	90.016584	3.620279	3.300196
std	4.759793	344.324223	4.998924	0.349956	0.136071
min	1.000000	0.000000	62.650000	2.240000	2.750000
25%	3.000000	157.000000	87.570000	3.410000	3.220000
50%	6.000000	283.000000	90.750000	3.610000	3.310000
75%	10.000000	476.000000	93.400000	3.820000	3.380000
max	20.000000	5226.000000	100.000000	5.420000	3.840000

Contd....



Observation 1: The variation in Protein is found very low when compared with remaining using coefficient of variation = $sd/mean$

Observation 2: The range in Protein is found very low when compared with remaining using range = $max-Min$

Observation 3: The middle range in Protein is found very low when compared with remaining using quartile range = $Q3(75\%) - Q1(25\%)$

Etc.....

Basic Probability

3) If $P(A) = 1/2$, $P(B) = 1/3$ and $P(A \cap B) = 1/5$ then find

a). $P(A \cup B)$

b). $P(A^c \cap B)$

c). $P(A \cap B^c)$

d). $P(A^c \cap B^c)$

e). $P(A^c \cup B^c)$

f). $P((A \cup B)^c)$

Solution

$$\text{a) } P(A \cup B) = P(A) + P(B) - P(A \cap B) = 1/2 + 1/3 - 1/5 = 19/30 = 0.6333$$

$$\text{b) } P(A^c \cap B) = P(B - A) = P(B - (A \cap B)) = P(B) - P(A \cap B) = 1/3 - 1/5 = 2/15 = 0.1333$$

$$\text{c) } P(A \cap B^c) = P(A - B) = P(A - (A \cap B)) = P(A) - P(A \cap B) = 1/2 - 1/5 = 3/10 = 0.3$$

$$\text{d) } P(A^c \cap B^c) = P((A \cup B)^c) = 1 - P(A \cup B) = 1 - 19/30 = 11/30 = 0.3667$$

$$\text{e) } P(A^c \cup B^c) = P((A \cap B)^c) = 1 - P(A \cap B) = 1 - 1/5 = 4/5 = 0.8$$

$$\text{f) } P((A \mid B)^c) = 1 - P(A \mid B) = 1 - 19/30 = 11/30 = 0.3667$$

Basic Probability

4) There is a 1% probability for a hard drive to crash. Therefore, it has two backups, each having a 2 % probability to crash, and all three components are independent of each other. The stored information is lost only in an unfortunate situation when all three devices crash. What is the probability that the information is saved?

Solution. Organize the data. Denote the events, say,

$$H = \{ \text{hard drive crashes} \},$$

$$B_1 = \{ \text{first backup crashes} \}, \quad B_2 = \{ \text{second backup crashes} \}.$$

It is given that H , B_1 , and B_2 are independent,

$$P\{H\} = 0.01, \text{ and } P\{B_1\} = P\{B_2\} = 0.02.$$

Applying rules for the complement and for the intersection of independent events,

$$\begin{aligned} P\{ \text{saved} \} &= 1 - P\{ \text{lost} \} = 1 - P\{H \cap B_1 \cap B_2\} \\ &= 1 - P\{H\} P\{B_1\} P\{B_2\} \\ &= 1 - (0.01)(0.02)(0.02) = 0.999996. \end{aligned}$$

Basic Probability

- 5) A political leader has submitted his nomination to compete in two different electoral constituencies namely A1 and A2. The probability of winning in constituency A1 and A2 is 0.80 and 0.65 respectively. The probability of losing at least one of the constituencies is 0.35. What will be the probability that he will win in exactly one of the constituencies?

We want:

$$P(\text{Win exactly one}) = P(\text{Win A1 and lose A2}) + P(\text{Lose A1 and win A2})$$

- $P(\text{Win A1 and lose A2}) = P(A \cap B^c)$
- $P(\text{Lose A1 and win A2}) = P(A^c \cap B)$

Solution:

A political leader has submitted his nomination to compete in two different electoral constituencies namely A1 and A2. The probability of winning in constituency A1 and A2 is 0.80 and 0.65 respectively. The probability of losing at least one of the constituencies is 0.35. What will be the probability that he will win in one of the constituencies?

Assume that A, B be the events defined as follows:

A : "Winning in constituency A1"

B : "Winning in constituency A2"

Given:

$P(A) = 0.80$, $P(B) = 0.65$

and $P(\bar{A} \cup \bar{B}) = 0.35$

Now, $\therefore P(\bar{A} \cup \bar{B}) = 0.35$

$\therefore P(\overline{A \cap B}) = 0.35$

$\Rightarrow 1 - P(A \cap B) = 0.35$

$\Rightarrow P(A \cap B) = 0.65$

Then, $P(A \cup B) = P(A) + P(B) - P(A \cap B)$

$= 0.80 + 0.65 - 0.65$

$\therefore P(A \cup B) = 0.80$

Then, $P(\text{He will win in one of the constituencies}) = P(A \cup B) - P(A \cap B)$
 $= 0.80 - 0.65$

$\therefore P(\text{He will win in one of the constituencies}) = 0.15$

$P(\text{He will win in constituency A1 ONLY}) = P(A) - P(A \cap B)$
 $= 0.80 - 0.65$
 $= 0.15$

$P(\text{He will win in constituency A2 ONLY}) = P(B) - P(A \cap B)$
 $= 0.65 - 0.65$
 $= 0$

Independent vs. Dependent Events



Using the bag of marbles on the left, what is the probability of pulling a black marble two times in a row? $P(\text{black}, \text{black})$

When you put 1st marble back in
(Independent Events)


$$\frac{2}{10} * \frac{2}{10}$$

$$\frac{1}{5} * \frac{1}{5} = \frac{1}{25}$$

When you KEEP 1st marble
(Dependent Events)

$$\frac{2}{10} * \frac{1}{9}$$

$$\frac{1}{5} * \frac{1}{9}$$





Independent events

6) Comment on the statement:

“If two events are mutually exclusive, then they are independent also and vice versa”

Two independent events cannot be mutually exclusive events - unless one or both events have a probability of zero (meaning one of the events is impossible).



Independent events

7) Let A and B be the two possible outcomes of an experiment and suppose $P(A) = 0.4$, $P(B) = p$ and $P(A \cup B) = 0.7$

- (i) For what choice of 'p' are A and B mutually exclusive?
- (ii) For what choice of 'p' are A and B independent?

Solution:

(i) If A and B are mutually exclusive then $P(A \cap B) = 0$

Thus $P(A \cup B) = P(A) + P(B) - P(A \cap B)$ becomes

$$0.7 = 0.4 + P(B) - 0$$

$$\therefore P(B) = 0.7 - 0.4 = 0.3$$

(ii) If A and B are independent then $P(A \cap B) = P(A) \cdot P(B)$

Thus $P(A \cup B) = P(A) + P(B) - P(A \cap B)$ becomes

$$P(A \cup B) = P(A) + P(B) - P(A) \cdot P(B)$$

$$0.7 = 0.4 + P - 0.4 \cdot P$$

$$= 0.4 + P(1 - 0.4)$$

$$= 0.4 + 0.6P$$

$$0.6P = 0.7 - 0.4 = 0.3$$

$$P = \frac{0.3}{0.6} = \frac{1}{2} = 0.5$$

$$\therefore P(B) = 0.5$$

Answer. (i) $p=0.3$, (ii) $p=0.5$

Conditional Probability

8. Ninety percent of flights depart on time. Eighty percent of flights arrive on time. Seventy-five percent of flights depart on time and arrive on time.
- You are meeting a flight that departed on time. What is the probability that it will arrive on time?
 - You have met a flight, and it arrived on time. What is the probability that it departed on time?
 - Are the events, departing on time and arriving on time, independent?

Let:

- D : Flight departs on time
 - A : Flight arrives on time
- $P(D) = 0.90$
 - $P(A) = 0.80$
 - $P(D \cap A) = 0.75$

Solution:



- a) You are meeting a flight that departed on time. What is the probability that it will arrive on time?

$$P(A | D) = \frac{P(D \cap A)}{P(D)}$$
$$P(A | D) = \frac{0.75}{0.90} = \frac{75}{90} = \frac{5}{6} \approx 0.833$$

- b) You have met a flight, and it arrived on time. What is the probability that it departed on time?

$$P(D | A) = \frac{P(D \cap A)}{P(A)}$$
$$P(D | A) = \frac{0.75}{0.80} = \frac{75}{80} = \frac{15}{16} = 0.9375$$

c) Are the events, departing on time and arriving on time, independent?

Two events A and B are independent if:

$$P(A \cap B) = P(A) \cdot P(B)$$

Check:

$$P(D) \cdot P(A) = (0.90)(0.80) = 0.72$$

But:

$$P(D \cap A) = 0.75 \neq 0.72$$

So, the events are NOT independent.



Conditional Probability

- 9) In an online shopping survey, 30% of persons made shopping in Flipkart, 45% of persons made shopping in Amazon and 5% made purchases in both. If a person is selected at random, find
- i) the probability that he makes shopping in at least one of two companies
 - ii) the probability that he makes shopping in Amazon given that he already made shopping in Flipkart.
 - iii) the probability that the person will not make shopping in Flipkart given that he already made purchase in Amazon.

Conditional Probability

Solution: Given $P(F) = 30\% = 0.30$

$$P(A) = 45\% = 0.45$$

$$P(F \cap A) = 5\% = 0.05$$

$$\text{i) } P(F \cup A) = P(F) + P(A) - P(F \cap A)$$

$$= 0.30 + 0.45 - 0.05 = 0.7$$

$$\begin{aligned} \text{ii) } P(A | F) &= \frac{P(A \cap F)}{P(F)} \\ &= \frac{0.05}{0.30} = 0.167 \end{aligned}$$

$$\begin{aligned} \text{iii) } P(F' | A) &= \frac{P(F' \cap A)}{P(A)} \\ P(F' \cap A) &= P(A) - P(A \cap F) \\ &= 0.45 - 0.05 \\ &= 0.40 \end{aligned}$$

$$P(F' | A) = \frac{0.40}{0.45} = 0.88$$

Total Probability

10) A businessman goes to hotels X, Y, Z -20%, 50%, 30% of the time, respectively. It is known that 5%, 4%, 8% of the rooms in X, Y, Z hotels have faulty plumbing. Determine the probability that the businessman goes to hotel with faulty plumbing.

Given: **Hotel Visit Probabilities:**

- $P(X) = 0.20$
- $P(Y) = 0.50$
- $P(Z) = 0.30$

Probability of Faulty Plumbing (given hotel):

- $P(F|X) = 0.05$
- $P(F|Y) = 0.04$
- $P(F|Z) = 0.08$

Find the **total probability** that the businessman goes to a hotel with faulty plumbing:

$$P(F) = P(X) \cdot P(F|X) + P(Y) \cdot P(F|Y) + P(Z) \cdot P(F|Z)$$

$$P(F) = (0.20 \times 0.05) + (0.50 \times 0.04) + (0.30 \times 0.08)$$

$$= 0.010 + 0.020 + 0.024 = \boxed{0.054}$$

Total Probability

11) A company receives 50% of customer complaints via phone, 30% via email, and 20% via chat.

Resolution rates:

- Phone: 90%
- Email: 80%
- Chat: 70%

What is the probability that a randomly selected complaint is resolved?

Given:

- $P(\text{Phone}) = 0.50$ (50% of complaints come via phone),
- $P(\text{Email}) = 0.30$ (30% of complaints come via email),
- $P(\text{Chat}) = 0.20$ (20% of complaints come via chat),
- $P(\text{Resolved} \mid \text{Phone}) = 0.90$ (90% of phone complaints are resolved),
- $P(\text{Resolved} \mid \text{Email}) = 0.80$ (80% of email complaints are resolved),
- $P(\text{Resolved} \mid \text{Chat}) = 0.70$ (70% of chat complaints are resolved).

To calculate $P(\text{Resolved})$, we'll use the **Law of Total Probability**. The formula is:

$$P(\text{Resolved}) = P(\text{Phone}) \cdot P(\text{Resolved} \mid \text{Phone}) + P(\text{Email}) \cdot P(\text{Resolved} \mid \text{Email}) + P(\text{Chat}) \cdot P(\text{Resolved} \mid \text{Chat})$$

Solution:

$$P(\text{Resolved}) = 0.50 \cdot 0.90 + 0.30 \cdot 0.80 + 0.20 \cdot 0.70 = 0.45 + 0.24 + 0.14 = 0.83$$

Conclusion:

So, there is an **83%** chance that a randomly selected customer complaint will be resolved, based on the given resolution rates for each channel.

Problem:


innovate

achieve

lead

12. A bag contains 7 green beads and 5 blue beads. A bead is taken from the bag at random, the colour is recorded and the bead is replaced. A second bead is then taken from the bag and its colour is recorded.
- Find the probability that one bead is green and the other is blue
 - Show that the event “the first bead is green” and “the second bead is green” are independent

Solution:



Problem:

innovate


achieve

lead

13. Suppose A, B and C are three mutually exclusive events in a sample space. Given that $S = A \cup B \cup C$, $P(A) = (1/5)P(B)$, and $P(C) = 4P(A)$, find $P(B \cup C)$.

Solution:

$5P(A) = P(B)$
Two or more events are mutually exclusive if they cannot happen at the same time. $P(X \cup Y) = P(X) + P(Y)$


$$\begin{aligned} P(A \cup B \cup C) &= 1 \\ P(A) + P(B) + P(C) &= 1 \\ P(A) + 5P(A) + 4P(A) &= 1 \\ 10P(A) &= 1 \\ P(A) &= \frac{1}{10} \end{aligned}$$
$$\begin{aligned} P(B) &= \frac{5}{10} = \frac{1}{2} \\ P(C) &= \frac{4}{10} = \frac{2}{5} \end{aligned}$$
$$\begin{aligned} P(B \cup C) &= P(B) + P(C) \\ &= \frac{5}{10} + \frac{4}{10} \\ &= \frac{9}{10} \end{aligned}$$

Practice Problems

1. If $P(A) = 1/3$, $P(B) = 1/2$, $P(A/B) = 1/6$ find i). $P(B/A)$ ii). $P(B/A')$ iii). $P(A \cup B / A)$ iv). $P(B/A)$.

2. Three machines A, B, C produce 50%, 30%, and 20% of the items in a factory. The percentage of defective outputs of these machines are 3, 4 and 5 respectively.
 - (i) If an item is selected at random, what is the probability that it is defective? (Ans: 0.037)
 - (ii) If a selected item is defective, what is the probability that it is from machine A? (Ans: 0.4054)



Practice Problems

3. A weather app predicts rain with 90% accuracy when it actually rains. It wrongly predicts rain 10% of the time when it doesn't rain. Suppose it rains only 10% of days in a year. You check the app and it predicts rain. What is the probability it will actually rain? (Ans: 0.5)

4. Consider the following data related to the employees, who are on travel. 40% check work email, 20% use cell phone to stay connected to work, 25% bring laptop with them, 23% check both work email and use cell phone to stay connected, and 50% neither check work email nor use a cell phone to stay connected nor bring a laptop. In addition, 88 out of every 100 who bring a laptop also check work email, and 70 out of every 100 who use a cell phone to stay connected also bring a laptop.

- i) What is the probability that a randomly selected traveller who checks work email also uses a cell phone to stay connected?
- ii) What is the probability that someone who brings a laptop on vacation also uses a cell phone to stay connected?
- iii) If the randomly selected traveller checked work email and brought a laptop, what is the probability that he/she uses a cell phone to stay connected?

TEXT BOOKS

T1 : Statistics for Data Scientists, An introduction to Probability,
Statistics and Data Analysis, Maurits Kaptein et al, Springer 2022

T2 : Probability and Statistics for Engineering and Sciences,
8th Edition, Jay L Devore, Cengage Learning

T3 : Introduction to Time Series and Forecasting, Second Edition,
Peter J Brockwell, Richard A Davis, Springer.

Questions please...



THANK YOU!



BITS Pilani

Pilani Campus

Introduction to Statistical Methods



BITS Pilani
Pilani Campus

Course No: DSECL ZC418 / AIML CZC418
Course Title: ISM
WEBINAR 2 : 05/06/2025

❖ Bayes' Theorem

❖ Naïve Bayes Classifier

❖ Random Variables

BAYES' THEOREM:

Suppose that $E_1, E_2, \dots, E_n$ are mutually exclusive and exhaustive events of a sample space “ S ”

such that $P(E_i) > 0$ for $i = 1, 2, 3, \dots, n$ and A is any arbitrary event of “ S ”

such that $P(A) > 0$ and $A \subseteq \bigcup_{i=1}^n E_i$ then the conditional probability of E_i given A

for $i = 1, 2, 3, \dots, n$ is given by
$$P(E_i/A) = \frac{P(E_i)P(A/E_i)}{\sum_{i=1}^n P(E_i)P(A/E_i)}$$

Bayes' theorem is also known as the formula for the **probability of “causes”**. **As we know, the E_i 's are a partition of the sample space S , and at any given time only one of the events E_i occurs.** Thus we conclude that the Bayes' theorem formula gives the probability of a particular E_i , given the event A has occurred.

Terms related to Bayes Theorem



- **Hypotheses:** Events happening in the sample space $E_1, E_2, \dots, E_n$ is called the hypotheses
 - **Priori Probability:** Priori Probability is the initial probability of an event occurring before any new data is taken into account. $P(E_i)$ is the priori probability of hypothesis E_i .
 - **Posterior Probability:** Posterior Probability is the updated probability of an event after considering new information. Probability $P(E_i|A)$ is considered as the posterior probability of hypothesis E_i .
-

Applications of Bayes Theorem



Bayesian inference is very important and has found application in various activities, including medicine, science, philosophy, engineering, sports, law, etc., and Bayesian inference is directly derived from Bayes' theorem.

Example: Bayes' theorem defines the accuracy of the medical test by taking into account how likely a person is to have a disease and what is the overall accuracy of the test.

Problem 1



A hospital has 4 nurses handling respectively 20%, 40%, 25% and 15% of the patients of all the doctors coming to the hospital. The probability that they misguide the patients are respectively 0.05, 0.03, 0.04 and 0.05. Find the probability that the misguided incident can be blamed on the third nurse.


Problem 1 - Solution

innovate

achieve

lead

Let A_1 : Event nurse 1 handles the patient
 A_2 : Event nurse 2 handles the patient
 A_3 : Event nurse 3 handles the patient
 A_4 : Event nurse 4 handles the patient
 X : Event that the patient is misguided



Given that,

$$P(A_1) = 0.20$$

$$P(A_2) = 0.40$$

$$P(A_3) = 0.25$$

$$P(A_4) = 0.15$$

$$P(X/A_1) = 0.05$$

$$P(X/A_2) = 0.03$$

$$P(X/A_3) = 0.04$$

$$P(X/A_4) = 0.05$$

By total probability,
P(a patient is misguided)

$$\begin{aligned} = P(X) &= P(A_1 \cap X) + P(A_2 \cap X) + P(A_3 \cap X) + P(A_4 \cap X) \\ &= P(A_1)P(X/A_1) + P(A_2)P(X/A_2) + P(A_3)P(X/A_3) \\ &\quad + P(A_4)P(X/A_4) \\ &= (0.20)(0.05) + (0.40)(0.03) + (0.25)(0.04) + (0.15)(0.05) \\ &= 0.01 + 0.012 + 0.01 + 0.0075 \\ &= 0.0395 \\ \therefore P(X) &= 0.0395 \end{aligned}$$

To find the probability that the misguided incident can be blamed on third Nurse:

By applying Bayes' theorem, we have

$$\begin{aligned} P(A_3/X) &= \frac{P(A_3 \cap X)}{P(X)} = \frac{P(A_3)P(X/A_3)}{P(X)} \\ &= \frac{(0.25)(0.04)}{0.0395} = \frac{0.01}{0.0395} = 0.2532 \end{aligned}$$

$\therefore$ The probability that the misguided incident can be blamed on third nurse is 0.2532 (or) 25.32%

Problem 2



Three machines A, B and C produce respectively 50%, 30% and 20% of the total number of items of a factory. The percentage of defective outputs of these machines are 3%, 5% and 2%. An item is selected at random and is found to be defective. (i) Find the probability that the item was produced by machine C? (ii) What is the probability that the item was produced by machine C or B? (iii) What is the probability that an item selected at random is found to be non-defective and also the probability that the item was produced by machine A given that it is non-defective.

Problem 2 - Solution

innovate

achieve

lead


Let D : Event that item is defective

A : Event item produced by machine A

B : Event item produced by machine B

C : Event item produced by machine C

ND : Event that item is non-defective



Given that,

$P(A) = 0.50$	$P(D A) = 0.03$	$P(ND A) = 0.97$
$P(B) = 0.30$	$P(D B) = 0.05$	$P(ND B) = 0.95$
$P(C) = 0.20$	$P(D C) = 0.02$	$P(ND C) = 0.98$

By total probability,

$$P(\text{Item found defective}) = P(D)$$

$$= P(A \cap D) + P(B \cap D) + P(C \cap D)$$

$$= P(A)P(D|A) + P(B)P(D|B) + P(C)P(D|C)$$

$$= (0.50)(0.03) + (0.30)(0.05) + (0.20)(0.02)$$

$$= 0.015 + 0.015 + 0.004$$

$$= 0.034$$

$$\therefore P(D) = 0.034$$

(i) To find the probability that the item was produced by machine C:

$$P(C|D) = \frac{P(C \cap D)}{P(D)} = \frac{P(C)P(D|C)}{P(D)}$$

$$= \frac{(0.20)(0.02)}{0.034} = \frac{0.004}{0.034} = 0.1176$$

$$\therefore P(C|D) = 0.1176$$

$\therefore$ Probability that the ^{given} defective item was produced by machine C = 0.1176 (or) 11.76%

Problem 2 - Solution

innovate

achieve

lead

ii) To find the probability that the item was produced by C or B
$$P(C \cup B | D) = P(C/D) + P(B/D)$$

$$P(B/D) = \frac{P(B \cap D)}{P(D)} = \frac{P(B)P(D/B)}{P(D)} = \frac{(0.30)(0.05)}{0.034}$$
$$= \frac{0.015}{0.034} \approx 0.4412$$

$$\therefore P(B/D) = 0.4412$$

$$\therefore P(C \cup B/D) = P(C/D) + P(B/D) = 0.1176 + 0.4412 = 0.5588$$

$\therefore$ Probability that item was produced by machine C or B, given that it is defective is app. 0.5588 (or 55.88%)

iii) By total probability,
$$P(\text{Item found Non-Defective}) = P(ND)$$
$$= P(A \cap ND) + P(B \cap ND) + P(C \cap ND)$$
$$= P(A)P(ND/A) + P(B)P(ND/B) + P(C)P(ND/C)$$
$$= (0.50)(0.97) + (0.30)(0.95) + (0.20)(0.98)$$
$$= 0.485 + 0.285 + 0.196$$
$$= 0.966$$

$$\therefore P(ND) = 0.966$$

To find the probability that the item was produced by machine A:

$$P(A/ND) = \frac{P(A \cap ND)}{P(ND)} = \frac{P(A)P(ND/A)}{P(ND)}$$
$$= \frac{(0.50)(0.97)}{0.966} = \frac{0.485}{0.966} \approx 0.502$$

$$\therefore P(A/ND) = 0.502$$

$\therefore$ Probability that the item was produced by machine A given that it is non-defective is 0.502 (or 50.2%)

Problem 3




Hardik Pandya, Rishabh Pant and Surya Kumar Yadav are in the race of leading Indian cricket team in the next world cup with probabilities 0.2, 0.5 and 0.3 respectively. The probabilities of getting an increase in the match fee by Hardik, Rishabh and Surya are 0.3, 0.6 and 0.5 respectively if they become the Captain.

If there is an increase in match fee then find the probability

- I. that it is because of Hardik
- II. that it is because of Rishabh
- III. that it is because of Surya Kumar

Problem 3 - Solution

Let, Event H: Hardik Pandya becomes the captain
Event R: Rishabh Pant becomes the captain
Event S: Suryakumar Yadav becomes the captain
Event F: There is an increase in the match fee



Given that, $P(H) = 0.2$ $P(F|H) = 0.3$
 $P(R) = 0.5$ $P(F|R) = 0.6$
 $P(S) = 0.3$ $P(F|S) = 0.5$

By Law of total probability, we have

$$\begin{aligned} P(F) &= P(H)P(F|H) + P(R)P(F|R) + P(S)P(F|S) \\ &= (0.2)(0.3) + (0.5)(0.6) + (0.3)(0.5) \\ &= 0.06 + 0.3 + 0.15 = 0.51 \Rightarrow P(F) = 0.51 \end{aligned}$$

(i) To find the probability that it is because of Hardik:

$$P(H|F) = \frac{P(H)P(F|H)}{P(F)} = \frac{(0.2)(0.3)}{0.51} = \frac{0.06}{0.51} = \frac{2}{17} = 0.1176$$

$\therefore P(H|F) = 0.1176$

$\therefore$ The probability that increase in matchfee is because of Hardik = 0.1176 (or) 11.76%

(ii) To find the probability that it is because of Rishabh:

$$P(R|F) = \frac{P(R)P(F|R)}{P(F)} = \frac{(0.5)(0.6)}{0.51} = \frac{0.30}{0.51} = \frac{10}{17} = 0.5882$$

$\therefore P(R|F) = 0.5882$

$\therefore$ The probability that increase in matchfee is because of Rishabh = 0.5882 (or) 58.82%

(iii) To find the probability that it is because of Surya Kumar:

$$P(S|F) = \frac{P(S)P(F|S)}{P(F)} = \frac{(0.3)(0.5)}{0.51} = \frac{0.15}{0.51} = \frac{5}{17} = 0.2941$$

$\therefore P(S|F) = 0.2941$

$\therefore$ The probability that increase in matchfee is because of Suryakumar yadav is = 0.2941 (or) 29.41%

Problem 4



In a manufacturing company Machine 1 produces 40% of the items and Machine 2 and Machine 3 produces 25% and 35% of the items respectively. But from the past records it is found that 15%, 20% and 25% of the items they produce are defective.

Then find

- i) Total percentage of defective items produced
- ii) If a defective item is selected randomly, then find the probability that it is produced by Machine 1
- iii) If a defective item is selected randomly, then find the probability that it is not produced by Machine 2


Problem 4 - Solution

innovate

achieve

lead

Let Event M_1 : Items produced by Machine M_1
Event M_2 : Items produced by Machine M_2
Event M_3 : Items produced by Machine M_3
Event D : Item is defective



Given that

$$P(M_1) = 0.40$$

$$P(M_2) = 0.25$$

$$P(M_3) = 0.35$$

$$P(D|M_1) = 0.15$$

$$P(D|M_2) = 0.20$$

$$P(D|M_3) = 0.25$$

- i) By the law of total probability, we have
- $$P(D) = P(M_1)P(D|M_1) + P(M_2)P(D|M_2) + P(M_3)P(D|M_3)$$
- $$= (0.40)(0.15) + (0.25)(0.20) + (0.35)(0.25)$$

$$= 0.06 + 0.05 + 0.0875 = 0.1975$$
$$\therefore P(D) = 0.1975$$

- ii) To find the probability that it is produced by machine M_1 :

$$P(M_1|D) = \frac{P(M_1)P(D|M_1)}{P(D)} = \frac{(0.40)(0.15)}{0.1975}$$

$$= \frac{0.06}{0.1975} \approx 0.3038$$

$$\therefore P(M_1|D) = 0.3038$$

$\therefore$ Probability that it is produced by Machine M_1 , given that it is defective = 0.3038 (or) 30.38%

- iii) To find the probability that it is not produced by Machine M_2 :

$$P(\bar{M}_2|D) = 1 - P(M_2|D)$$
$$= 1 - \left(\frac{P(M_2)P(D|M_2)}{P(D)} \right)$$


$$= 1 - \left(\frac{(0.25)(0.20)}{0.1975} \right) = 1 - \left(\frac{0.05}{0.1975} \right)$$

$$= 1 - 0.2532 = 0.7468$$

$$\therefore P(\bar{M}_2|D) = 0.7468$$

$\therefore$ The probability that it is not produced by Machine M_2 , given that it is defective = 0.7468 (or) 74.68%

Problem 5



A product is manufactured by four companies A, B, C and D. They produce 20%, 30%, 40% and 10% of the products released in the market. But it is observed that 30%, 20%, 10% and 25% are defective items manufactured by them respectively. Then find


- I. Total percentage of defective items manufactured
- II. The percentage that the selected defective item is produced by company A
- III. The percentage that the selected defective item is not produced by company B
- IV. The percentage that the selected defective item is not produced by company D

Problem 5 - Solution

innovate

achieve

lead



Given that, $P(A) = 0.20$ $P(X/A) = 0.30$

$P(B) = 0.30$ $P(X/B) = 0.20$

$P(C) = 0.40$ $P(X/C) = 0.10$

$P(D) = 0.10$ $P(X/D) = 0.25$

i) By the law of total probability, we have
$$P(X) = P(A)P(X/A) + P(B)P(X/B) + P(C)P(X/C) + P(D)P(X/D)$$

$$\begin{aligned} &= (0.20)(0.30) + (0.30)(0.20) + (0.40)(0.10) + (0.10)(0.25) \\ &= 0.06 + 0.06 + 0.04 + 0.025 = 0.185 \\ &\therefore P(X) = 0.185 \end{aligned}$$

$\therefore$ Total percentage of the defective items manufactures
 $= 18.5\%$

ii) To find the percentage that the selected defective item is produced by company A

$$P(A/X) = \frac{P(A)P(X/A)}{P(X)} = \frac{(0.20)(0.30)}{0.185} = \frac{0.06}{0.185} = 0.3243$$

$$\therefore P(A/X) = 0.3243$$

$\therefore$ The percentage that the selected defective item is produced by company A $= 32.43\%$

Problem 5 - Solution

innovate

achieve

lead

(iii) To find the percentage that the selected defective item is not produced by company B.

$$P(B^c/x) = 1 - P(B/x) = 1 - \left(\frac{P(B)P(x/B)}{P(x)} \right)$$

$$= 1 - \left(\frac{(0.30)(0.20)}{0.185} \right) = 1 - \left(\frac{0.06}{0.185} \right)$$

$$= 1 - 0.3243 = 0.6757$$

$$\therefore P(B^c/x) = 0.6757$$

$\therefore$ The percentage that the selected defective item is not produced by B = 67.57%.

(iv) To find the percentage that the selected defective item is not produced by company D.

$$P(D^c/x) = 1 - P(D/x) = 1 - \left(\frac{P(D)P(x/D)}{P(x)} \right)$$

$$= 1 - \left(\frac{(0.10)(0.25)}{0.185} \right) = 1 - \left(\frac{0.025}{0.185} \right)$$

$$= 1 - 0.1351 = 0.8649$$

$$\therefore P(D^c/x) = 0.8649$$

$\therefore$ The percentage that the selected defective item is not produced by company D = 86.49%.

Naïve Bayes Classifier



- Naïve Bayes classifiers are a collection of algorithms based on Bayes' Theorem. It is not a single algorithm but a family of algorithms where all of them share a common principle, i.e., every pair of features being classified as independent of each other.
- Naïve Bayes classifier is one of the most simple and effective classification algorithms, that aids in the rapid development of machine learning models with rapid prediction capabilities.
- It is highly used in text classification tasks where data contains high dimension (as each word represent one feature in the data). It is fast and making prediction easy with high dimension data.
- It is a probabilistic classifier, that predicts the probability of an instance belongs to a class with a given set of feature value. It uses Bayes theorem in the algorithm for training and prediction.

Why is it called Naïve Bayes?



- The “Naïve” part of the name indicates the simplifying assumption made by the Naïve Bayes Classifier. The classifier assumes that the features used to describe an observation are conditionally independent, given the class label.
-

Assumptions of Naïve Bayes



The fundamental Naive Bayes assumption is that each feature makes an:

- **Feature independence:** The features of the data are conditionally independent of each other, given the class label.
- **Continuous features are normally distributed:** If a feature is continuous, then it is assumed to be normally distributed within each class.
- **Discrete features have multinomial distributions:** If a feature is discrete, then it is assumed to have a multinomial distribution within each class.
- **Features are equally important:** All features are assumed to contribute equally to the prediction of the class label.
- **No missing data:** The data should not contain any missing values.

Types of Naïve Bayes Model



- ❑ **Gaussian Naïve Bayes Classifier** : The Gaussian Naive Bayes classifier assumes that the attributes of a dataset have a normal distribution. Here, if the attributes have continuous values, the classification model assumes that the values are sampled from a Gaussian distribution.
 - ❑ **Multinomial Naïve Bayes**: When the input data is multinomially distributed, we use the multinomial naive Bayes classifier. This algorithm is primarily used for document classification problems like sentiment analysis.
 - ❑ **Bernoulli Naïve Bayes**: The Bernoulli Naive Bayes classification works in a similar manner to the multinomial classification. The difference is that the attributes of the dataset contain boolean values representing the presence or absence of a particular attribute in a data point.
-

Advantages of Naïve Bayes Classifier



- Easy to implement and computationally efficient.
 - Effective in cases with a large number of features.
 - Performs well even with limited training data.
 - It performs well in the presence of categorical features.
 - For numerical features data is assumed to come from normal distributions
-

Disadvantages of Naive Bayes Classifier



- Assumes that features are independent, which may not always hold in real-world data.
 - Can be influenced by irrelevant attributes.
 - May assign zero probability to unseen events, leading to poor generalization.
-

Implementation of Naïve Bayes classifier



To implement a Naive Bayes classifier, we perform three steps.

- First, we calculate the probability of each class label in the training dataset.
 - Next, we calculate the conditional probability of each attribute of the training data for each class label given in the training data.
 - Finally, we use the Bayes theorem and the calculated probabilities to predict class labels for new data points. For this, we will calculate the probability of the new data point belonging to each class. The class with which we get the maximum probability is assigned to the new data point.
-

Problem 6



Consider the following data set:

Find out whether the object with attribute **Confident = Yes, Sick = No** will Fail or Pass using Bayesian Classification.

Confident	Studied	Sick	Result
Yes	No	No	Fail
Yes	No	Yes	Pass
No	Yes	Yes	Fail
No	Yes	No	Pass
Yes	Yes	Yes	Pass

Data set for classification

Problem 6 - Solution

Confident	Studied	Sick	Result	
Yes	No	No	Fail	
Yes	No	Yes	Pass	
No	Yes	Yes	Fail	$P(\text{Pass}) = \frac{3}{5}$
No	Yes	No	Pass	
Yes	Yes	Yes	Pass	$P(\text{Fail}) = \frac{2}{5}$

(Data set for classification)

Let $X = \{\text{Confident} = \text{Yes}, \text{Sick} = \text{No}\}$

Confident \ Result	Pass	Fail
Yes	$\frac{2}{3}$	$\frac{1}{2}$
No	$\frac{1}{3}$	$\frac{1}{2}$

Sick \ Result	Pass	Fail
Yes	$\frac{2}{3}$	$\frac{1}{3}$
No	$\frac{1}{3}$	$\frac{1}{2}$

$$\begin{aligned}
 P(\text{Result} = \text{Pass} / X) &= P(\text{Result} = \text{Pass} / \text{Confident} = \text{Yes}, \text{Sick} = \text{No}) \\
 &= P(\text{Result} = \text{Pass}) \times P(\text{Confident} = \text{Yes} / \text{Result} = \text{Pass}) \times \\
 &\quad P(\text{Sick} = \text{No} / \text{Result} = \text{Pass}) \\
 &= \frac{3}{5} \times \frac{2}{3} \times \frac{1}{3} = \frac{2}{15} = 0.1333
 \end{aligned}$$

$$\therefore P(\text{Result} = \text{Pass} / x) = 0.1333$$

$$\begin{aligned}
 P(\text{Result} = \text{Fail} / x) &= P(\text{Result} = \text{Fail} / \text{Confident} = \text{Yes}, \text{Sick} = \text{No}) \\
 &= P(\text{Result} = \text{Fail}) \times P(\text{Confident} = \text{Yes} / \text{Result} = \text{Fail}) \\
 &\quad \times P(\text{Sick} = \text{No} / \text{Result} = \text{Fail}) \\
 &= \frac{2}{5} \times \frac{1}{2} \times \frac{1}{2} = \frac{1}{10} = 0.1
 \end{aligned}$$

$$\therefore P(\text{Result} = \text{Fail} / x) = 0.1$$

$$\therefore P(\text{Result} = \text{Pass} / x) > P(\text{Result} = \text{Fail} / x)$$

$\therefore$ The new object with attribute X ,
 $X = \{\text{Confident} = \text{Yes}, \text{Sick} = \text{No}\}$ belongs to pass class.

Problem 7

innovate

achieve

lead

Example No.	Color	Type	Origin	Stolen?
1	Red	Sports	Domestic	Yes
2	Red	Sports	Domestic	No
3	Red	Sports	Domestic	Yes
4	Yellow	Sports	Domestic	No
5	Yellow	Sports	Imported	Yes
6	Yellow	SUV	Imported	No
7	Yellow	SUV	Imported	Yes
8	Yellow	SUV	Domestic	No
9	Red	SUV	Imported	No
10	Red	Sports	Imported	Yes

New Instance = (Red, SUV, Domestic) → (Yes or No)

Problem 7 - Solution

innovate

achieve

lead

Let the New instance be

$X = \{ \text{Red, SUV, Domestic} \}$

$$P(\text{Yes}) = \frac{5}{10} = \frac{1}{2}$$

Stolen

$$P(\text{No}) = \frac{5}{10} = \frac{1}{2}$$

Type	Yes	No
Sports	4/5	1/5
SUV	1/5	3/5

Stolen

Colour	Yes	No
Yellow	2/5	3/5
Red	3/5	2/5

Stolen

Origin	Yes	No
Domestic	2/5	3/5
Imported	3/5	2/5

$$P(\text{Yes}/x) = P(\text{Yes}/\text{Red, SUV, Domestic})$$

$$= P(\text{Yes}) \times P(\text{Red}/\text{Yes}) \times P(\text{SUV}/\text{Yes}) \times P(\text{Domestic}/\text{Yes})$$

$$= \frac{1}{2} \times \frac{3}{5} \times \frac{1}{5} \times \frac{2}{5} = \frac{3}{125} = 0.024$$

$$\therefore P(\text{Yes}/\text{Red, SUV, Domestic}) = 0.024$$

$$P(\text{No}/x) = P(\text{No}/\text{Red, SUV, Domestic})$$

$$= P(\text{No}) \times P(\text{Red}/\text{No}) \times P(\text{SUV}/\text{No}) \times P(\text{Domestic}/\text{No})$$

$$= \frac{1}{2} \times \frac{2}{5} \times \frac{3}{5} \times \frac{3}{5} = \frac{9}{125} = 0.072$$

$$\therefore P(\text{No}/\text{Red, SUV, Domestic}) = 0.072$$

$$P(\text{No}/x) > P(\text{Yes}/x)$$

$\therefore$ The new instance belongs to 'No' class



Random Variables

Motivation:

In most statistical problems we are concerned with one number or a few numbers that are associated with the outcomes of experiments.

When inspecting a manufactured product we may be interested only in the number of defectives; in the analysis of a road test we may be interested only in the average speed and the average fuel consumption; and in the study of the performance of a miniature rechargeable battery we may be interested only in its power and life length. All these numbers are associated with situations involving an element of chance—in other words, they are values of random variables.



Random Variables

A random variable is a real valued function defined on the sample space

- The random variable is the observation.
- The random variable is a function of the observation.

A **random variable** consists of an experiment with a probability measure $P[\cdot]$ defined on a sample space S and a function that assigns a real number to each outcome in the sample space of the experiment.

Advantages:

- When the elements of the sample space are non-numeric, they can be quantified by assigning a real number to every event of the sample space. A real number X associated with the outcome of a random experiment.
- Infinite sample space can be converted to finite



Random Variables - Classification

Random variables are usually classified according to the number of values they can assume. Random variable can be classified as

- 1) discrete random variables, which can take on only a finite number
- 2) an uncountable infinity of values; called continuous random variables

Examples:

- **Number of accidents on a highway – Discrete**
- **Weights of the students of the class – Continuous**

Discrete Random Variables – Problem 8

A random variable X can assume five values: 0, 1, 2, 3, 4. A portion of the probability distribution is shown here:

X	0	1	2	3	4
$P(x)$	0.1	0.25	0.25	?	0.2

- Find $p(3)$
- Calculate the mean
- What is the probability that X is greater than 3?
- What is the probability that X is 3 or less?
- What is the probability that X is in between 1 and 3?

Problem 8 - Solution

innovate

achieve

lead

Given probability distⁿ is

X	0	1	2	3	4
P(X=x)	0.1	0.25	0.25	?	0.2

i) We know that, $\sum P(x) = 1$

$$\Rightarrow 0.1 + 0.25 + 0.25 + P(3) + 0.2 = 1$$

$$\Rightarrow 0.8 + P(3) = 1$$

$$\Rightarrow P(3) = 1 - 0.8$$

$$\Rightarrow P(3) = 0.2$$

ii) Mean = $E(X) = \sum x P(x)$

$$= 0(0.1) + 1(0.25) + 2(0.25) + 3(0.2) + 4(0.2)$$

$$= 0 + 0.25 + 0.5 + 0.6 + 0.8$$

$$= 2.15$$

$$\therefore \text{Mean} = 2.15$$


iii) $P(X > 3) = P(X=4) = 0.2$
 $\therefore P(X > 3) = 0.2$

iv) $P(X \leq 3) = P(X=0) + P(X=1) + P(X=2) + P(X=3)$
 $= 0.1 + 0.25 + 0.25 + 0.2$
 $= 0.8$

$$\therefore P(X \leq 3) = 0.8$$

v) $P(1 < X < 3) = P(X=2) = 0.25$
 $\therefore P(1 < X < 3) = 0.25$

Discrete Random Variables – Problem 9



The probabilities that number of times in a day certain computer may malfunction is as follows:

No. of malfunctions in a day	0	1	2	3	4	5	6
Probability	0.17	k	0.27	0.16	0.07	0.03	0.01

Find

(a) the value of k

(b) mean and variance of the malfunction of certain computer

Problem 9 - Solution

8. Given probability distribution is

No. of calls in a day	0	1	2	3	4	5	6
probability	0.17	K	0.27	0.16	0.07	0.03	0.01

(a) We know that $\sum P(x) = 1$

$$\Rightarrow 0.17 + K + 0.27 + 0.16 + 0.07 + 0.03 + 0.01 = 1$$

$$\Rightarrow 0.71 + K = 1$$

$$\Rightarrow K = 1 - 0.71$$

$$\Rightarrow K = 0.29$$

(b) Mean = $E(X) = \sum xP(x)$

$$= 0(0.17) + 1(0.29) + 2(0.27) + 3(0.16) + 4(0.07) + 5(0.03) + 6(0.01)$$

$$= 0 + 0.29 + 0.54 + 0.48 + 0.28 + 0.15 + 0.06$$

$$= 1.8$$

$$\therefore \text{Mean} = 1.8$$

$$\text{Var}(x) = E(x^2) - (E(x))^2$$

$$E(x^2) = \sum x^2 P(x)$$

$$= (0)^2(0.17) + (1)^2(0.29) + (2)^2(0.27) + (3)^2(0.16) + (4)^2(0.07) + (5)^2(0.03) + (6)^2(0.01)$$

$$= 0 + 0.29 + 4(0.27) + 9(0.16) + 16(0.07) + 25(0.03) + 36(0.01)$$

$$= 0.29 + 1.08 + 1.44 + 1.44 + 0.75 + 0.36$$

$$= 5.04$$

$$E(x^2) = 5.04$$

$$\text{Var}(x) = E(x^2) - (E(x))^2$$

$$= 5.04 - (1.8)^2$$

$$= 5.04 - 3.24$$

$$= 1.8$$

$$\therefore \text{Var}(x) = 1.8$$

Random Variable – Probability Distribution - Problem 10

A student takes two courses. In each course, the student will earn a B with probability 0.6 or a C with probability 0.4, independent of the other course. To calculate a grade point average (GPA), a B is worth 3 points and a C is worth 2 points. The student's GPA is the sum of the GPA for each course divided by 2. Make a table of the sample space of the experiment and the corresponding values of the student's GPA, G .

The sample space, probabilities and corresponding grades for the experiment are

Outcome	$P[\cdot]$	G
BB	0.36	3.0
BC	0.24	2.5
CB	0.24	2.5
CC	0.16	2

Continuous Random Variables – Problem 11



A data scientist is working on a machine learning model that predicts house prices based on various features such as size, location, and number of bedrooms. The model has some error in its predictions, represented by the random variable X . The error X follows the following probability density function (PDF):

$$f(x) = \begin{cases} \frac{k}{6} & -3 \leq x \leq 3 \\ 0 & \text{otherwise} \end{cases}$$

Calculate *i)* the value of k , *ii)* the expected value of error *iii)* Compute the variance of the error.

Problem 11 - Solution



i) For a valid probability distribution, the total probability over the range must equal 1.

$$\therefore \int_{-3}^3 f(x) dx = 1 \rightarrow \int_{-3}^3 \frac{k}{6} dx = 1 \rightarrow \frac{k}{6} (3 - (-3)) = 1 \rightarrow k = 1.$$

$$ii) E(X) = \int_{-3}^3 x \cdot \frac{1}{6} dx = \left[\frac{x^2}{12} \right]_{-3}^3 = 0$$

$$iii) Var(x) = E(X^2) - (E(X))^2.$$

$$E(X^2) = \int_{-3}^3 \frac{x^2}{6} dx = \left[\frac{x^3}{18} \right]_{-3}^3 = \frac{1}{6} \left(\frac{27}{3} - \left(-\frac{27}{3} \right) \right) = \frac{18}{6} = 3.$$

$$Var(X) = 3 - (0)^2 = 3.$$

Continuous Random Variables – Problem 12



EC2R
JULY 2024

Let $f(x)$ be a continuous random variable defined in $[0,2]$ as

$f(x) = k(x + 1)/4$, then find

- (a) k value
- (b) mean of x
- (c) Expectation of x^2
- (d) Variance
- (e) $P(0.5 < x < 1.5)$

Problem 12 - Solution



$$a) \int_0^2 \frac{k(x+1)}{4} dx = 1 \Rightarrow \frac{k}{4} \left[\frac{(x+1)^2}{2} \right]_0^2 = 1 \quad K=1$$

$$b) E(x) = \int_0^2 \frac{x(x+1)}{4} dx = \frac{1}{4} \left[\frac{x^3}{3} + \frac{x^2}{2} \right]_0^2 = \frac{7}{6} = 1.1666$$

$$c) E(x^2) = \int_0^2 \frac{x^2(x+1)}{4} dx = \frac{1}{4} \left[\frac{x^4}{4} + \frac{x^3}{3} \right]_0^2 = \frac{5}{3} = 1.6666$$

$$d) Var(x) = E(x^2) - [E(x)]^2 = \frac{11}{36} = 0.3055$$

$$e) P(0.5 < x < 1.5) = \int_{0.5}^{1.5} \frac{x+1}{4} dx \\ = \left[\frac{(x+1)^2}{8} \right]_{0.5}^{1.5} = 0.5$$

Continuous Random Variables – Problem 13



EC2R
JAN 2024

Let $f(x) = k(x + 1)/2$ where $0 < x < 2$,
= 0, otherwise

Then find

i) k value

ii) $P(1/2 < x < 2)$

iii) $P(X > 1/4)$

Problem 13 - Solution



i) To find Value of k:

$$\int_0^2 \frac{k(x+1)}{2} dx = 1$$
$$\frac{k}{2} \left[\frac{x^2}{2} + x \right]_0^2 = 1$$
$$\frac{k}{2} [(2+2) - (0+0)] = 1$$

Therefore $k = \frac{1}{2}$

ii) $P(1/2 < x < 2)$

$$P\left(\frac{1}{2} < x < 2\right) = \int_{1/2}^2 \frac{1}{2} \cdot \frac{(x+1)}{2} dx$$
$$P\left(\frac{1}{2} < x < 2\right) = \frac{1}{4} \left[\frac{x^2}{2} + x \right]_{1/2}^2$$
$$= \frac{27}{32} = 0.84375$$

iii) $P(X > 1/4)$

$$P(X > 1/4) = \frac{1}{4} \left[\frac{x^2}{2} + x \right]_{1/4}^2$$
$$= \frac{119}{128} = 0.9297$$

Joint Probability Distribution Function



- ❖ Let $X = \{x_1, x_2, \dots, x_m\}$ and $Y = \{y_1, y_2, \dots, y_n\}$ be two discrete random variables. Then $P(x, y) = J_{ij}$ is called joint probability function of X and Y if it satisfies the conditions:

$$(i) J_{ij} \geq 0 \quad (ii) \sum_{i=1}^m \sum_{j=1}^n J_{ij} = 1$$

- ❖ Set of values of this joint probability function J_{ij} is called joint probability distribution of X and Y.

	y_1	y_2	...	y_n	Sum
x_1	J_{11}	J_{12}	...	J_{1n}	$f(x_1)$
x_2	J_{21}	J_{22}	...	J_{2n}	$f(x_2)$
...	...	...	...	...	...
x_m	J_{m1}	J_{m2}	...	J_{mn}	$f(x_m)$
Sum	$g(y_1)$	$g(y_2)$	...	$g(y_n)$	Total = 1

Joint Probability Distribution Function



- ❖ If X and Y are discrete random variables, the joint probability distribution of X and Y is a description of the set of points (x,y) in the range of (X,Y) along with the probability of each point.
- ❖ The joint probability distribution of two discrete random variables is sometimes referred to as the **bivariate probability distribution** or **bivariate distribution**.
- ❖ Thus, we can describe the joint probability distribution of two discrete random variables is through a **joint probability mass function**

$$f(x,y) = P(X=x, Y=y)$$

Joint Density Function



When X and Y are continuous random variables, the **joint density function** $f(x, y)$ is a surface lying above the xy plane, and $P[(X, Y) \in A]$, where A is any region in the xy plane, is equal to the volume of the right cylinder bounded by the base A and the surface.

The function $f(x, y)$ is a **joint density function** of the continuous random variables X and Y if

1. $f(x, y) \geq 0$, for all (x, y) ,
2. $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) \, dx \, dy = 1$,
3. $P[(X, Y) \in A] = \int \int_A f(x, y) \, dx \, dy$, for any region A in the xy plane.

Marginal Distributions



The **marginal distributions** of X alone and of Y alone are

$$g(x) = \sum_y f(x, y) \quad \text{and} \quad h(y) = \sum_x f(x, y)$$

for the discrete case, and

$$g(x) = \int_{-\infty}^{\infty} f(x, y) \, dy \quad \text{and} \quad h(y) = \int_{-\infty}^{\infty} f(x, y) \, dx$$

for the continuous case.

Independent Random Variables



□ Let X and Y are two random variables with joint probability function $f(x, y)$ are said to be independent if following condition satisfied:

□ $f(x, y) = g(x) \cdot h(y)$ for all x and y , where $g(x)$ is marginal probability function of X and $h(y)$ is marginal probability function of Y .

Example: Suppose $f(x, y) = e^{-(x+y)}$, $x \geq 0, y \geq 0$. For this probability function,

marginal probability function for X is $g(x) = e^{-x}$, $x \geq 0$ and

marginal probability function for Y is $h(y) = e^{-y}$, $y \geq 0$.

Clearly $f(X, Y) = g(X) \cdot h(Y)$ So X and Y are independent variables.

Conditional Probability Distribution Function

innovate

achieve

lead

- **Discrete Case:**

$$P[X = x_i / Y = y_j] = \frac{P[X = x_i / Y = y_j]}{P[Y = y_j]} = \frac{p_{ij}}{p_{.j}}$$

The set $\{x_i, p_{ij} / p_{.j}\}$, $i = 1, 2, 3, \dots$ is called the conditional probability distribution of X given $Y = y_j$.

The conditional probability function of Y given $X = x_i$ is given by

$$P[Y = y_i / X = x_j] = \frac{P[Y = y_i / X = x_j]}{P[X = x_j]} = \frac{p_{ij}}{p_{i.}}$$

The set $\{y_i, p_{ij} / p_{i.}\}$, $j = 1, 2, 3, \dots$ is called the conditional probability distribution of Y given $X = x_i$.

Conditional Probability Density Function

$$f(y | x) = \frac{f(x, y)}{f_X(x)} \quad \text{or} \quad f(x | y) = \frac{f(x, y)}{f_Y(y)}$$

- the continuous version of Bayes' theorem

$$f(y | x) = \frac{f(x | y)f_Y(y)}{f_X(x)}$$

- another expression of the marginal pdf

$$f_X(x) = \int_{-\infty}^{\infty} f(x, y)dy = \int_{-\infty}^{\infty} f(x | y)f_Y(y)dy$$

$$f_Y(y) = \int_{-\infty}^{\infty} f(x, y)dx = \int_{-\infty}^{\infty} f(y | x)f_X(x)dx$$

Joint Discrete – Problem 14



Consider the following probability distribution of X and Y.

Y	X			
	0	1	2	3
0	0.05	0.25	0.05	0
1	0.05	0.15	0.05	0.05
2	0.10	0.15	0.10	0

- i). Find marginal distribution of X
- ii). Find marginal distribution of Y
- iii). Find $P(X < 2 / Y < 2)$
- iv). Find $P(X < 2 / Y = 1)$

Problem 14 – Solution

innovate

achieve

lead

Solution:

i). Find marginal distribution of X

X	0	1	2	3	Total
P(x)	0.2	0.55	0.2	0.05	1

ii). Find marginal distribution of Y

Y	0	1	2	Total
P(y)	0.35	0.3	0.35	1

iii). Find $P(X < 2 / Y < 2)$

$$\begin{aligned} P(X < 2 | Y < 2) &= \frac{P(X < 2 \cap Y < 2)}{P(Y < 2)} \\ &= \frac{P(X = 0, Y = 0) + P(X = 1, Y = 0) + P(X = 0, Y = 1) + P(X = 1, Y = 1)}{P(Y = 0) + P(Y = 1)} \\ &= \frac{0.05 + 0.05 + 0.25 + 0.15}{0.35 + 0.3} = \frac{0.5}{0.65} = 0.7692 \end{aligned}$$

Problem 14 – Solution



iv). Find $P(X < 2 \mid Y = 1)$

$$\begin{aligned} P(X < 2 \mid Y = 1) &= \frac{P(X < 2 \cap Y = 1)}{P(Y = 1)} \\ &= \frac{P(X = 0, Y = 1) + P(X = 1, Y = 1)}{P(Y = 1)} \\ &= \frac{0.05 + 0.15}{0.3} = \frac{2}{3} = 0.67 \end{aligned}$$

Joint Discrete – Problem 15



The discrete random variables X and Y have joint probability mass function

$$f(x, y) = \begin{cases} cxy & x = 1, 2, 3, \ y = 1, 2 \\ 0 & \text{otherwise} \end{cases}.$$

- i) Find the value of constant c .
- ii) Find the marginal probability function for X and Y .

Problem 15 – Solution

innovate

achieve

lead

$$\text{i) } \sum_x \sum_y cxy = 1$$

$$\therefore c + 2c + 2c + 4c + 3c + 6c = 1$$

$$\therefore 18c = 1$$

$$\therefore c = \frac{1}{18}$$

ii) Let $g(x)$ be marginal probability function of x .

$$\therefore g(x) = \sum_y f(x,y) = \sum_y cxy = cx(1+2) = \frac{3x}{18} = \frac{x}{6}, x = 1,2,3.$$

Let $h(y)$ be marginal probability function for y :

$$h(y) = \sum_x f(x,y) = \sum_x cxy = cy(1+2+3) = 6cy = \frac{6y}{18} = \frac{y}{3}, y = 1,2.$$

Joint Discrete – Problem 16



Let X and Y are two independent random variables with the probability distributions given as

X	-1	0	1	2
$P(X)$	0.25	0.15	0.35	0.25

Y	0	1	2	3
$P(Y)$	0.10	0.20	0.30	0.40

Then find the following

- If possible find the joint distribution of X and Y . If not, justify it.
- If possible ,find $P(x < 1, Y < 2)$
- If possible, find $P(X < 1 / Y < 2)$

Problem 16 – Solution



Part (a): Joint Distribution of X and Y

$P(X, Y)$	$Y = 0$	$Y = 1$	$Y = 2$	$Y = 3$
$X = -1$	0.25×0.10 $= 0.025$	0.25×0.20 $= 0.05$	0.25×0.30 $= 0.075$	0.25×0.40 $= 0.10$
$X = 0$	0.15×0.10 $= 0.015$	0.15×0.20 $= 0.03$	0.15×0.30 $= 0.045$	0.15×0.40 $= 0.06$
$X = 1$	0.35×0.10 $= 0.035$	0.35×0.20 $= 0.07$	0.35×0.30 $= 0.105$	0.35×0.40 $= 0.14$
$X = 2$	0.25×0.10 $= 0.025$	0.25×0.20 $= 0.05$	0.25×0.30 $= 0.075$	0.25×0.40 $= 0.10$

Part (b): $P(X < 1, Y < 2)$

$$\begin{aligned}P(X < 1, Y < 2) &= P(X = -1, Y = 0) + P(X = -1, Y = 1) + P(X = 0, Y = 0) + P(X = 0, Y = 1) \\&= 0.025 + 0.05 + 0.015 + 0.03 \\&= 0.12\end{aligned}$$

Problem 16 – Solution

innovate

achieve

lead

C) $P(X < 1 | Y < 2)$

$$P(X < 1 | Y < 2) = \frac{P(X < 1 \text{ and } Y < 2)}{P(Y < 2)}$$

We already calculated $P(X < 1 \text{ and } Y < 2) = 0.12$.

Next, we need to find $P(Y < 2)$:

$$P(Y < 2) = P(Y = 0) + P(Y = 1) = 0.10 + 0.20 = 0.30$$

Now, we can calculate the conditional probability:

$$\begin{aligned} P(X < 1 | Y < 2) &= \frac{P(X < 1 \text{ and } Y < 2)}{P(Y < 2)} \\ &= \frac{0.12}{0.30} \\ &= 0.4 \end{aligned}$$

Joint Continuous – Problem 17

innovate

create

lead

EC2 M JULY
2024

Consider the probability density function $f(X,Y) = k (X + Y)/8$ where $0 < X < 2$ and $0 < Y < 2$. Then find

- a) k value
- b) Marginal probability density function of X
- c) Marginal probability density function of Y
- d) Are X and Y are independent? Justify it

Problem 17 – Solution



$$a) \int_0^2 \int_0^2 \frac{k(x+y)}{8} dx dy = 1,$$

Solving this with proper steps yield $K = 1$

$$b) f(x) = \int_0^2 \frac{x+y}{8} dy = \frac{x+1}{4}$$

$$c) f(y) = \int_0^2 \frac{x+y}{8} dx = \frac{y+1}{4}$$

d) $f(xy) \neq f(x) \cdot f(y)$. Hence not independent

Joint Continuous – Problem 18

innovate

create

lead

EC2 M FEB
2024

Consider the following probability distribution.

$$f(x, y) = \frac{k}{2} (e^{-(x+y)}, 0 < x < \infty, 0 < y < \infty)$$

- i) Find k value
- ii) Check whether x and y are independent or not.
- iii) Find $P(0 < x < 2, 0 < y < 5)$
- iv) $P(0 < x < 2 / 0 < y < 5)$

Problem 18 – Solution

innovate

achieve

lead

$$\int_0^{\infty} \int_0^{\infty} \frac{k}{2} e^{-(x+y)} dx dy = 1$$

$$\int_0^{\infty} \left(\frac{k}{2} e^{-y} \int_0^{\infty} e^{-x} dx \right) dy = 1$$

$$\int_0^{\infty} \left(\frac{k}{2} e^{-y} (0 - (-1)) \right) dy = 1$$

$$\int_0^{\infty} \frac{k}{2} e^{-y} dy = 1$$

$$\frac{k}{2} [-e^{-y}]_0^{\infty} = 1$$

$$\frac{k}{2} (0 - (-1)) = 1$$

$$\frac{k}{2} = 1$$

$$k = 2$$

i) Check whether x and y are independent or not.

As Joint pdf is $f(x, y) = e^{-(x+y)} = f(x) \cdot f(y)$

And marginal PDF are

$$f(x) = \int_0^{\infty} e^{-(x+y)} dy$$

$$f(y) = \int_0^{\infty} e^{-(x+y)} dx$$

Which gives

$$f(x) = -e^{-x}$$

$$f(y) = -e^{-y}$$

Thus $f(x, y) = f(x) \cdot f(y)$

Hence x and y are independent variables.

ii) Find $P(0 < x < 2, 0 < y < 5)$

$$P(0 < x < 2, 0 < y < 5) = \int_0^2 \int_0^5 e^{-(x+y)} dy dx$$
$$= (e^{-2} - 1) \cdot (e^{-5} - 1) = 0.859$$

iii) $P(0 < x < 2 / 0 < y < 5)$

$$P(0 < x < 2 / 0 < y < 5) = \frac{P(0 < x < 2, 0 < y < 5)}{P(0 < y < 5)} = \frac{0.859}{0.993} = 0.865$$

innovate

achieve

lead

THANK YOU