

Value Iteration Convergence

Last Time

- How do we reason about the **future consequences** of actions in an MDP?
- What are the basic **algorithms for solving MDPs**?

Guiding Questions

- Does value iteration always converge?
- Is the value function unique?

Value Iteration: The Bellman Operator

Algorithm: Value Iteration

while $\|V - V'\|_\infty > \epsilon$

$V \leftarrow V'$

$V' \leftarrow B[V]$

return V'

$$B[V](s) = \max_{a \in A} (R(s, a) + \gamma E[V(s')])$$

Value Iteration Convergence

Theorem 1: Let $\{V_1, \dots, V_\infty\}$ be a sequence of value functions for a discrete MDP generated by the recurrence $V_{k+1} = B[V_k]$. If $\gamma < 1$, then $\lim_{k \rightarrow \infty} V_k = V^*$.

Metrics

Definition: Let M be a set. A *metric* on M is a function $d : M \times M \rightarrow [0, \infty)$ which satisfies the following three conditions for all $x, y, z \in M$:

1. $d(x, y) = 0$ if and only if $x = y$
2. $d(x, y) = d(y, x)$
3. $d(x, y) \leq d(x, z) + d(z, y)$

Contraction Mappings

Definition: A *contraction mapping* on metric space (M, d) is a function $f : M \rightarrow M$ satisfying

$$d(f(x), f(y)) \leq \alpha d(x, y)$$

for some α , $0 \leq \alpha \leq 1$ and all x and y in M .

Definition: x^* is said to be a *fixed point* of f if $f(x^*) = x^*$.

Script: contraction_mapping.jl

Banach's Theorem

Theorem (Banach): If f is a contraction mapping on metric space (M, d) , then

1. f has a single, unique fixed point x^* .
2. If $\{x_k\}$ is a sequence defined by $x_{k+1} = f(x_k)$, then $\lim_{k \rightarrow \infty} x_k = x^*$.

Max Norm

Lemma 1: $(\mathbb{R}^{|S|}, \|\cdot\|_\infty)$ is a metric space.

Definition: Let M be a set. A *metric* on M is a function $d : M \times M \rightarrow [0, \infty)$ which satisfies the following three conditions for all $x, y, z \in M$:

1. $d(x, y) = 0$ if and only if $x = y$
2. $d(x, y) = d(y, x)$
3. $d(x, y) \leq d(x, z) + d(z, y)$

Proof: Note: $\|x - y\|_\infty = \max_i |x_i - y_i|$

$$1. \max |x - y| = 0 \text{ iff } x_i = y_i \quad \forall i$$

$$2. |x - y| = |-(x - y)| = |y - x|$$
$$\therefore \max |x - y| = \max |y - x|$$

$$3. \max |x - z| = \max |x - y + y - z|$$
$$\leq \max(|x - y| + |y - z|)$$
$$\leq \max |x - y| + \max |y - z|$$

Bellman Operator Contraction

Lemma 2: B is a γ contraction mapping on $(\mathbb{R}^{|S|}, \|\cdot\|_\infty)$.

Proof

$$\begin{aligned} \|B[V_1] - B[V_2]\|_\infty &= \max_{s \in S} |B[V_1](s) - B[V_2](s)| \\ &= \max_{s \in S} \left| \max_{a \in A} \left(R(s, a) + \gamma \sum_{s' \in S} T(s'|s, a) V_1(s') \right) - \max_{a \in A} \left(R(s, a) + \gamma \sum_{s' \in S} T(s'|s, a) V_2(s') \right) \right| \\ &\leq \max_{s \in S} \left| \max_{a \in A} \left(R(s, a) + \gamma \sum_{s' \in S} T(s'|s, a) V_1(s') - R(s, a) - \gamma \sum_{s' \in S} T(s'|s, a) V_2(s') \right) \right| \\ &\leq \max_{s \in S, a \in A} \left| \gamma \sum_{s' \in S} T(s'|s, a) (V_1(s') - V_2(s')) \right| \quad | \max(x) | \leq \max |x| \\ &\leq \max_{s \in S, a \in A} \gamma \sum_{s' \in S} T(s'|s, a) |V_1(s') - V_2(s')| \\ &\leq \max_{s \in S, a \in A} \gamma \sum_{s' \in S} T(s'|s, a) \|V_1 - V_2\|_\infty \\ &= \gamma \|V_1 - V_2\|_\infty \max_{s \in S, a \in A} \sum_{s' \in S} T(s'|s, a) \\ &= \gamma \|V_1 - V_2\|_\infty \end{aligned}$$

Value Iteration Convergence

Theorem 1: Let $\{V_1, \dots, V_\infty\}$ be a sequence of value functions for a discrete MDP generated by the recurrence $V_{k+1} = B[V_k]$. If $\gamma < 1$, then $\lim_{k \rightarrow \infty} V_k = V^*$.

Proof:

Lemma 2: B is a γ contraction mapping on $(\mathbb{R}^{|S|}, \|\cdot\|_\infty)$.

Theorem (Banach): If f is a contraction mapping on metric space (M, d) , then

1. f has a single, unique fixed point x^* .
2. If $\{x_k\}$ is a sequence defined by $x_{k+1} = f(x_k)$, then $\lim_{k \rightarrow \infty} x_k = x^*$.

By Lemma 2 and Banach's theorem (part 2), repeated application of the Bellman operator always has a fixed point limit, \hat{V} .

By Banach's theorem (part 1), $\hat{V} = B[\hat{V}]$. Since \hat{V} satisfies Bellman's equation, it is optimal and $\hat{V} = V^*$.

Does Policy Iteration Converge?

Theorem: Policy iteration converges to an optimal policy for a finite MDP in finite time.

Proof (sketch):

1. The policy will either improve or stay the same at each iteration
2. The policy will stay the same if and only if $V^\pi = V^*$
3. There are a finite number of possible policies
4. By (1), (2), and (3), the policy will improve until it finds the optimal policy, and it will always find the optimal policy.

Guiding Questions

- Does value iteration always converge?
- Is the value function unique?

Break

Conservation MDP

