Supporting Information for Synoptic Pathways of Mid-depth Salinity Maximum Intrusions in the South Brazil Bight

D. A. Silva[1], A. Gangopadhyay[2], D. K. Sasaki[1,3], M. Dottori[1], I. C. A. da Silveira[1], W. C. Belo[4], R. P. Martins[4]

[1]Instituto Oceanográfico, Universidade de Sao Paulo, Praca do Oceanográfico, 191 - SP 05508-120, São Paulo, Brasil

[2]School for Marine Sciences and Technology, University of Massachusetts, 836 South Rodney French Blvd, New Bedford, MA 02744-1221

[3]Centro de Pesquisas, Desenvolvimento e Inovação Leopoldo Américo Miguez de Mello - CENPES/PETROBRAS, Av. Horácio Macedo, 950 - Cidade Universitária da Universidade Federal do Rio de Janeiro, Rio de Janeiro - RJ, 21941-915

[4]Department of Marine and Environmental Sciences, Northeastern University, 430 Nahant Rd, Nahant, MA 01908

Contents of this file

1. Figures S1 to S3

Additional Supporting Information (Files uploaded separately)

1. Captions for Movie V1




Figure S1. (a) Location of the virtual along-shelf edge transect, represented by a straight line over the shelf edge. (b) time by latitude plot, with cross-shelf edge geostrophic velocities, in $m\ s^{-1}$. Blue (negative) values are shoreward and red (positive) offshoreward. The method description for this analysis is in Section 2.5.1.




Figure S2. Vertical profiles of practical salinity extracted from the GLORYS12V1 reanalysis locations of interested (a-g), depicted in subpanel (h). The profiles are colored by the date of occurrence, from January to May of 2022.

1. Lagrangian analysis - Forward propagation experiments

From our Lagrangian analysis, we have found that onlye particles released at north of the region of interest reach the shelf, via two pathways. To further investigate this advection, we used the FPN experiment to understand the relative importance of each pathway. To this end, we assumed that particles advected by the alongshelf pathway would reach the shelf area (a square box between Guanarabara and Cabo Frio, Figure S3) faster than those advected by a cross-shelf pathway. The rationale for this assumption is that the alongshelf pathway, associated with the BC non-meandering state, would provide a more direct route to the region of interest, whereas the cross-shelf pathway would involve more complex interactions with mesoscale features, such as eddies, potentially slowing transport. We have found that 31% of the particles that reached the shelf area did so within the first 10 days of the simulation, on average, covering an average cumulative distance of 335 km, suggesting they were likely advected along the shelf. In contrast, 21% of the particles reached the shelf area after 20 days on average, with a cumulative distance covered of 541 km, indicating they were likely transported via a cross-shelf pathway (Figure S3 a). The alongshelf pathway predominantly enters the region of interest through the eastern boundary (100%), whereas the cross-shelf pathway particles enter through both the eastern (84%) and southern (16%) boundaries (Figure S3 b). Particles with a longer pathway spend significantly more time under high EKE activity than those particles with a fast advection (Mann-Whitney Test, p-value < 0.001, Figure S3 c). We also run additional experiments releasing particles along the isopycnal, similar to BPII, but for forward propagation (not shown). The results were consistent with FPN, with particles

reaching the shelf region only when released from the north, following both pathways described above.




Figure S3. (a) Fast (blue) and slow (orange) regimes pathways of particles that entered the region of interest (dashed black square), for the experiment FPN only. Black dots indicate the location of the sequence of profiles selected on the main analysis. Note we have extended the region of interest further north to consider particles advected over the shelf and along the shelfbreak and south to consider the shelf break, since in this last case particles drifting along the shelf might be injected into the shelf. (b) Percentage of particles entering the region of interest through each boundary, computed considering only particles that effectively enter the box and separated by pathway, the colors follow the same from (a). (c) Fraction of time particles were found under high EKE activity for

the fast (left) and slow (right) regimes. The Mann-Whitney U-Test indicate particles on the slow regime are under eddy activity influence.

Video V1. Practical salinity horizontal fields from GLORYS12V1 (Jean-Michel et al., 2021) in colors, bathymetry from GEBCO2019 (Weatherall et al., 2015) is thin gray lines and 36 and 37.3 PSU isohalines in thicker black lines. (a) is the horizontal field at th 50-m layer and (b) is the salinity horizontal distribution projected onto the 25.5 kg m-3 isopycnal surface, with the selected profiles, presented in SectionXXX in the manuscript, in black scatters. The video presents the salinity evolution from January 1st to May 31st of 2022.

do not specify file extension