

Simulation of Proof Systems

Bachelor's Thesis

Nils Wisiol

Lehrstuhl für Informatik IV
Institut für Informatik
Julius-Maximilians-Universität Würzburg

04. Juni 2012

Outline

Preliminaries

Languages with Optimal Proof Systems

Languages without Optimal Proof Systems

Consequences

Literature

Optimal Proof Systems

A \mathcal{FP} -function h is called *proof system* for a language L , if $f(\Sigma^*) = L$. If $h(w) = x$ holds, we call w a *h -proof* for x .

Let h and h' be proof systems for a language L , and let p be a polynomial and a let f be a function such that for all h' -proofs w it holds that

$$h(f(w)) = h'(w),$$

where $|f(w)| \leq p(|w|)$ holds. In this case, we say h *simulates* the proof system h' .

If a proof system h for a language L simulates every other proof system for the same language L , we call it *optimal*. Let OPT be the complexity class of all languages possessing a optimal proof system.

Optimal Proof Systems

A \mathcal{FP} -function h is called *proof system* for a language L , if $f(\Sigma^*) = L$. If $h(w) = x$ holds, we call w a *h -proof* for x .

Let h and h' be proof systems for a language L , and let p be a polynomial and a let f be a function such that for all h' -proofs w it holds that

$$h(f(w)) = h'(w),$$

where $|f(w)| \leq p(|w|)$ holds. In this case, we say h *simulates* the proof system h' .

If a proof system h for a language L simulates every other proof system for the same language L , we call it *optimal*. Let OPT be the complexity class of all languages possessing a optimal proof system.

Optimal Proof Systems

A \mathcal{FP} -function h is called *proof system* for a language L , if $f(\Sigma^*) = L$. If $h(w) = x$ holds, we call w a *h -proof* for x .

Let h and h' be proof systems for a language L , and let p be a polynomial and a let f be a function such that for all h' -proofs w it holds that

$$h(f(w)) = h'(w),$$

where $|f(w)| \leq p(|w|)$ holds. In this case, we say h *simulates* the proof system h' .

If a proof system h for a language L simulates every other proof system for the same language L , we call it *optimal*. Let OPT be the complexity class of all languages possessing a optimal proof system.

Optimal Proof Systems

A \mathcal{FP} -function h is called *proof system* for a language L , if $f(\Sigma^*) = L$. If $h(w) = x$ holds, we call w a *h -proof* for x .

Let h and h' be proof systems for a language L , and let p be a polynomial and a let f be a function such that for all h' -proofs w it holds that

$$h(f(w)) = h'(w),$$

where $|f(w)| \leq p(|w|)$ holds. In this case, we say h *simulates* the proof system h' .

If a proof system h for a language L simulates every other proof system for the same language L , we call it *optimal*. Let OPT be the complexity class of all languages possessing a optimal proof system.

Optimal Proof Systems

A \mathcal{FP} -function h is called *proof system* for a language L , if $f(\Sigma^*) = L$. If $h(w) = x$ holds, we call w a *h -proof* for x .

Let h and h' be proof systems for a language L , and let p be a polynomial and a let f be a function such that for all h' -proofs w it holds that

$$h(f(w)) = h'(w),$$

where $|f(w)| \leq p(|w|)$ holds. In this case, we say h *simulates* the proof system h' .

If a proof system h for a language L simulates every other proof system for the same language L , we call it *optimal*. Let OPT be the complexity class of all languages possessing a optimal proof system.

Optimal Proof Systems

A \mathcal{FP} -function h is called *proof system* for a language L , if $f(\Sigma^*) = L$. If $h(w) = x$ holds, we call w a h -proof for x .

Let h and h' be proof systems for a language L , and let p be a polynomial and let f be a function such that for all h' -proofs w it holds that

$$h(f(w)) = h'(w),$$

where $|f(w)| \leq p(|w|)$ holds. In this case, we say h *simulates* the proof system h' .

If a proof system h for a language L simulates every other proof system for the same language L , we call it *optimal*. Let OPT be the complexity class of all languages possessing an optimal proof system.

Optimal Proof Systems

A \mathcal{FP} -function h is called *proof system* for a language L , if $f(\Sigma^*) = L$. If $h(w) = x$ holds, we call w a *h -proof* for x .

Let h and h' be proof systems for a language L , and let p be a polynomial and a let f be a function such that for all h' -proofs w it holds that

$$h(f(w)) = h'(w),$$

where $|f(w)| \leq p(|w|)$ holds. In this case, we say h *simulates* the proof system h' .

If a proof system h for a language L simulates every other proof system for the same language L , we call it *optimal*. Let OPT be the complexity class of all languages possessing a optimal proof system.

Optimal Proof Systems

A \mathcal{FP} -function h is called *proof system* for a language L , if $f(\Sigma^*) = L$. If $h(w) = x$ holds, we call w a *h -proof* for x .

Let h and h' be proof systems for a language L , and let p be a polynomial and let f be a function such that for all h' -proofs w it holds that

$$h(f(w)) = h'(w),$$

where $|f(w)| \leq p(|w|)$ holds. In this case, we say h *simulates* the proof system h' .

If a proof system h for a language L simulates every other proof system for the same language L , we call it *optimal*. Let OPT be the complexity class of all languages possessing a optimal proof system.

Optimal Proof Systems

A \mathcal{FP} -function h is called *proof system* for a language L , if $f(\Sigma^*) = L$. If $h(w) = x$ holds, we call w a *h -proof* for x .

Let h and h' be proof systems for a language L , and let p be a polynomial and let f be a function such that for all h' -proofs w it holds that

$$h(f(w)) = h'(w),$$

where $|f(w)| \leq p(|w|)$ holds. In this case, we say h *simulates* the proof system h' .

If a proof system h for a language L simulates every other proof system for the same language L , we call it *optimal*. Let OPT be the complexity class of all languages possessing a optimal proof system.

Languages with Optimal Proof Systems

- ▶ $P \subseteq \text{OPT}$: $L \in P$ has a proof system f : choose $f : \Sigma^* \rightarrow L$ with $f(x) = x$ if $x \in L$, otherwise f is undefined. For any other proof system g of L , g itself translates g -proofs into f -proofs in polynomial time.
- ▶ $\text{NP} \subseteq \text{OPT}$: For any $L \in \text{NP}$, there is a nondet. TM that accepts L in poly. time. Let $f(\langle x, i \rangle)$ be the function defined by

$$f(\langle x, i \rangle) = \begin{cases} x & \text{if } M \text{ accepts } x \text{ on the } i\text{-th path,} \\ \perp & \text{otherwise.} \end{cases}$$

f can be calculated in polynomial time and f has range L . Hence, f is a proof system for L . Let g be an arbitrary proof system for L . Then g itself translates g -proofs into f -proofs. Hence, f is an optimal proof system for L .

Languages with Optimal Proof Systems

- ▶ $P \subseteq \text{OPT}$: $L \in P$ has a proof system f : choose $f : \Sigma^* \rightarrow L$ with $f(x) = x$ if $x \in L$, otherwise f is undefined. For any other proof system g of L , g itself translates g -proofs into f -proofs in polynomial time.
- ▶ $\text{NP} \subseteq \text{OPT}$: For any $L \in \text{NP}$, there is a nondet. TM that accepts L in poly. time. Let $f(\langle x, i \rangle)$ be the function defined by

$$f(\langle x, i \rangle) = \begin{cases} x & \text{if } M \text{ accepts } x \text{ on the } i\text{-th path,} \\ \perp & \text{otherwise.} \end{cases}$$

f can be calculated in polynomial time and f has range L . Hence, f is a proof system for L . Let g be an arbitrary proof system for L . Then g itself translates g -proofs into f -proofs. Hence, f is an optimal proof system for L .

Languages with Optimal Proof Systems

- ▶ $P \subseteq \text{OPT}$: $L \in P$ has a proof system f : choose $f : \Sigma^* \rightarrow L$ with $f(x) = x$ if $x \in L$, otherwise f is undefined. For any other proof system g of L , g itself translates g -proofs into f -proofs in polynomial time.
- ▶ $\text{NP} \subseteq \text{OPT}$: For any $L \in \text{NP}$, there is a nondet. TM that accepts L in poly. time. Let $f(\langle x, i \rangle)$ be the function defined by

$$f(\langle x, i \rangle) = \begin{cases} x & \text{if } M \text{ accepts } x \text{ on the } i\text{-th path,} \\ \perp & \text{otherwise.} \end{cases}$$

f can be calculated in polynomial time and f has range L . Hence, f is a proof system for L . Let g be an arbitrary proof system for L . Then g itself translates g -proofs into f -proofs. Hence, f is an optimal proof system for L .

Languages with Optimal Proof Systems

- ▶ $P \subseteq \text{OPT}$: $L \in P$ has a proof system f : choose $f : \Sigma^* \rightarrow L$ with $f(x) = x$ if $x \in L$, otherwise f is undefined. For any other proof system g of L , g itself translates g -proofs into h -proofs in polynomial time.
- ▶ $\text{NP} \subseteq \text{OPT}$: For any $L \in \text{NP}$, there is a nondet. TM that accepts L in poly. time. Let $f(\langle x, i \rangle)$ be the function defined by

$$f(\langle x, i \rangle) = \begin{cases} x & \text{if } M \text{ accepts } x \text{ on the } i\text{-th path,} \\ \perp & \text{otherwise.} \end{cases}$$

f can be calculated in polynomial time and f has range L . Hence, f is a proof system for L . Let g be an arbitrary proof system for L . Then g itself translates g -proofs into f -proofs. Hence, f is an optimal proof system for L .

Languages with Optimal Proof Systems

- ▶ $P \subseteq \text{OPT}$: $L \in P$ has a proof system f : choose $f : \Sigma^* \rightarrow L$ with $f(x) = x$ if $x \in L$, otherwise f is undefined. For any other proof system g of L , g itself translates g -proofs into h -proofs in polynomial time.
- ▶ $\text{NP} \subseteq \text{OPT}$: For any $L \in \text{NP}$, there is a nondet. TM that accepts L in poly. time. Let $f(\langle x, i \rangle)$ be the function defined by

$$f(\langle x, i \rangle) = \begin{cases} x & \text{if } M \text{ accepts } x \text{ on the } i\text{-th path,} \\ \perp & \text{otherwise.} \end{cases}$$

f can be calculated in polynomial time and f has range L . Hence, f is a proof system for L . Let g be an arbitrary proof system for L . Then g itself translates g -proofs into f -proofs. Hence, f is an optimal proof system for L .

Languages with Optimal Proof Systems

- ▶ $P \subseteq \text{OPT}$: $L \in P$ has a proof system f : choose $f : \Sigma^* \rightarrow L$ with $f(x) = x$ if $x \in L$, otherwise f is undefined. For any other proof system g of L , g itself translates g -proofs into h -proofs in polynomial time.
- ▶ $\text{NP} \subseteq \text{OPT}$: For any $L \in \text{NP}$, there is a nondet. TM that accepts L in poly. time. Let $f(\langle x, i \rangle)$ be the function defined by

$$f(\langle x, i \rangle) = \begin{cases} x & \text{if } M \text{ accepts } x \text{ on the } i\text{-th path,} \\ \perp & \text{otherwise.} \end{cases}$$

f can be calculated in polynomial time and f has range L . Hence, f is a proof system for L . Let g be an arbitrary proof system for L . Then g itself translates g -proofs into f -proofs. Hence, f is an optimal proof system for L .

Languages with Optimal Proof Systems

- ▶ $P \subseteq \text{OPT}$: $L \in P$ has a proof system f : choose $f : \Sigma^* \rightarrow L$ with $f(x) = x$ if $x \in L$, otherwise f is undefined. For any other proof system g of L , g itself translates g -proofs into h -proofs in polynomial time.
- ▶ $\text{NP} \subseteq \text{OPT}$: For any $L \in \text{NP}$, there is a nondet. TM that accepts L in poly. time. Let $f(\langle x, i \rangle)$ be the function defined by

$$f(\langle x, i \rangle) = \begin{cases} x & \text{if } M \text{ accepts } x \text{ on the } i\text{-th path,} \\ \perp & \text{otherwise.} \end{cases}$$

f can be calculated in polynomial time and f has range L . Hence, f is a proof system for L . Let g be an arbitrary proof system for L . Then g itself translates g -proofs into f -proofs. Hence, f is an optimal proof system for L .

Languages with Optimal Proof Systems

- ▶ $P \subseteq \text{OPT}$: $L \in P$ has a proof system f : choose $f : \Sigma^* \rightarrow L$ with $f(x) = x$ if $x \in L$, otherwise f is undefined. For any other proof system g of L , g itself translates g -proofs into f -proofs in polynomial time.
- ▶ $\text{NP} \subseteq \text{OPT}$: For any $L \in \text{NP}$, there is a nondet. TM that accepts L in poly. time. Let $f(\langle x, i \rangle)$ be the function defined by

$$f(\langle x, i \rangle) = \begin{cases} x & \text{if } M \text{ accepts } x \text{ on the } i\text{-th path,} \\ \perp & \text{otherwise.} \end{cases}$$

f can be calculated in polynomial time and f has range L . Hence, f is a proof system for L . Let g be an arbitrary proof system for L . Then g itself translates g -proofs into f -proofs. Hence, f is an optimal proof system for L .

Languages with Optimal Proof Systems

- ▶ $P \subseteq \text{OPT}$: $L \in P$ has a proof system f : choose $f : \Sigma^* \rightarrow L$ with $f(x) = x$ if $x \in L$, otherwise f is undefined. For any other proof system g of L , g itself translates g -proofs into f -proofs in polynomial time.
- ▶ $\text{NP} \subseteq \text{OPT}$: For any $L \in \text{NP}$, there is a nondet. TM that accepts L in poly. time. Let $f(\langle x, i \rangle)$ be the function defined by

$$f(\langle x, i \rangle) = \begin{cases} x & \text{if } M \text{ accepts } x \text{ on the } i\text{-th path,} \\ \perp & \text{otherwise.} \end{cases}$$

f can be calculated in polynomial time and f has range L . Hence, f is a proof system for L . Let g be an arbitrary proof system for L . Then g itself translates g -proofs into f -proofs. Hence, f is an optimal proof system for L .

Languages without Optimal Proof Systems

Theorem

There exists a language $L \in \text{co-NTIME}(2^n)$, that does not possess an optimal proof system.

Proof Overview

1. Let f_1, f_2, \dots be an enumeration of all polynomial time functions

2. $L_i = 0^i 10^*$

Let L'_i be the language of all strings L_i without any “short” f_i -proof

$$L = \bigcup_i L'_i \in \text{co-NTIME}(2^n)$$

3. We will show that for L -proof-systems f_i it holds $L'_i = L_i$. As a consequence, there are only long f_i -proofs for $L'_i \subset L$
4. This will contradict to the assumption, that f_i is an optimal proof system for L .

Proof Overview

1. Let f_1, f_2, \dots be an enumeration of all polynomial time functions

2. $L_i = 0^i 10^*$

Let L'_i be the language of all strings L_i without any “short” f_i -proof

$$L = \bigcup_i L'_i \in \text{co-NTIME}(2^n)$$

3. We will show that for L -proof-systems f_i it holds $L'_i = L_i$. As a consequence, there are only long f_i -proofs for $L'_i \subset L$
4. This will contradict to the assumption, that f_i is an optimal proof system for L .

Proof Overview

1. Let f_1, f_2, \dots be an enumeration of all polynomial time functions

2. $L_i = 0^i 10^*$

Let L'_i be the language of all strings L_i without any “short” f_i -proof

$$L = \bigcup_i L'_i \in \text{co-NTIME}(2^n)$$

3. We will show that for L -proof-systems f_i it holds $L'_i = L_i$. As a consequence, there are only long f_i -proofs for $L'_i \subset L$
4. This will contradict to the assumption, that f_i is an optimal proof system for L .

Proof Overview

1. Let f_1, f_2, \dots be an enumeration of all polynomial time functions
2. $L_i = 0^i 10^*$
Let L'_i be the language of all strings L_i without any “short” f_i -proof
 $L = \bigcup_i L'_i \in \text{co-NTIME}(2^n)$
3. We will show that for L -proof-systems f_i it holds $L'_i = L_i$. As a consequence, there are only long f_i -proofs for $L'_i \subset L$
4. This will contradict to the assumption, that f_i is an optimal proof system for L .

Proof Overview

1. Let f_1, f_2, \dots be an enumeration of all polynomial time functions
2. $L_i = 0^i 10^*$
Let L'_i be the language of all strings L_i without any “short” f_i -proof
 $L = \bigcup_i L'_i \in \text{co-NTIME}(2^n)$
3. We will show that for L -proof-systems f_i it holds $L'_i = L_i$. As a consequence, there are only long f_i -proofs for $L'_i \subset L$
4. This will contradict to the assumption, that f_i is an optimal proof system for L .

Proof Overview

1. Let f_1, f_2, \dots be an enumeration of all polynomial time functions
2. $L_i = 0^i 10^*$
Let L'_i be the language of all strings L_i without any “short” f_i -proof
 $L = \bigcup_i L'_i \in \text{co-NTIME}(2^n)$
3. We will show that for L -proof-systems f_i it holds $L'_i = L_i$. As a consequence, there are only long f_i -proofs for $L'_i \subset L$
4. This will contradict to the assumption, that f_i is an optimal proof system for L .

Enumerating \mathcal{FP}

- ▶ Gödel: M_1, M_2, \dots
- ▶ We define M'_1, M'_2, \dots as the M_i with a clock that stops the calculation after $n^i + i$ steps
- ▶ Let f_i the function calculated by M_i
- ▶ As for unbounded i , the runtime $n^i + i$ is unbounded, we obtain all \mathcal{FP} functions



Kurt Gödel
1906 – 1978

Enumerating \mathcal{FP}

- ▶ Gödel: M_1, M_2, \dots
- ▶ We define M'_1, M'_2, \dots as the M_i with a clock that stops the calculation after $n^i + i$ steps
- ▶ Let f_i the function calculated by M_i
- ▶ As for unbounded i , the runtime $n^i + i$ is unbounded, we obtain all \mathcal{FP} functions



Kurt Gödel
1906 – 1978

Enumerating \mathcal{FP}

- ▶ Gödel: M_1, M_2, \dots
- ▶ We define M'_1, M'_2, \dots as the M_i with a clock that stops the calculation after $n^i + i$ steps
- ▶ Let f_i the function calculated by M_i
- ▶ As for unbounded i , the runtime $n^i + i$ is unbounded, we obtain all \mathcal{FP} functions



Kurt Gödel
1906 – 1978

Enumerating \mathcal{FP}

- ▶ Gödel: M_1, M_2, \dots
- ▶ We define M'_1, M'_2, \dots as the M_i with a clock that stops the calculation after $n^i + i$ steps
- ▶ Let f_i the function calculated by M_i
- ▶ As for unbounded i , the runtime $n^i + i$ is unbounded, we obtain all \mathcal{FP} functions



Kurt Gödel
1906 – 1978

Enumerating \mathcal{FP}

- ▶ Gödel: M_1, M_2, \dots
- ▶ We define M'_1, M'_2, \dots as the M_i with a clock that stops the calculation after $n^i + i$ steps
- ▶ Let f_i the function calculated by M_i
- ▶ As for unbounded i , the runtime $n^i + i$ is unbounded, we obtain all \mathcal{FP} functions



Kurt Gödel
1906 – 1978

Construction of L

- ▶ $L_i = 0^i 10^*$
- ▶ take $x \in L'_i$, that do not have f_i -proofs

$$L'_i = \{x \in L_i : \forall_{y \in \Sigma^*} |y|^{2^i} \leq 2^{|x|} \implies f_i(y) \neq x\}$$

- ▶ union

$$L = \bigcup_{i \geq 0} L'_i$$

Construction of L

- ▶ $L_i = 0^i 10^*$
- ▶ take $x \in L'_i$, that do not have f_i -proofs

$$L'_i = \{x \in L_i : \forall_{y \in \Sigma^*} |y|^{2^i} \leq 2^{|x|} \implies f_i(y) \neq x\}$$

- ▶ union

$$L = \bigcup_{i \geq 0} L'_i$$

Construction of L

- ▶ $L_i = 0^i 10^*$
- ▶ take $x \in L'_i$, that do not have f_i -proofs

$$L'_i = \{x \in L_i : \forall_{y \in \Sigma^*} |y|^{2^i} \leq 2^{|x|} \implies f_i(y) \neq x\}$$

- ▶ union

$$L = \bigcup_{i > 0} L'_i$$

L is member of $\text{co-NTIME}(2^n)$

- ▶ As $L \in \text{co-NTIME}(2^n) \Leftrightarrow \bar{L} \in \text{NTIME}(2^n)$, we will analyze the complexity of \bar{L} .
- ▶ $\bar{L} = \overline{\bigcup_{i>0} L_i} = \bigcap_{i>0} \bar{L}_i$
- ▶ claim: $\bigcap_{i>0} \bar{L}_i \in \text{NTIME}(2^n)$

$$\bar{L}_i = \{x \in \Sigma^* : x \notin L_i \vee (\exists y \in \Sigma^* (|y|^{2^i} \leq 2^{|x|}) \wedge (f_i(y) = x))\}$$

Let x be an arbitrary string.

- ▶ Check, if x is member of any L_i : if not, then $x \in \bar{L}$
- ▶ Otherwise, choose i^* such that $x \in L_{i^*}$
- ▶ $x \in \bar{L}_j$ for any $j \neq i^*$
- ▶ for any y such that $|y|^{2^i} \leq 2^{|x|}$: calculate $f_{i^*}(y)$. If and only if there is a y such that $f_{i^*}(y) = x$, then $x \in \bar{L}$.

L is member of $\text{co-NTIME}(2^n)$

- ▶ As $L \in \text{co-NTIME}(2^n) \Leftrightarrow \bar{L} \in \text{NTIME}(2^n)$, we will analyze the complexity of \bar{L} .
- ▶ $\bar{L} = \overline{\bigcup_{i>0} L_i} = \bigcap_{i>0} \bar{L}_i$
- ▶ claim: $\bigcap_{i>0} \bar{L}_i \in \text{NTIME}(2^n)$

$$\bar{L}_i = \{x \in \Sigma^* : x \notin L_i \vee (\exists y \in \Sigma^* (|y|^{2^i} \leq 2^{|x|}) \wedge (f_i(y) = x))\}$$

Let x be an arbitrary string.

- ▶ Check, if x is member of any L_i : if not, then $x \in \bar{L}$
- ▶ Otherwise, choose i^* such that $x \in L_{i^*}$
- ▶ $x \in \bar{L}_j$ for any $j \neq i^*$
- ▶ for any y such that $|y|^{2^i} \leq 2^{|x|}$: calculate $f_{i^*}(y)$. If and only if there is a y such that $f_{i^*}(y) = x$, then $x \in \bar{L}$.

L is member of $\text{co-NTIME}(2^n)$

- ▶ As $L \in \text{co-NTIME}(2^n) \Leftrightarrow \bar{L} \in \text{NTIME}(2^n)$, we will analyze the complexity of \bar{L} .
- ▶ $\bar{L} = \overline{\bigcup_{i>0} L_i} = \bigcap_{i>0} \bar{L}_i$
- ▶ claim: $\bigcap_{i>0} \bar{L}_i \in \text{NTIME}(2^n)$

$$\bar{L}_i = \{x \in \Sigma^* : x \notin L_i \vee (\exists y \in \Sigma^* (|y|^{2i} \leq 2^{|x|}) \wedge (f_i(y) = x))\}$$

Let x be an arbitrary string.

- ▶ Check, if x is member of any L_i : if not, then $x \in \bar{L}$
- ▶ Otherwise, choose i^* such that $x \in L_{i^*}$
- ▶ $x \in \bar{L}_j$ for any $j \neq i^*$
- ▶ for any y such that $|y|^{2i} \leq 2^{|x|}$: calculate $f_{i^*}(y)$. If and only if there is a y such that $f_{i^*}(y) = x$, then $x \in \bar{L}$.

L is member of $\text{co-NTIME}(2^n)$

- ▶ As $L \in \text{co-NTIME}(2^n) \Leftrightarrow \bar{L} \in \text{NTIME}(2^n)$, we will analyze the complexity of \bar{L} .
- ▶ $\bar{L} = \overline{\bigcup_{i>0} L'_i} = \bigcap_{i>0} \bar{L}'_i$
 - ▶ claim: $\bigcap_{i>0} \bar{L}'_i \in \text{NTIME}(2^n)$

$$\bar{L}'_i = \{x \in \Sigma^* : x \notin L_i \vee (\exists y \in \Sigma^* (|y|^{2^i} \leq 2^{|x|}) \wedge (f_i(y) = x))\}$$

Let x be an arbitrary string.

- ▶ Check, if x is member of any L_i : if not, then $x \in \bar{L}$
- ▶ Otherwise, choose i^* such that $x \in L_{i^*}$
- ▶ $x \in \bar{L}'_j$ for any $j \neq i^*$
- ▶ for any y such that $|y|^{2^i} \leq 2^{|x|}$: calculate $f_{i^*}(y)$. If and only if there is a y such that $f_{i^*}(y) = x$, then $x \in \bar{L}$.

L is member of $\text{co-NTIME}(2^n)$

- ▶ As $L \in \text{co-NTIME}(2^n) \Leftrightarrow \bar{L} \in \text{NTIME}(2^n)$, we will analyze the complexity of \bar{L} .
- ▶ $\bar{L} = \overline{\bigcup_{i>0} L'_i} = \bigcap_{i>0} \bar{L}'_i$
- ▶ claim: $\bigcap_{i>0} \bar{L}'_i \in \text{NTIME}(2^n)$

$$\bar{L}'_i = \{x \in \Sigma^* : x \notin L_i \vee (\exists y \in \Sigma^* (|y|^{2^i} \leq 2^{|x|}) \wedge (f_i(y) = x))\}$$

Let x be an arbitrary string.

- ▶ Check, if x is member of any L_i : if not, then $x \in \bar{L}$
- ▶ Otherwise, choose i^* such that $x \in L_{i^*}$
- ▶ $x \in \bar{L}'_j$ for any $j \neq i^*$
- ▶ for any y such that $|y|^{2^i} \leq 2^{|x|}$: calculate $f_{i^*}(y)$. If and only if there is a y such that $f_{i^*}(y) = x$, then $x \in \bar{L}$.

L is member of $\text{co-NTIME}(2^n)$

- ▶ As $L \in \text{co-NTIME}(2^n) \Leftrightarrow \bar{L} \in \text{NTIME}(2^n)$, we will analyze the complexity of \bar{L} .
- ▶ $\bar{L} = \overline{\bigcup_{i>0} L'_i} = \bigcap_{i>0} \bar{L}'_i$
- ▶ claim: $\bigcap_{i>0} \bar{L}'_i \in \text{NTIME}(2^n)$

$$\bar{L}'_i = \{x \in \Sigma^* : x \notin L_i \vee (\exists y \in \Sigma^* (|y|^{2^i} \leq 2^{|x|}) \wedge (f_i(y) = x))\}$$

Let x be an arbitrary string.

- ▶ Check, if x is member of any L_i : if not, then $x \in \bar{L}$
- ▶ Otherwise, choose i^* such that $x \in L_{i^*}$
- ▶ $x \in \bar{L}'_j$ for any $j \neq i^*$
- ▶ for any y such that $|y|^{2^i} \leq 2^{|x|}$: calculate $f_{i^*}(y)$. If and only if there is a y such that $f_{i^*}(y) = x$, then $x \in \bar{L}$.

L is member of $\text{co-NTIME}(2^n)$

- ▶ As $L \in \text{co-NTIME}(2^n) \Leftrightarrow \bar{L} \in \text{NTIME}(2^n)$, we will analyze the complexity of \bar{L} .
- ▶ $\bar{L} = \overline{\bigcup_{i>0} L'_i} = \bigcap_{i>0} \bar{L}'_i$
- ▶ claim: $\bigcap_{i>0} \bar{L}'_i \in \text{NTIME}(2^n)$

$$\bar{L}'_i = \{x \in \Sigma^* : x \notin L_i \vee (\exists y \in \Sigma^* (|y|^{2^i} \leq 2^{|x|}) \wedge (f_i(y) = x))\}$$

Let x be an arbitrary string.

- ▶ Check, if x is member of any L_i : if not, then $x \in \bar{L}$
- ▶ Otherwise, choose i^* such that $x \in L_{i^*}$
- ▶ $x \in \bar{L}'_j$ for any $j \neq i^*$
- ▶ for any y such that $|y|^{2^i} \leq 2^{|x|}$: calculate $f_{i^*}(y)$. If and only if there is a y such that $f_{i^*}(y) = x$, then $x \in \bar{L}$.

L is member of $\text{co-NTIME}(2^n)$

- ▶ As $L \in \text{co-NTIME}(2^n) \Leftrightarrow \bar{L} \in \text{NTIME}(2^n)$, we will analyze the complexity of \bar{L} .
- ▶ $\bar{L} = \overline{\bigcup_{i>0} L'_i} = \bigcap_{i>0} \bar{L}'_i$
- ▶ claim: $\bigcap_{i>0} \bar{L}'_i \in \text{NTIME}(2^n)$

$$\bar{L}'_i = \{x \in \Sigma^* : x \notin L_i \vee (\exists y \in \Sigma^* (|y|^{2^i} \leq 2^{|x|}) \wedge (f_i(y) = x))\}$$

Let x be an arbitrary string.

- ▶ Check, if x is member of any L_i : if not, then $x \in \bar{L}$
 - ▶ Otherwise, choose i^* such that $x \in L_{i^*}$
 - ▶ $x \in \bar{L}'_j$ for any $j \neq i^*$
 - ▶ for any y such that $|y|^{2^{i^*}} \leq 2^{|x|}$: calculate $f_{i^*}(y)$. If and only if there is a y such that $f_{i^*}(y) = x$, then $x \in \bar{L}$.

L is member of $\text{co-NTIME}(2^n)$

- ▶ As $L \in \text{co-NTIME}(2^n) \Leftrightarrow \bar{L} \in \text{NTIME}(2^n)$, we will analyze the complexity of \bar{L} .
- ▶ $\bar{L} = \overline{\bigcup_{i>0} L'_i} = \bigcap_{i>0} \bar{L}'_i$
- ▶ claim: $\bigcap_{i>0} \bar{L}'_i \in \text{NTIME}(2^n)$

$$\bar{L}'_i = \{x \in \Sigma^* : x \notin L_i \vee (\exists y \in \Sigma^* (|y|^{2i} \leq 2^{|x|}) \wedge (f_i(y) = x))\}$$

Let x be an arbitrary string.

- ▶ Check, if x is member of any L_i : **if not, then $x \in \bar{L}$**
- ▶ Otherwise, choose i^* such that $x \in L_{i^*}$
- ▶ $x \in \bar{L}'_j$ for any $j \neq i$
- ▶ for any y such that $|y|^{2i} \leq 2^{|x|}$: calculate $f_{i^*}(y)$. If and only if there is a y such that $f_{i^*}(y) = x$, then $x \in \bar{L}$.

L is member of $\text{co-NTIME}(2^n)$

- ▶ As $L \in \text{co-NTIME}(2^n) \Leftrightarrow \bar{L} \in \text{NTIME}(2^n)$, we will analyze the complexity of \bar{L} .
- ▶ $\bar{L} = \overline{\bigcup_{i>0} L'_i} = \bigcap_{i>0} \bar{L}'_i$
- ▶ claim: $\bigcap_{i>0} \bar{L}'_i \in \text{NTIME}(2^n)$

$$\bar{L}'_i = \{x \in \Sigma^* : x \notin L_i \vee (\exists y \in \Sigma^* (|y|^{2^i} \leq 2^{|x|}) \wedge (f_i(y) = x))\}$$

Let x be an arbitrary string.

- ▶ Check, if x is member of any L_i : if not, then $x \in \bar{L}$
- ▶ Otherwise, choose i^* such that $x \in L_{i^*}$
- ▶ $x \in \bar{L}'_j$ for any $j \neq i$
- ▶ for any y such that $|y|^{2^i} \leq 2^{|x|}$: calculate $f_{i^*}(y)$. If and only if there is a y such that $f_{i^*}(y) = x$, then $x \in \bar{L}$.

L is member of $\text{co-NTIME}(2^n)$

- ▶ As $L \in \text{co-NTIME}(2^n) \Leftrightarrow \bar{L} \in \text{NTIME}(2^n)$, we will analyze the complexity of \bar{L} .
- ▶ $\bar{L} = \overline{\bigcup_{i>0} L'_i} = \bigcap_{i>0} \bar{L}'_i$
- ▶ claim: $\bigcap_{i>0} \bar{L}'_i \in \text{NTIME}(2^n)$

$$\bar{L}'_i = \{x \in \Sigma^* : x \notin L_i \vee (\exists y \in \Sigma^* (|y|^{2^i} \leq 2^{|x|}) \wedge (f_i(y) = x))\}$$

Let x be an arbitrary string.

- ▶ Check, if x is member of any L_i : if not, then $x \in \bar{L}$
- ▶ Otherwise, choose i^* such that $x \in L_{i^*}$
- ▶ $x \in \bar{L}'_j$ for any $j \neq i^*$
- ▶ for any y such that $|y|^{2^i} \leq 2^{|x|}$: calculate $f_{i^*}(y)$. If and only if there is a y such that $f_{i^*}(y) = x$, then $x \in \bar{L}$.

L is member of $\text{co-NTIME}(2^n)$

- ▶ As $L \in \text{co-NTIME}(2^n) \Leftrightarrow \bar{L} \in \text{NTIME}(2^n)$, we will analyze the complexity of \bar{L} .
- ▶ $\bar{L} = \overline{\bigcup_{i>0} L'_i} = \bigcap_{i>0} \bar{L}'_i$
- ▶ claim: $\bigcap_{i>0} \bar{L}'_i \in \text{NTIME}(2^n)$

$$\bar{L}'_i = \{x \in \Sigma^* : x \notin L_i \vee (\exists_{y \in \Sigma^*} (|y|^{2^i} \leq 2^{|x|}) \wedge (f_i(y) = x))\}$$

Let x be an arbitrary string.

- ▶ Check, if x is member of any L_i : if not, then $x \in \bar{L}$
- ▶ Otherwise, choose i^* such that $x \in L_{i^*}$
- ▶ $x \in \bar{L}'_j$ for any $j \neq i^*$
- ▶ for any y such that $|y|^{2^i} \leq 2^{|x|}$: calculate $f_{i^*}(y)$. If and only if there is a y such that $f_{i^*}(y) = x$, then $x \in \bar{L}$.

L is member of $\text{co-NTIME}(2^n)$

- ▶ As $L \in \text{co-NTIME}(2^n) \Leftrightarrow \bar{L} \in \text{NTIME}(2^n)$, we will analyze the complexity of \bar{L} .
- ▶ $\bar{L} = \overline{\bigcup_{i>0} L'_i} = \bigcap_{i>0} \bar{L}'_i$
- ▶ claim: $\bigcap_{i>0} \bar{L}'_i \in \text{NTIME}(2^n)$

$$\bar{L}'_i = \{x \in \Sigma^* : x \notin L_i \vee (\exists y \in \Sigma^* (|y|^{2^i} \leq 2^{|x|}) \wedge (f_i(y) = x))\}$$

Let x be an arbitrary string.

- ▶ Check, if x is member of any L_i : if not, then $x \in \bar{L}$
- ▶ Otherwise, choose i^* such that $x \in L_{i^*}$
- ▶ $x \in \bar{L}'_j$ for any $j \neq i^*$
- ▶ for any y such that $|y|^{2^i} \leq 2^{|x|}$: calculate $f_{i^*}(y)$. If and only if there is a y such that $f_{i^*}(y) = x$, then $x \in \bar{L}$.

L is member of $\text{co-NTIME}(2^n)$

- ▶ As $L \in \text{co-NTIME}(2^n) \Leftrightarrow \bar{L} \in \text{NTIME}(2^n)$, we will analyze the complexity of \bar{L} .
- ▶ $\bar{L} = \overline{\bigcup_{i>0} L'_i} = \bigcap_{i>0} \bar{L}'_i$
- ▶ claim: $\bigcap_{i>0} \bar{L}'_i \in \text{NTIME}(2^n)$

$$\bar{L}'_i = \{x \in \Sigma^* : x \notin L_i \vee (\exists y \in \Sigma^* (|y|^{2^i} \leq 2^{|x|}) \wedge (f_i(y) = x))\}$$

Let x be an arbitrary string.

- ▶ Check, if x is member of any L_i : if not, then $x \in \bar{L}$
- ▶ Otherwise, choose i^* such that $x \in L_{i^*}$
- ▶ $x \in \bar{L}'_j$ for any $j \neq i^*$
- ▶ for any y such that $|y|^{2^i} \leq 2^{|x|}$: calculate $f_{i^*}(y)$. If and only if there is a y such that $f_{i^*}(y) = x$, then $x \in \bar{L}$.

L is member of $\text{co-NTIME}(2^n)$

- ▶ As $L \in \text{co-NTIME}(2^n) \Leftrightarrow \bar{L} \in \text{NTIME}(2^n)$, we will analyze the complexity of \bar{L} .
- ▶ $\bar{L} = \overline{\bigcup_{i>0} L'_i} = \bigcap_{i>0} \bar{L}'_i$
- ▶ claim: $\bigcap_{i>0} \bar{L}'_i \in \text{NTIME}(2^n)$

$$\bar{L}'_i = \{x \in \Sigma^* : x \notin L_i \vee (\exists y \in \Sigma^* (|y|^{2^i} \leq 2^{|x|}) \wedge (f_i(y) = x))\}$$

Let x be an arbitrary string.

- ▶ Check, if x is member of any L_i : if not, then $x \in \bar{L}$
- ▶ Otherwise, choose i^* such that $x \in L_{i^*}$
- ▶ $x \in \bar{L}'_j$ for any $j \neq i^*$
- ▶ for any y such that $|y|^{2^i} \leq 2^{|x|}$: calculate $f_{i^*}(y)$. If and only if there is a y such that $f_{i^*}(y) = x$, then $x \in \bar{L}$.

A Property for Proof Systems of L

Recall $L'_i = \{x \in L_i : \forall_{y \in \Sigma^*} |y|^{2^i} \leq 2^{|x|} \implies f_i(y) \neq x\}$
and $\overline{L}'_i = \{x \in \Sigma^* : x \notin L_i \vee (\exists_{y \in \Sigma^*} |y|^{2^i} \leq 2^{|x|} \wedge f_i(y) = x)\}$

- ▶ any proof system of L is one of the f_i
- ▶ claim: for any $f_i(\Sigma^*) = L$ it holds $L_i = L'_i$
- ▶ let f_i be a proof system for L
- ▶ assume there is a $x = 0^i 1 z \in L_i$ that is not a member of L'_i
- ▶ then, there is a y such that $|y|^{2^i} \leq 2^{|x|}$ and $f_i(y) = x$
- ▶ Hence, y is a f_i -proof for x . It follows that $x \in L$, and therefore $x \in L'_i$. This is a contradiction.

A Property for Proof Systems of L

Recall $L'_i = \{x \in L_i : \forall_{y \in \Sigma^*} |y|^{2^i} \leq 2^{|x|} \implies f_i(y) \neq x\}$
and $\overline{L}'_i = \{x \in \Sigma^* : x \notin L_i \vee (\exists_{y \in \Sigma^*} |y|^{2^i} \leq 2^{|x|} \wedge f_i(y) = x)\}$

- ▶ any proof system of L is one of the f_i
- ▶ claim: for any $f_i(\Sigma^*) = L$ it holds $L_i = L'_i$
- ▶ let f_i be a proof system for L
- ▶ assume there is a $x = 0^i 1 z \in L_i$ that is not a member of L'_i
- ▶ then, there is a y such that $|y|^{2^i} \leq 2^{|x|}$ and $f_i(y) = x$
- ▶ Hence, y is a f_i -proof for x . It follows that $x \in L$, and therefore $x \in L'_i$. This is a contradiction.

A Property for Proof Systems of L

Recall $L'_i = \{x \in L_i : \forall_{y \in \Sigma^*} |y|^{2^i} \leq 2^{|x|} \implies f_i(y) \neq x\}$
and $\overline{L}'_i = \{x \in \Sigma^* : x \notin L_i \vee (\exists_{y \in \Sigma^*} |y|^{2^i} \leq 2^{|x|} \wedge f_i(y) = x)\}$

- ▶ any proof system of L is one of the f_i
- ▶ claim: for any $f_i(\Sigma^*) = L$ it holds $L_i = L'_i$
- ▶ let f_i be a proof system for L
- ▶ assume there is a $x = 0^i 1 z \in L_i$ that is not a member of L'_i
- ▶ then, there is a y such that $|y|^{2^i} \leq 2^{|x|}$ and $f_i(y) = x$
- ▶ Hence, y is a f_i -proof for x . It follows that $x \in L$, and therefore $x \in L'_i$. This is a contradiction.

A Property for Proof Systems of L

Recall $L'_i = \{x \in L_i : \forall_{y \in \Sigma^*} |y|^{2^i} \leq 2^{|x|} \implies f_i(y) \neq x\}$
and $\overline{L}'_i = \{x \in \Sigma^* : x \notin L_i \vee (\exists_{y \in \Sigma^*} |y|^{2^i} \leq 2^{|x|} \wedge f_i(y) = x)\}$

- ▶ any proof system of L is one of the f_i
- ▶ claim: for any $f_i(\Sigma^*) = L$ it holds $L_i = L'_i$
- ▶ let f_i be a proof system for L
- ▶ assume there is a $x = 0^i 1 z \in L_i$ that is not a member of L'_i
- ▶ then, there is a y such that $|y|^{2^i} \leq 2^{|x|}$ and $f_i(y) = x$
- ▶ Hence, y is a f_i -proof for x . It follows that $x \in L$, and therefore $x \in L'_i$. This is a contradiction.

A Property for Proof Systems of L

Recall $L'_i = \{x \in L_i : \forall_{y \in \Sigma^*} |y|^{2^i} \leq 2^{|x|} \implies f_i(y) \neq x\}$
and $\overline{L}'_i = \{x \in \Sigma^* : x \notin L_i \vee (\exists_{y \in \Sigma^*} |y|^{2^i} \leq 2^{|x|} \wedge f_i(y) = x)\}$

- ▶ any proof system of L is one of the f_i
- ▶ claim: for any $f_i(\Sigma^*) = L$ it holds $L_i = L'_i$
- ▶ let f_i be a proof system for L
- ▶ assume there is a $x = 0^i 1z \in L_i$ that is not a member of L'_i
- ▶ then, there is a y such that $|y|^{2^i} \leq 2^{|x|}$ and $f_i(y) = x$
- ▶ Hence, y is a f_i -proof for x . It follows that $x \in L$, and therefore $x \in L'_i$. This is a contradiction.

A Property for Proof Systems of L

Recall $L'_i = \{x \in L_i : \forall_{y \in \Sigma^*} |y|^{2^i} \leq 2^{|x|} \implies f_i(y) \neq x\}$
and $\overline{L}'_i = \{x \in \Sigma^* : x \notin L_i \vee (\exists_{y \in \Sigma^*} |y|^{2^i} \leq 2^{|x|} \wedge f_i(y) = x)\}$

- ▶ any proof system of L is one of the f_i
- ▶ claim: for any $f_i(\Sigma^*) = L$ it holds $L_i = L'_i$
- ▶ let f_i be a proof system for L
- ▶ assume there is a $x = 0^i 1 z \in L_i$ that is not a member of L'_i
- ▶ then, there is a y such that $|y|^{2^i} \leq 2^{|x|}$ and $f_i(y) = x$
- ▶ Hence, y is a f_i -proof for x . It follows that $x \in L$, and therefore $x \in L'_i$. This is a contradiction.

A Property for Proof Systems of L

Recall $L'_i = \{x \in L_i : \forall_{y \in \Sigma^*} |y|^{2^i} \leq 2^{|x|} \implies f_i(y) \neq x\}$
and $\overline{L'_i} = \{x \in \Sigma^* : x \notin L_i \vee (\exists_{y \in \Sigma^*} |y|^{2^i} \leq 2^{|x|} \wedge f_i(y) = x)\}$

- ▶ any proof system of L is one of the f_i
- ▶ claim: for any $f_i(\Sigma^*) = L$ it holds $L_i = L'_i$
- ▶ let f_i be a proof system for L
- ▶ assume there is a $x = 0^i 1 z \in L_i$ that is not a member of L'_i
- ▶ then, there is a y such that $|y|^{2^i} \leq 2^{|x|}$ and $f_i(y) = x$
- ▶ Hence, y is a f_i -proof for x . It follows that $x \in L$, and therefore $x \in L'_i$. This is a contradiction.

A Property for Proof Systems of L

Recall $L'_i = \{x \in L_i : \forall_{y \in \Sigma^*} |y|^{2^i} \leq 2^{|x|} \implies f_i(y) \neq x\}$
and $\overline{L}'_i = \{x \in \Sigma^* : x \notin L_i \vee (\exists_{y \in \Sigma^*} |y|^{2^i} \leq 2^{|x|} \wedge f_i(y) = x)\}$

- ▶ any proof system of L is one of the f_i
- ▶ claim: for any $f_i(\Sigma^*) = L$ it holds $L_i = L'_i$
- ▶ let f_i be a proof system for L
- ▶ assume there is a $x = 0^i 1 z \in L_i$ that **is not a member of L'_i**
- ▶ then, there is a y such that $|y|^{2^i} \leq 2^{|x|}$ and $f_i(y) = x$
- ▶ Hence, y is a f_i -proof for x . It follows that $x \in L$, and therefore **$x \in L'_i$** . This is a contradiction.

These results contradict the existence of an optimal proof system for L

Recall $L'_i = \{x \in L_i : \forall_{y \in \Sigma^*} |y|^{2i} \leq 2^{|x|} \implies f_i(y) \neq x\}$.

- ▶ Let f_i be an optimal proof system for L

- ▶ Let g be defined by

$$g(bx) = \begin{cases} f_i(x) & (b = 0) \\ x & (b = 1 \text{ and } x = 0^i 10^* \in L_i = L'_i) \end{cases}$$

- ▶ g is a proof system for L
- ▶ as f_i is optimal, there is a f^* , such that $f_i(f^*(x)) = g(x)$. f^* is polynomial bounded: $|f^*(x)| \leq p(|x|)$
- ▶ for any x of L'_i , such that $p(|1x|)^{2i} \leq 2^{|x|}$.
- ▶ $1x$ is a g -proof for x : $g(1x) = x$
- ▶ For $y = f^*(1x)$ it holds $|y| = |f^*(1x)| \leq p(|1x|) \leq p(|1x|)^{2i} \leq 2^{|x|}$.
- ▶ By Definition of L'_i : $f_i(y) \neq x$, hence $y = f^*(1x)$ is no f_i -proof for x
- ▶ This is a contradiction \square

These results contradict the existence of an optimal proof system for L

Recall $L'_i = \{x \in L_i : \forall_{y \in \Sigma^*} |y|^{2i} \leq 2^{|x|} \implies f_i(y) \neq x\}$.

► Let f_i be an optimal proof system for L

► Let g be defined by

$$g(bx) = \begin{cases} f_i(x) & (b = 0) \\ x & (b = 1 \text{ and } x = 0^i 10^* \in L_i = L'_i) \end{cases}$$

► g is a proof system for L

► as f_i is optimal, there is a f^* , such that $f_i(f^*(x)) = g(x)$. f^* is polynomial bounded: $|f^*(x)| \leq p(|x|)$

► for any x of L'_i , such that $p(|1x|)^{2i} \leq 2^{|x|}$.

► $1x$ is a g -proof for x : $g(1x) = x$

► For $y = f^*(1x)$ it holds

$$|y| = |f^*(1x)| \leq p(|1x|) \leq p(|1x|)^{2i} \leq 2^{|x|}.$$

► By Definition of L'_i : $f_i(y) \neq x$, hence $y = f^*(1x)$ is no f_i -proof for x

► This is a contradiction \square

These results contradict the existence of an optimal proof system for L

Recall $L'_i = \{x \in L_i : \forall_{y \in \Sigma^*} |y|^{2i} \leq 2^{|x|} \implies f_i(y) \neq x\}$.

► Let f_i be an optimal proof system for L

► Let g be defined by

$$g(bx) = \begin{cases} f_i(x) & (b = 0) \\ x & (b = 1 \text{ and } x = 0^i 10^* \in L_i = L'_i) \end{cases}$$

► g is a proof system for L

► as f_i is optimal, there is a f^* , such that $f_i(f^*(x)) = g(x)$. f^* is polynomial bounded: $|f^*(x)| \leq p(|x|)$

► for any x of L'_i , such that $p(|1x|)^{2i} \leq 2^{|x|}$.

► $1x$ is a g -proof for x : $g(1x) = x$

► For $y = f^*(1x)$ it holds

$$|y| = |f^*(1x)| \leq p(|1x|) \leq p(|1x|)^{2i} \leq 2^{|x|}.$$

► By Definition of L'_i : $f_i(y) \neq x$, hence $y = f^*(1x)$ is no f_i -proof for x

► This is a contradiction \square

These results contradict the existence of an optimal proof system for L

Recall $L'_i = \{x \in L_i : \forall_{y \in \Sigma^*} |y|^{2i} \leq 2^{|x|} \implies f_i(y) \neq x\}$.

- ▶ Let f_i be an optimal proof system for L

- ▶ Let g be defined by

$$g(bx) = \begin{cases} f_i(x) & (b = 0) \\ x & (b = 1 \text{ and } x = 0^i 10^* \in L_i = L'_i) \end{cases}$$

- ▶ g is a proof system for L

- ▶ as f_i is optimal, there is a f^* , such that $f_i(f^*(x)) = g(x)$. f^* is polynomial bounded: $|f^*(x)| \leq p(|x|)$

- ▶ for any x of L'_i , such that $p(|1x|)^{2i} \leq 2^{|x|}$.

- ▶ $1x$ is a g -proof for x : $g(1x) = x$

- ▶ For $y = f^*(1x)$ it holds

$$|y| = |f^*(1x)| \leq p(|1x|) \leq p(|1x|)^{2i} \leq 2^{|x|}.$$

- ▶ By Definition of L'_i : $f_i(y) \neq x$, hence $y = f^*(1x)$ is no f_i -proof for x

- ▶ This is a contradiction \square

These results contradict the existence of an optimal proof system for L

Recall $L'_i = \{x \in L_i : \forall_{y \in \Sigma^*} |y|^{2i} \leq 2^{|x|} \implies f_i(y) \neq x\}$.

- ▶ Let f_i be an optimal proof system for L

- ▶ Let g be defined by

$$g(bx) = \begin{cases} f_i(x) & (b = 0) \\ x & (b = 1 \text{ and } x = 0^i 10^* \in L_i = L'_i) \end{cases}$$

- ▶ g is a proof system for L

- ▶ as f_i is optimal, there is a f^* , such that $f_i(f^*(x)) = g(x)$. f^* is polynomial bounded: $|f^*(x)| \leq p(|x|)$

- ▶ for any x of L'_i , such that $p(|1x|)^{2i} \leq 2^{|x|}$.

- ▶ $1x$ is a g -proof for x : $g(1x) = x$

- ▶ For $y = f^*(1x)$ it holds

$$|y| = |f^*(1x)| \leq p(|1x|) \leq p(|1x|)^{2i} \leq 2^{|x|}.$$

- ▶ By Definition of L'_i : $f_i(y) \neq x$, hence $y = f^*(1x)$ is no f_i -proof for x

- ▶ This is a contradiction \square

These results contradict the existence of an optimal proof system for L

Recall $L'_i = \{x \in L_i : \forall_{y \in \Sigma^*} |y|^{2i} \leq 2^{|x|} \implies f_i(y) \neq x\}$.

- ▶ Let f_i be an optimal proof system for L

- ▶ Let g be defined by

$$g(bx) = \begin{cases} f_i(x) & (b = 0) \\ x & (b = 1 \text{ and } x = 0^i 10^* \in L_i = L'_i) \end{cases}$$

- ▶ g is a proof system for L

- ▶ as f_i is optimal, there is a f^* , such that $f_i(f^*(x)) = g(x)$. f^* is polynomial bounded: $|f^*(x)| \leq p(|x|)$

- ▶ for any x of L'_i , such that $p(|1x|)^{2i} \leq 2^{|x|}$.

- ▶ $1x$ is a g -proof for x : $g(1x) = x$

- ▶ For $y = f^*(1x)$ it holds

$$|y| = |f^*(1x)| \leq p(|1x|) \leq p(|1x|)^{2i} \leq 2^{|x|}.$$

- ▶ By Definition of L'_i : $f_i(y) \neq x$, hence $y = f^*(1x)$ is no f_i -proof for x

- ▶ This is a contradiction \square

These results contradict the existence of an optimal proof system for L

Recall $L'_i = \{x \in L_i : \forall_{y \in \Sigma^*} |y|^{2i} \leq 2^{|x|} \implies f_i(y) \neq x\}$.

- ▶ Let f_i be an optimal proof system for L

- ▶ Let g be defined by

$$g(bx) = \begin{cases} f_i(x) & (b = 0) \\ x & (b = 1 \text{ and } x = 0^i 10^* \in L_i = L'_i) \end{cases}$$

- ▶ g is a proof system for L

- ▶ as f_i is optimal, there is a f^* , such that $f_i(f^*(x)) = g(x)$. f^* is polynomial bounded: $|f^*(x)| \leq p(|x|)$

- ▶ for any x of L'_i , such that $p(|1x|)^{2i} \leq 2^{|x|}$.

- ▶ $1x$ is a g -proof for x : $g(1x) = x$

- ▶ For $y = f^*(1x)$ it holds

$$|y| = |f^*(1x)| \leq p(|1x|) \leq p(|1x|)^{2i} \leq 2^{|x|}.$$

- ▶ By Definition of L'_i : $f_i(y) \neq x$, hence $y = f^*(1x)$ is no f_i -proof for x

- ▶ This is a contradiction \square

These results contradict the existence of an optimal proof system for L

Recall $L'_i = \{x \in L_i : \forall_{y \in \Sigma^*} |y|^{2i} \leq 2^{|x|} \implies f_i(y) \neq x\}$.

► Let f_i be an optimal proof system for L

► Let g be defined by

$$g(bx) = \begin{cases} f_i(x) & (b = 0) \\ x & (b = 1 \text{ and } x = 0^i 10^* \in L_i = L'_i) \end{cases}$$

► g is a proof system for L

► as f_i is optimal, there is a f^* , such that $f_i(f^*(x)) = g(x)$. f^* is polynomial bounded: $|f^*(x)| \leq p(|x|)$

► for any x of L'_i , such that $p(|1x|)^{2i} \leq 2^{|x|}$.

► $1x$ is a g -proof for x : $g(1x) = x$

► For $y = f^*(1x)$ it holds

$$|y| = |f^*(1x)| \leq p(|1x|) \leq p(|1x|)^{2i} \leq 2^{|x|}.$$

► By Definition of L'_i : $f_i(y) \neq x$, hence $y = f^*(1x)$ is no f_i -proof for x

► This is a contradiction \square

These results contradict the existence of an optimal proof system for L

Recall $L'_i = \{x \in L_i : \forall_{y \in \Sigma^*} |y|^{2i} \leq 2^{|x|} \implies f_i(y) \neq x\}$.

- ▶ Let f_i be an optimal proof system for L

- ▶ Let g be defined by

$$g(bx) = \begin{cases} f_i(x) & (b = 0) \\ x & (b = 1 \text{ and } x = 0^i 10^* \in L_i = L'_i) \end{cases}$$

- ▶ g is a proof system for L

- ▶ as f_i is optimal, there is a f^* , such that $f_i(f^*(x)) = g(x)$. f^* is polynomial bounded: $|f^*(x)| \leq p(|x|)$

- ▶ for any x of L'_i , such that $p(|1x|)^{2i} \leq 2^{|x|}$.

- ▶ $1x$ is a g -proof for x : $g(1x) = x$

- ▶ For $y = f^*(1x)$ it holds

$$|y| = |f^*(1x)| \leq p(|1x|) \leq p(|1x|)^{2i} \leq 2^{|x|}.$$

- ▶ By Definition of L'_i : $f_i(y) \neq x$, hence $y = f^*(1x)$ is no f_i -proof for x

- ▶ This is a contradiction \square

These results contradict the existence of an optimal proof system for L

Recall $L'_i = \{x \in L_i : \forall_{y \in \Sigma^*} |y|^{2i} \leq 2^{|x|} \implies f_i(y) \neq x\}$.

- ▶ Let f_i be an optimal proof system for L

- ▶ Let g be defined by

$$g(bx) = \begin{cases} f_i(x) & (b = 0) \\ x & (b = 1 \text{ and } x = 0^i 10^* \in L_i = L'_i) \end{cases}$$

- ▶ g is a proof system for L

- ▶ as f_i is optimal, there is a f^* , such that $f_i(f^*(x)) = g(x)$. f^* is polynomial bounded: $|f^*(x)| \leq p(|x|)$

- ▶ for any x of L'_i , such that $p(|1x|)^{2i} \leq 2^{|x|}$.

- ▶ $1x$ is a g -proof for x : $g(1x) = x$

- ▶ For $y = f^*(1x)$ it holds

$$|y| = |f^*(1x)| \leq p(|1x|) \leq p(|1x|)^{2i} \leq 2^{|x|}.$$

- ▶ By Definition of L'_i : $f_i(y) \neq x$, hence $y = f^*(1x)$ is no f_i -proof for x

- ▶ This is a contradiction \square

Consequences

What do we know about L ?

Theorem

*If $L \subseteq 0^*10^*$ has no optimal proof system, then there is a polynomial time equivalent $T \in TALLY$, that does not possess an optimal proof system.*

Corollary

Let $u : \mathbb{N} \rightarrow \mathbb{N}$ be a strictly increasing polynomial time function. Then there is a language $L \in co-NTIME(2^n)$, that has no optimal proof system. Additionally, every length of a string in L is in the range of u .

Theorem

For every language L that possesses no optimal proof system there is a polynomial time equivalent, many-one-mitotic language, that does not possess an optimal proof system.

Consequences

What do we know about L ?

Theorem

*If $L \subseteq 0^*10^*$ has no optimal proof system, then there is a polynomial time equivalent $T \in \text{TALLY}$, that does not possess an optimal proof system.*

Corollary

Let $u : \mathbb{N} \rightarrow \mathbb{N}$ be a strictly increasing polynomial time function. Then there is a language $L \in \text{co-NTIME}(2^n)$, that has no optimal proof system. Additionally, every length of a string in L is in the range of u .

Theorem

For every language L that possesses no optimal proof system there is a polynomial time equivalent, many-one-mitotic language, that does not possess an optimal proof system.

Consequences

What do we know about L ?

Theorem

*If $L \subseteq 0^*10^*$ has no optimal proof system, then there is a polynomial time equivalent $T \in TALLY$, that does not possess an optimal proof system.*

Corollary

Let $u : \mathbb{N} \rightarrow \mathbb{N}$ be a strictly increasing polynomial time function. Then there is a language $L \in co-NTIME(2^n)$, that has no optimal proof system. Additionally, every length of a string in L is in the range of u .

Theorem

For every language L that possesses no optimal proof system there is a polynomial time equivalent, many-one-mitotic language, that does not possess an optimal proof system.

Consequences

What do we know about L ?

Theorem

*If $L \subseteq 0^*10^*$ has no optimal proof system, then there is a polynomial time equivalent $T \in TALLY$, that does not possess an optimal proof system.*

Corollary

Let $u : \mathbb{N} \rightarrow \mathbb{N}$ be a strictly increasing polynomial time function. Then there is a language $L \in co-NTIME(2^n)$, that has no optimal proof system. Additionally, every length of a string in L is in the range of u .

Theorem

For every language L that possesses no optimal proof system there is a polynomial time equivalent, many-one-mitotic language, that does not possess an optimal proof system.

Consequences

What do we know about L ?

Theorem

*If $L \subseteq 0^*10^*$ has no optimal proof system, then there is a polynomial time equivalent $T \in TALLY$, that does not possess an optimal proof system.*

Corollary

Let $u : \mathbb{N} \rightarrow \mathbb{N}$ be a strictly increasing polynomial time function. Then there is a language $L \in co-NTIME(2^n)$, that has no optimal proof system. Additionally, every length of a string in L is in the range of u .

Theorem

For every language L that possesses no optimal proof system there is a polynomial time equivalent, many-one-mitotic language, that does not possess an optimal proof system.

Literature I

- [AB09] Sanjeev Arora and Boaz Barak, *Computational complexity: A modern approach*, 1st ed., Cambridge University Press, New York, NY, USA, 2009.
- [Coo71] Stephen A. Cook, *The complexity of theorem-proving procedures*, Proceedings of the third annual ACM symposium on Theory of computing (New York, NY, USA), STOC '71, ACM, 1971, pp. 151–158.
- [CR79] Stephen A. Cook and Robert A. Reckhow, *The relative efficiency of propositional proof systems*, Journal of Symbolic Logic **44** (1979), 36–50.
- [For09] Lance Fortnow, *The status of the P versus NP problem*, Commun. ACM **52** (2009), no. 9, 78–86.

Literature II

- [KM00] Johannes Köbler and Jochen Messner, *Is the standard proof system for SAT p -optimal?*, Proceedings of the 20th Conference on Foundations of Software Technology and Theoretical Computer Science (London, UK, UK), FST TCS 2000, Springer-Verlag, 2000, pp. 361–372.
- [KMT03] Johannes Köbler, Jochen Messner, and Jacobo Torán, *Optimal proof systems imply complete sets for promise classes*, Inf. Comput. **184** (2003), no. 1, 71–92.
- [KP89] Jan Krajíček and Pavel Pudlák, *Propositional proof systems, the consistency of first order theories and the complexity of computations*, J. Symb. Log. **54** (1989), no. 3, 1063–1079.

Literature III

- [Mes99] Jochen Messner, *On optimal algorithms and optimal proof systems*, Proceedings of the 16th annual conference on Theoretical aspects of computer science (Berlin, Heidelberg), STACS'99, Springer-Verlag, 1999, pp. 541–550.
- [Pap94] Christos H. Papadimitriou, *Computational complexity*, Addison-Wesley, 1994.

Images Sources I

- ▶ Kurt Gödel: Wikipedia, de.wikipedia.org