

The Complexity of Finding Tarski Fixed Points

Master Thesis

August 25, 2024

Nils Jensen

ADVISED BY

PROF. DR. BERND GÄRTNER

SEBASTIAN HASLEBACHER

Abstract

Insert the abstract here.

Acknowledgements

Insert the acknowledgements here.

Contents

Contents	v
List of Figures	vii
List of Tables	vii
1 Introduction	1
1.1 Total Search Problems	1
1.2 The TFNP landscape	2
1.3 The TARSKI problem	2
1.4 Current algorithms for solving TARSKI	3
1.5 Location of TARSKI in TFNP	4
1.6 Thesis Outline	4
2 Preliminaries	5
2.1 Total search problems	5
2.1.1 Search problems	5
2.1.2 Reductions	7
2.1.3 Promise Problems	7
2.1.4 Representation of sets	8
2.1.5 Representation of functions	9
2.1.6 Complexity of boolean circuits	11
2.2 Subclasses of TFNP	12
2.2.1 Polynomial Local Search (PLS)	12
2.2.2 Polynomial Parity Argument on Directed Graphs (PPAD)	13
2.2.3 End of Potential Line (EOPL)	14
2.3 The TARSKI Problem	15
2.3.1 Definition of the TARSKI Problem	15
2.3.2 Two algorithms for solving TARSKI	17
2.3.3 Lower bounds for TARSKI	20
2.3.4 Location of TARSKI in TFNP	20
2.3.5 Variants of TARSKI	21
3 Reducing TARSKI to PPAD	23
3.1 Presentation of the known reduction of TARSKI to PPAD	23
3.2 Introducing TARSKI*	24
3.3 Sperner's Lemma	26
3.3.1 Sperner's Lemma for Simplices	27
3.3.2 Sperner's Lemma for an integer lattice	28
3.3.3 Reducing SPERNER to ENDOFLINE	29
3.4 Reducing TARSKI* to SPERNER	30

4	Reducing TARSKI to EOPL	33
4.1	Warmup: No cycles in two-dimensional TARSKI*-instances	33
4.2	Choosing a simplicial decomposition of the lattice — Freudenthal’s Simplicial Decomposition	35
4.3	Sequences of simplices	39
4.4	A side note on super-unique TARSKI instances	40
4.5	Orientation of a the simplicial decomposition	40
4.5.1	Orientation of a simplex	41
4.5.2	Orientation of a simplicial complex	43
4.5.3	Orienting colored simplices	46
4.6	Properties of colored of oriented simplicial sequences	47
4.6.1	General properties of the coloring	47
4.6.2	Properties of sequences of simplices	47
4.7	No cycles in the ENDOFLINE instance	50
4.8	Discussing the reduction of TARSKI* to ENDOFPOTENTIALLINE	51
	 APPENDIX	 53
	Bibliography	54
	Alphabetical Index	57

List of Figures

2.1	Example of a Boolean Circuit	10
2.2	Computing a function with circuits	11
2.3	Example of a LOCALOPT Problem	12
2.4	Example of an END-OF-LINE Problem	14
2.5	Example of an EOPL Problem	15
2.6	Example of a TARSKI instance	16
3.1	Setup for SPERNER'S LEMMA	27
3.2	Example of SPERNER'S LEMMA	28
3.3	Example of a simplicial decomposition of a lattice	29
3.4	Reduction of SPERNER to ENDOFLINE	30
3.5	Coloring of a TARSKI*instance	31
4.1	Sketch of the setting for the two dimensional proof	34
4.2	Orientation of a simplex	43
4.3	Orientation of a simplicial complex	44

List of Tables

List of Algorithms

1	Iterative Algorithm for TARSKI	17
2	Recursive Algorithm for TARSKI	20

1.1 Total Search Problems

The study of computational complexity is central to computer science. Its primary goal is to establish lower bounds on the complexity of various problems. Specifically, complexity theory attempts to prove that certain problems cannot be solved faster than a given time as a function of the size of the input. This endeavor has proven particularly challenging for many problems, with a significant gap between the best-known upper bounds, determined by existing algorithms, and the best-known lower bound.

A fundamental tool in complexity theory is the concept of reduction, which makes it possible to compare the difficulty of two problems. We say that a problem P_1 is reducible to another problem P_2 if P_1 can be solved efficiently by solving P_2 . This concept underlies the classification of problems into complexity classes: groups of problems that reduce onto the same fundamental problem.

Traditionally, complexity theory has focused on decision problems, which involve determining whether a given object has a given property. Examples include determining whether a graph contains a k -clique or whether a number is prime. These problems typically require a decision about whether an object belongs to a set of objects — a language — defined by a particular property.

However, real-world problems often extend beyond simple decision-making into the realm of search problems. In practical scenarios, the existence of a solution is typically assumed, and the task is not just to verify its existence but to compute the solution itself. For example, instead of just detecting the existence of a k -clique in a graph, it is likely one would wish to explicitly identify this clique or verify its absence. Similarly, in addition to recognizing a number as prime, one might want to determine its prime factors. Instead of simply deciding whether a function has a global minimum, the objective would be to compute it efficiently.

Within this broader category of search problems lies a special subclass known as *total search problems*. These are characterized by the guaranteed existence of a solution, often proven by

1.1 Total Search Problems . . .	1
1.2 The TFNP landscape . . .	2
1.3 The TARSKI problem . . .	2
1.4 Current algorithms for solving TARSKI	3
1.5 Location of TARSKI in TFNP	4
1.6 Thesis Outline	4

Here, *efficiently* generally means in polynomial time. We will define this and related concepts more strictly later.

mathematical theorems. A notable example within this subclass is the problem of identifying a sink in a directed acyclic graph. This is a *total* problem because every such graph has a sink.

1.2 The TFNP landscape

The class of **TFNP** is the pendant of **NP**, in the sense that it is the class of all total search problems, where a solution can be checked for validity in polynomial time. Studying this complexity class has been an active research subject in recent years, giving rise to many exciting results.

Because it is unexpected that we can find **TFNP**-complete problems, the class has been studied using other tools. The primary method which has been established is the use of syntactic subclasses. The idea is to build subclasses of **TFNP**, which are created by using very classical and almost obvious existence results. Three of these subclasses are particularly relevant to this thesis.

The first is the class **PPAD**, which is the class of total search problems where the existence of a solution is guaranteed by *Brouwer's fixed point theorem*. The problems in **PPAD** can be solved by walking along a directed graph, starting at an unbalanced vertex and ending at an unbalanced vertex [1].

The second class of interest is **PLS**, which is the class of total search problems that can be expressed as starting at a vertex of a directed acyclic graph and finding a sink of this graph [2].

Finally, the class **EOPL** is the class of total search problems, which can be expressed as starting at a source of a directed acyclic graph and finding a vertex which is a sink [3].

[1]: Papadimitriou (1994), *On the complexity of the parity argument and other inefficient proofs of existence*

[2]: Johnson et al. (1988), *How easy is local search?*

[3]: Fearnley et al. (2018), *End of Potential Line*

1.3 The TARSKI problem

The main problem we study in this thesis is the **TARSKI** problem. The namesake of the **TARSKI** problem is *Tarski's fixed point theorem*, which states that every monotone function on a complete lattice has a fixed point [4]. The **TARSKI** problem is the problem of finding such a fixed point for a given function f on a complete lattice L , or to find a violation of monotonicity of this function [5]. According to *Tarski's theorem*, this problem is guaranteed to have a solution, and hence, it is a total search problem.

[4]: Tarski (1955), *A lattice-theoretical fixpoint theorem and its applications*.

[5]: Etessami et al. (2020), *Tarski's Theorem, Supermodular Games, and the Complexity of Equilibria*

The TARSKI-problem has numerous applications in various fields. For example, it can be shown that supermodular games, which model certain economic situation, have a equilibrium by *Tarski's Theorem* [6, 7]. These equilibria can be found by solving TARSKI-instances [5]. The existence of equilibria in some stochastic games can be found using *Tarski's Theorem*, and finding this equilibria can be reduced to solving a TARSKI instance [8].

Another application of this problem can be found when studying the celebrated ARRIVAL problem. It can be shown that ARRIVAL reduces to TARSKI [9] hence studying the TARSKI problem can help understand the complexity of ARRIVAL.

[6]: Topkis (1979), *Equilibrium Points in Nonzero-Sum n -Person Submodular Games*

[7]: Milgrom and Roberts (1990), *Rationalizability, Learning, and Equilibrium in Games with Strategic Complementarities*

[8]: Condon (1992), *The complexity of stochastic games*

[9]: Gärtner et al. (2021), *A subexponential algorithm for ARRIVAL*

1.4 Current algorithms for solving TARSKI

We want to give an overview of the different known strategies for solving TARSKI-instances. This has a theoretical interest, as the state of these algorithms often describe graphs, which can be seen as instances of **TFNP**-complete problems, and hence can help construct reductions.

The most fundamental approach to solving the Tarski problem is a simple iterative algorithm that leverages the monotonicity of the function to converge to a fixed point iteratively. Starting from the smallest point within the lattice, the Algorithm applies the function repeatedly until a fixed point is reached [5]. This method is straightforward but can be computationally expensive, as it may require a large number of iterations to converge, particularly for functions defined over large lattices in the worst case for a lattice $L = [N]^d$, this Algorithm requires time $\mathcal{O}(N \cdot d)$.

[5]: Etessami et al. (2020), *Tarski's Theorem, Supermodular Games, and the Complexity of Equilibria*

A more sophisticated approach involves a binary search technique, where the lattice is systematically divided, and the function's monotonicity is used to eliminate regions that cannot contain a fixed point. This is done by recursively solving lower-dimensional subproblems until the fixed point is found [10]. This method can significantly reduce the search space and converges faster than the iterative Algorithm, with a runtime of $\mathcal{O}(\log^d(N))$.

[10]: Dang et al. (2020), *Computations and Complexities of Tarski's Fixed Points and Supermodular Games*

The latest developments in solving the TARSKI problem involve advanced decomposition methods that reduce the search space. These methods decompose the problem into smaller instances that can be more easily managed and solved. Using these techniques a runtime of $\mathcal{O}(\log^{\lceil \frac{d-1}{2} \rceil} N)$ can be achieved [11].

[11]: Chen and Li (2022), *Improved Upper Bounds for Finding Tarski Fixed Points*

1.5 Location of TARSKI in TFNP

It is known that the TARSKI problem lies in **PPAD** and in **PLS**. A recent breakthrough has shown that the class $\mathbf{PPAD} \cap \mathbf{PLS} = \mathbf{EOPL}$ [12]. This result immediately implies that the TARSKI problem is in **EOPL**, which in turn means that there must be a reduction from TARSKI to **EOPL**-complete problems, in particular to the **ENDOFPOTENTIALLINE** problem. The main goal of this thesis is to understand why TARSKI lies in **EOPL** and to construct a reduction from TARSKI to the **ENDOFPOTENTIALLINE** problem, which is **EOPL**-complete.

[12]: Goos et al. (2022), *Further Collapses in TFNP*

1.6 Thesis Outline

TODO: Write this section.

Preliminaries 2

This Chapter aims to establish the complexity framework used throughout this thesis to study the TARSKI problem. It formally introduces the concept of total search problems, the complexity class **TFNP**, and its subclasses **PLS**, **PPAD**, and **EOPL**. In addition, in this Chapter, we will describe how we represent sets and functions in this framework and how their complexity is measured. Finally, we give a formal introduction to the TARSKI problem and a presentation of the known algorithms for solving it and its location in the **TFNP** landscape.

2.1 Total search problems

The study of complexity classes has traditionally focused on *decision problems*, which involve determining whether an object belongs to a set, also called a *language*. Notable examples include determining whether a Boolean formula is satisfiable or whether a k -clique exists in a given graph [13]. However, real-world questions often require explicit answers rather than existence results. For example, while deciding whether a function has a global minimum is a decision problem, the practical interest lies in identifying that minimum, which goes beyond mere existence. Here, so-called *search problems* come into play.

2.1.1 Search problems

Definition 2.1 — Search Problem.

A *search problem* is given by a relation $R \subset \{0, 1\}^* \times \{0, 1\}^*$. For a given *instance* $I \in \{0, 1\}^*$ the computational problem is, to find a *solution* $s \in \{0, 1\}^*$ that satisfies: $(I, s) \in R$, or output “No” if no such s exists.

We can view these search problems as decision problems by looking at the corresponding decision problem given by the language:

$$\mathcal{L}_R = \{I \in \{0, 1\}^* \mid \exists s \in \{0, 1\}^* : (I, s) \in R\}$$

The above shows that every search problem can be seen as a decision problem of a broader language. This perspective allows

2.1	Total search problems	5
2.1.1	Search problems	5
2.1.2	Reductions	7
2.1.3	Promise Problems	7
2.1.4	Representing sets	8
2.1.5	Representing functions	9
2.1.6	Complexity of circuits	11
2.2	Subclasses of TFNP	12
2.2.1	PLS	12
2.2.2	PPAD	13
2.2.3	EOPL	14
2.3	TARSKI Problem	15
2.3.1	TARSKI Definition	15
2.3.2	TARSKI Algorithms	17
2.3.3	Lower bounds for TARSKI	20
2.3.4	TARSKI in TFNP	20
2.3.5	TARSKI Variants	21

[13]: Arora and Barak (2009), *Computational complexity : a modern approach*

The “No” case can be encoded as some special binary string.

Here, we have rephrased the valid language as the pair of a problem instance and a valid solution.

us to ask classical complexity questions about search problems: Are these problems in **P** or **NP**? Are they **NP**-hard? It is evident that search problems are at least as complex as their decision counterparts since solving a search problem inherently solves the associated decision question.

Similarly to decision problems, we want to study which problems can be solved efficiently and which cannot. This question leads us to the definition of the complexity class **FNP**, which is pendant for search problems, to **NP** for decision problems. We introduce **FNP** formally as in [14].

Definition 2.2 — Function NP (FNP).

We say that a relation $R \subset \{0,1\}^* \times \{0,1\}^*$ is in **FNP** if it is *polynomially balanced*, i.e. there exists a polynomial p such that for every $(I, s) \in R$ implies $|y| \leq p(|x|)$. The class **FNP** is the set of all relations R that are polynomially balanced.

[14]: Megiddo and Papadimitriou (1991), *On total functions, existence theorems and computational complexity*

This means that checking a solution to a search problem can be done in polynomial time. What cannot be done in polynomial time is checking whether no solution exists. This leads to the question of what happens if we remove this decision problem from the search problem. This question is what leads to the definition of a total search problem [14]:

Definition 2.3 — Total search problems.

A *total search problem* is a search problem given by a relation $R \subset \{0,1\}^* \times \{0,1\}^*$, such that for every given instance $I \in \{0,1\}^*$ there is a solution $s \in \{0,1\}^*$ that satisfies: $(I, s) \in R$.

This means that a solution always exists for any input, i.e. “No” is never a valid answer.

The complexity class **TFNP** is simply the class of all *total* problems in **FNP**.

Definition 2.4 — Total Function NP (TFNP).

The class **TFNP** is the set of all problems given by total relations polynomially bounded relations R .

Similarly to **P**, **FP** is the class of all search problems that can be solved in polynomial time. It is not known whether **FP** is equal to **TFNP**, but it is widely believed — similarly to **P** and **NP** — that they are different. Examples of **TFNP** problems are:

- **FACTORING**, the problem of finding the prime factors of a number. Every number admits a factorization into prime numbers, which can be checked in polynomial time;
- **NASH**, the problem of finding a Nash equilibrium in a bimatrix game [15];

[15]: Daskalakis et al. (2009), *The Complexity of Computing a Nash Equilibrium*

- MINIMIZE, the problem of finding the global minimum of a convex function [16].

[16]: Daskalakis and Papadimitriou (2011), *Continuous Local Search*

2.1.2 Reductions

Similarly to decision problems, we can also define reductions inside **TFNP**.

Definition 2.5 — Many-to-one Reduction.

For two problem $R, S \in \mathbf{TFNP}$, we say that R *reduces* (many to one) to S if there exist polynomial time computable functions $f : \{0, 1\}^* \rightarrow \{0, 1\}^*$ and $g : \{0, 1\}^* \times \{0, 1\}^* \rightarrow \{0, 1\}^*$ such that for $I, s \in \{0, 1\}^*$:

$$\text{If } (f(I), s) \in S \text{ then } (I, g(I, s)) \in R.$$

This means that if s is a solution to the instance $f(I)$ in S , we can compute a solution $g(I, s)$ to an instance I in R

Saying that *one can reduce R onto S* can be understood as saying that *if one can solve S efficiently, then one can solve R efficiently*.

Notice that many-to-one reductions map *instances* to *instances*, if we instead assume that we can compute a solution, we use a *Turing reduction*, which we introduce analogously to the classical Turing reduction.

Definition 2.6 — Turing Reduction.

For two problems $R, S \in \mathbf{TFNP}$, we say that R *Turing reduces* to S if a polynomial-time oracle Turing machine that solves R given access to an oracle for S exists.

An *oracle* is a black-box which solves S .

2.1.3 Promise Problems

We have defined total search problems as problems where a solution always exists for *any* input in $\{0, 1\}^*$. However, in practice, we often study problems where a solution is guaranteed to exist only for a subset of the inputs. For instance, every convex function has a global minimum, but this existence result relies on the fact that we are given a convex function. This leads us to the notion of *promise problems* as introduced in [17]. Formally, we restrict the instance space to some subset $\mathcal{X} \subset \{0, 1\}^*$. We only require our algorithm to solve the problem for instances in \mathcal{X} , and it can behave arbitrarily on instances outside of \mathcal{X} .

[17]: Hollender (2021), *Structural Results for Total Search Complexity Classes with Applications to Game Theory and Optimisation*

We highlight that formally **TFNP** does not contain promise problems where $\mathcal{X} \neq \{0, 1\}^*$. We still want to study these problems inside the **TFNP**-Framework. There is a trick for restricting the input space to a subset $\mathcal{X} \subset \{0, 1\}^*$, where the language \mathcal{X}

can be decided in polynomial time. For a search problem R on $X \subset \{0,1\}^*$, we can then define the promise problem R' on $\{0,1\}^*$ by adding a solution (I, \star) to R for all $I \in \{0,1\}^* \setminus R$, where \star is some special binary string. Because it can be decided in polynomial time whether an instance is in \mathcal{X} , we can solve R' by checking whether the instance is in \mathcal{X} and then solving R , hence obtaining a problem in **TFNP**.

For example, in this thesis, we use syntactic validation when the instances are a function or a boolean circuit to validate that the given input is indeed an encoding of a function or circuit. This verification can be done in polynomial time [18], and a special binary string can be outputted if this verification fails. For example, this is the case for the TARSKI problem, where the instances are boolean circuits, and the validity of the instances can be checked in polynomial time. For the sake of simplicity, we will assume that this step implicitly when defining **TFNP**-Problems, and allow instances which have an input space $\mathcal{X} \subset \{0,1\}^*$ if it can be validated in polynomial time.

[18]: Greenlaw and Hoover (1998),
Chapter 9 - Circuit Complexity

Additionally to this syntactic validation, we can construct **TFNP**-instances by adding *violations* to the solution space. For example, if we are interested in finding the global minima of convex functions, we can construct a total search problem by:

- (1) Checking syntactically that the input defines a function;
- (2) Adding a violation of convexity to the solution space. Formally, this is done by changing the relation R to ensure that a solution exists for every instance I ; this can be thought of as allowing more solutions.

A violation of convexity is given by a $x, y \in \mathcal{D}_f$, and $t \in \{0,1\}$ such that $tf(x) + (1-t)f(y) < f(tx + (1-t)y)$.

This means we can often construct a **TFNP**-problem starting out with a promise-problem by checking the validity of the input syntactically and adding violations to the solution space. However, it is essential to note that this is only sometimes the case and that constructing a **TFNP** problem from a promise problem can be a non-trivial task. Also, there is no unique way of constructing **TFNP**-Problems from promise problems, and care has to be taken to introduce the studied problem rigorously.

2.1.4 Representation of sets

In this thesis, we will work with sets of the form $S = \{0, \dots, 2^n - 1\}$, which we will denote by $[2^n]$. Notice that this set can be identified with the set of binary strings of length n . We will denote the set of binary strings of length n by $\{0,1\}^n$. Formally, the functions and the model we will use to represent the functions will use the underlying binary strings in $\{0,1\}^n$. We often denote the integer

$x \in [2^n]$ interchangeably with its representation as a binary string.

Similarly, when considering the d -dimensional case, we can represent the set $L = [2^n]^d$, which corresponds to a d -dimensional lattice with side length 2^n , as the set of binary strings of length $n \cdot d$, i.e. $\{0, 1\}^{nd}$. Again, for simplicity, while the underlying functions rely on the binary strings, these naturally correspond to a unique point $(x_1, \dots, x_d) \in [2^n]^d$. We will use both notations interchangeably.

2.1.5 Representation of functions

Now that we have described the sets, we can describe how we represent the functions. We will represent the functions by using so-called boolean circuits. In this section, we will rely on the presentation of boolean circuits described in [18] and refer an interested reader to this source for a more detailed description.

[18]: Greenlaw and Hoover (1998),
Chapter 9 - Circuit Complexity

On a high level, a boolean circuit is a directed acyclic graph, where the nodes are called *gates*, and the edges are called *wires*. The sinks of the graphs are the output gates, and the sources are the input gates. We want to start by defining a gate formally.

Definition 2.7 — Gate.

A gate is a function $g : \{0, 1\}^k \rightarrow \{0, 1\}$, where k is the number of input wires of the gate.

This corresponds to the gate node, having k incoming edges, and one outgoing edge.

In this thesis, we will only consider the following types of gates:

- **AND-gate:** $g(x_1, x_2) = x_1 \wedge x_2$,
- **OR-gate:** $g(x_1, x_2) = x_1 \vee x_2$,
- **NOT-gate:** $g(x) = \neg x$.

Notice that we only consider gates with at most two inputs, as we can always represent a gate with k inputs as a composition of gates with at most two inputs.

Now, we can describe a boolean circuit formally as follows:

Definition 2.8 — Boolean circuit.

A boolean circuit C is a labeled finite directed acyclic graph, where each vertex has a *type* τ , with

$$\tau(v) \in \{\text{INPUT}\} \cup \{\text{OUTPUT}\} \cup \{\text{AND, OR, NOT}\}$$

and with the following properties:

- If $\tau(v) = \text{INPUT}$, then v has no incoming edges. We call these vertices the *inputs gates*.
- If $\tau(v) = \text{OUTPUT}$, then v has one incoming edge. We call these vertices the *output gates*.



Figure 2.1: example of a Boolean circuit with three input and four output gates.

- If $\tau(v) = \text{AND}$, then v has two incoming edges. We call these vertices the *AND-gates*.
- If $\tau(v) = \text{OR}$, then v has two incoming edges. We call these vertices the *OR-gates*.
- If $\tau(v) = \text{NOT}$, then v has one incoming edge. We call these vertices the *NOT-gates*.

The inputs of C are given by a tuple (x_1, \dots, x_k) of distinct input gates. The output of C is given by a tuple (y_1, \dots, y_l) of distinct output gates.

We give an example of a boolean circuit in Figure 2.1. Of course, we now want to use a boolean circuit to represent a function. To do this, we need to define the function computed by a boolean circuit formally.

Definition 2.9 — Computed function of a boolean circuit.

A boolean circuit C with inputs x_1, \dots, x_n and outputs y_1, \dots, y_m computes a function $f : \{0, 1\}^n \rightarrow \{0, 1\}^m$ as follows:

- The input x_i is assigned the value of the i -th bit of the argument to the function.
- Every other vertex v is assigned the value of the gate g of the vertex, applied to the values of the incoming edges of v .
- The i -th bit of the output of the function is the value of the output gate y_i .

In Figure 2.2, we give an example of using a boolean circuit to compute a function, in particular for a function that is a TARSKI instance. From now on, we will formally represent all functions used in problems by boolean circuits.



Figure 2.2: example of how a function $f : \{0,1\}^2 \rightarrow \{0,1\}^2$ (on the top), can be computed using boolean circuits (on the bottom).

2.1.6 Complexity of boolean circuits

Of course, formally, the complexity of a problem is defined in terms of the *size* of the input. This means we also need to define what we mean by the size of a boolean circuit. We will use the following definition:

Definition 2.10 — Size of a boolean circuit.

The size of a boolean circuit C is the number of gates in the circuit.

The size of the boolean circuits is a measure of the input complexity, i.e. it gives us an indication of how many bits we need to represent the input, it also tells us how many computations are made when computing the function output. We also define the depth of a boolean circuit as follows:

Definition 2.11 — Depth of a boolean circuit.

The depth of a boolean circuit C is the length of the longest path from an input gate to an output gate.

The depth of a boolean circuit is a measure of the time complexity of the computation, i.e. it tells us how many time steps are needed to compute the output of the function. This is especially true in a parallel setting, where all gates can be seen as setting off at the same time.

It can be shown that $\text{poly}(\text{size}(n))$ bits suffice to encode any boolean circuit.




Figure 2.3: Example of a LOCALOPT Problem with $n = 3$ (8 vertices). Solid lines represent the circuit S . The valid solutions are colored green.

2.2 Subclasses of TFNP

Complete **FNP** problems within **TFNP** would imply that **NP** = **coNP** [14], a highly unlikely scenario. Consequently, complete problems are not expected within **TFNP**, necessitating alternative approaches to investigate its structure.

TFNP is known as a *semantic* class [19]. This is known to mean that it is unlikely that we can find complete **TFNP** problems [20]. We refer the reader to Papadimitriou's work for a more detailed discussion of these terms and their implications. We want to explore *syntactic* subclasses of **TFNP** to address this challenge. One approach, proposed by Papadimitriou [19], categorizes search problems based on existence proofs confirming their totalness. This basic strategy leads to the detailed study of specific complexity classes discussed in the following sections.

[14]: Megiddo and Papadimitriou (1991), *On total functions, existence theorems and computational complexity*

[19]: Papadimitriou (1994), *Computational complexity*

[20]: Sipser (1982), *On relativization and the existence of complete sets*

2.2.1 Polynomial Local Search (PLS)

The existence result which gives rise to **PLS** is:

"Every directed acyclic graph has a sink."

We can then construct the class **PLS** by defining it as all problems which reduce to finding the sink of a directed acyclic graph (DAG). Formally we first define the problem LOCALOPT as in [2]:

LOCALOPT

Input: Two boolean circuits $S, V : [2^n] \rightarrow [2^n]$.

Output: A vertex $v \in [2^n]$ such that $P(S(v)) \geq P(v)$.

[2]: Johnson et al. (1988), *How easy is local search?*

S can be seen as a proposed successor, and V as a potential. The goal is to find a local minima v of the potential.

Let us discuss why solving a LOCALOPT instance is equivalent to finding the sink of a DAG. The circuit S defines a directed graph, which might contain cycles. Only keeping the edges on which the potential decreases (strictly) leads to a DAG, with as sinks exactly the v such that $P(S(v)) \geq P(v)$. We give an example of a LOCALOPT instance in Figure 2.3. Now we can define **PLS**:

Definition 2.12 — Polynomial Local Search (PLS).

The class **PLS** is the set of all **TFNP** problems that reduce to **LOCALOPT**.

Studying “simple” problems such as **PLS** is particularly insightful because we strongly believe that these problems cannot be solved by any method more efficiently than simply traversing the graph, even though there might be a very clever way of analyzing the input circuit, which leads to a quicker result. Hence, given a graph of exponentially large size, it appears highly improbable that an efficient solution can be found. Therefore, all problems in **PLS** inherently embody the fundamental challenge of not being able to surpass the basic strategy of navigating through the directed acyclic graph. Of course — and here lies the difficulty of complexity theory — we cannot prove this statement; it could be that some very clever analysis of the boolean circuits could lead to an efficient algorithm for finding sinks of exponentially large directed acyclic graphs.

2.2.2 Polynomial Parity Argument on Directed Graphs (PPAD)

Now we want to discuss the complexity class **PPAD**, introduced by Papadimitriou [1] as one of the first syntactic subclasses of **TFNP**. The existence result giving rise to this class is:

“If a directed graph has an unbalanced vertex, then it has at least one other unbalanced vertex.”

PPAD can be defined using the problem **END-OF-LINE** as introduced in [15].

END-OF-LINE (EOL)

Input: Boolean circuit $S, P : \{0, 1\}^n \rightarrow \{0, 1\}^n$ such that $P(0^n) = 0^n \neq S(0^n)$ (0^n is a source.)

Output: An $x \in \{0, 1\}^n$ such that either:

- $P(S(x)) \neq x$ (x is a sink) or
- $S(P(x)) \neq x \neq 0^n$ (x is a non non-standard source)

These boolean circuits represent a directed graph with maximal in and out-degree of one by having an edge from x to y if and only if $S(x) = y$ and $P(y) = x$. The goal is to find a sink in the graph or another source. It can be shown that the general case of finding a second imbalanced vertex in a directed graph (a problem called **IMBALANCE**) can be reduced to **END-OF-LINE** [21]. Now we can define the complexity class **PPAD** as follows:

[1]: Papadimitriou (1994), *On the complexity of the parity argument and other inefficient proofs of existence*

[15]: Daskalakis et al. (2009), *The Complexity of Computing a Nash Equilibrium*

Here, S can be thought of as giving the successor of a vertex, and P as giving the predecessor of a vertex.

Notice that **END-OF-LINE** allows cycles and that these do not induce solutions.

[21]: Goldberg and Hollender (2021), *The Hairy Ball problem is PPAD-complete*




Figure 2.4: Example of an END-OF-LINE Problem with $n = 3$ (8 vertices). Solid lines represent the circuit S , and dashed lines represent the circuit P . The solutions are the sinks $x = 5$, $x = 7$ and $x = 1$, as well as the sources $x = 4$ and $x = 6$.

Definition 2.13 — PPAD.

The class **PPAD** is the set of all **TFNP** problems that reduce to END-OF-LINE.

2.2.3 End of Potential Line (EOPL)

Next, we discuss the complexity class **EOPL** introduced in [3]. The existence results giving rise to **EOPL** is:

[3]: Fearnley et al. (2018), *End of Potential Line*

“In a directed acyclic graph, there must be at least two unbalanced vertices.”

Similarly to **PLS**, acyclicity will be enforced using a potential.

END OF POTENTIAL LINE

Input: Two boolean circuits $S, P : \{0, 1\}^n \rightarrow \{0, 1\}^n$, and a boolean circuit $V : \{0, 1\}^n \rightarrow [2^n - 1]$, such that 0^n is a source, (i.e. $P(0^n) = 0^n \neq S(0^n)$).

Output: An $x \in \{0, 1\}^n$ such that either:

- ▶ $P(S(x)) \neq x$ (x is a sink)
- ▶ $S(P(x)) \neq x \neq 0^n$ (x is a non-standard source)
- ▶ $S(x) \neq x$, $P(S(x)) = x$ and $V(S(x)) \leq V(x)$ (violation of the monotonicity of the potential)

Here, S can be thought of as giving the successor of a vertex, and P as giving the predecessor of a vertex. V is a potential that is supposed to increase monotonously along the line.

S and P can be thought of as representing a directed line. Finding another source (a non-standard source) is a violation, as a directed line only has one source. The potential serves as a guarantee of acyclicity. Now, we can define the complexity class **EOPL**.

Definition 2.14 — EOPL.

The class **EOPL** is the set of all **TFNP** problems that reduce to END OF POTENTIAL LINE.




Figure 2.5: Example of an EOPL Problem with $n = 3$ (8 vertices). Solid lines represent the circuit S , and dashed lines represent the circuit P . The solutions are the sink $x = 7$, the violation of potential at $x = 5$, and the non-standard source $x = 4$.

2.3 The TARSKI Problem

2.3.1 Definition of the TARSKI Problem

Next, we introduce the TARSKI Problem. Before we do this, we recall that there is a partial order on the d dimensional lattice $[N]^d$, given by $x \leq y$ if and only if $x_i \leq y_i$ for all $i \in \{1, \dots, d\}$. We can now define functions on this lattice, and in particular, we can define monotone functions.

Definition 2.15 — Monotone function.

A function $f : [N]^d \rightarrow [N]^d$ is *monotone* if for all $x, y \in [N]^d$ we have $x \leq y$ implies $f(x) \leq f(y)$.

Notice that $x \not\leq y$ does *not* imply $x \geq y$. In particular, two points are not always comparable.

Such functions are also called *order preserving* functions in the literature.

The TARSKI-problem originates from Tarski's fixed point Theorem, introduced in [4]. In our setting the Theorem states the following:

Theorem 2.16 — Tarski's fixed point Theorem.

Let $f : [N]^d \rightarrow [N]^d$ a function on the d -dimensional lattice. If f is monotone (for the previously discussed partial order), then f has a fixed point, i.e. there is an $x \in [N]^d$ such that $f(x) = x$.

[4]: Tarski (1955), *A lattice-theoretical fixpoint theorem and its applications*.

This Theorem is also known as the Knaster–Tarski Theorem in the literature.

A proof of this Theorem can be found in the previously mentioned work [4]. Without surprise, the TARSKI problem, defined in [5], is now to find such a fixed point. Formally, we define the problem as follows:

[5]: Etesami et al. (2020), *Tarski's Theorem, Supermodular Games, and the Complexity of Equilibria*

TARSKI

Input: A boolean circuit $f : [N]^d \rightarrow [N]^d$.

Output: Either:

- An $x \in [N]^d$ such that $f(x) = x$ (fixed point) or
- $x, y \in [N]^d$ such that $x \leq y$ and $f(x) \not\leq f(y)$ (violation of monotonicity).




Figure 2.6: Example of a 2 dimensional TARSKI instance. A fixed point is located at $x = (4, 5)$. The path to the fixed point which Algorithm 1 finds is colored in green.

This is a total search problem, as there will always either be a fixed point or a point violating monotonicity. We give an example of a two-dimensional TARSKI instance in Figure 2.6. Before we discuss the location of TARSKI in the **TFNP** landscape and two known algorithms for solving TARSKI, we want to discuss a useful Lemma, which allows us to simplify the study of Tarski instances. The definition of TARSKI instances allows for the image of a point to be located anywhere in the lattice; we will show that we can reduce to the cases where the image of a point is in the immediate neighborhood of the point.

Lemma 2.17 — Simplifying TARSKI.

Let $f : [2^n]^d \rightarrow [2^n]^d$ be a TARSKI instance on a complete lattice $[2^n]^d$. Consider $\tilde{f} : [2^n]^d \rightarrow [2^n]^d$ given by:

$$\tilde{f}(x)[i] = \begin{cases} x[i] + 1 & \text{if } f(x)[i] > x[i], \\ x[i] & \text{if } f(x)[i] = x[i], \\ x[i] - 1 & \text{if } f(x)[i] < x[i]. \end{cases} \quad \text{for all } i \in \{1, \dots, d\}$$

Then, for any two points $x, y \in [2^n]^d$, $f(x) \leq f(y)$ if and only if $\tilde{f}(x) \leq \tilde{f}(y)$.

Notice that given a circuit C which computes f , we can construct a circuit \tilde{C} which computes \tilde{f} by adding $\mathcal{O}(\text{poly } d)$ gates to C . This means that both problems are equivalent in terms of complexity.

Proof. The lemma follows directly by observing that for all $i \in \{1, \dots, d\}$ we have: $f(x)[i] \leq f(y)[i]$ if and only if $\tilde{f}(x)[i] \leq \tilde{f}(y)[i]$. \square

This means that in this thesis, we can consider the simplified version of the TARSKI problem, where for every $x \in [2^n]^d$, we have $\|x - f(x)\|_\infty \leq 1$, which we will implicitly assume from now on.

Also note that given a circuit that computes f , we can construct a circuit with at most $\mathcal{O}(\text{poly } d)$ additional gates which computes \tilde{f} .

2.3.2 Two algorithms for solving TARSKI

We briefly discuss the most common algorithms for solving TARSKI instances. We begin with a straightforward algorithm, which is based on the following observation:

Remark 2.18

Let f be a TARSKI instance on a complete lattice L . If f is monotone, and for some $x \in L$ we have $f(x) \geq x$, then $f(f(x)) \geq f(x)$.

Now, note that by starting at the point $\mathbf{0} = 0^d$ and iterating the function f , we will eventually reach a fixed point. This means that we can construct an iterative algorithm for solving TARSKI, as described in Algorithm 1.

Algorithm 1: Iterative Algorithm for TARSKI

Data: A boolean circuit $f : L \rightarrow L$

Result: A fixed point of f

```

 $x \leftarrow \mathbf{0}$  ;
while  $f(x) \neq x$  do
  if  $f(x) \not\geq x$  then
    return " $x_{\text{old}}, x$  are a violation of monotonicity." ;
  else
     $x_{\text{old}} \leftarrow x$   $x \leftarrow f(x)$  ;
return  $x$  ;

```

The path which Algorithm 1 takes to solve a TARSKI instance is colored green in Figure 2.6. While Algorithm 1 might not be very efficient — it runs in worst-case time $\mathcal{O}(d \cdot N)$ for $L = [N]^d$ — it does have some theoretical applications for locating TARSKI inside **TFNP**. Previous work [5] showed that TARSKI lies in **PLS** by considering the set of possible states of the previously described algorithm, together with a potential function given by $V(x) = \sum_{i=1}^d x[i]$, and showing that this potential is monotone along the states of the algorithm. The circuit S associates to the state of the algorithm the next state it will be in.

[5]: Etessami et al. (2020), *Tarski's Theorem, Supermodular Games, and the Complexity of Equilibria*

Next, we describe a more advanced algorithm, due to [10], for solving TARSKI instances and also give an alternative presentation and simplified proof of its correctness. The following notation aims to make the argument as clear as possible. For a given complete lattice $L = [N_1] \times \dots \times [N_d]$ and some dimension $x \in L$ we define the following sublattices:

$$L_{\leq x} = [x[1] + 1] \times \dots \times [x[d] + 1],$$

$$L_{\geq x} = \llbracket x[1], N_1 \rrbracket \times \dots \times \llbracket x[d], N_d \rrbracket$$

[10]: Dang et al. (2020), *Computations and Complexities of Tarski's Fixed Points and Supermodular Games*

We denote by $\llbracket a, b \rrbracket$ the set of whole numbers $\{a, a + 1, \dots, b\}$.

and for a given dimension $k \in \{1, \dots, d\}$ and $K \in [N_k]$, we define the following sublattices:

$$L_{k < K} = [N_1] \times \dots \times [N_{k-1}] \times [K] \times [N_{k+1}] \times \dots \times [N_d],$$

$$L_{k = K} = [N_1] \times \dots \times [N_{k-1}] \times \{K\} \times [N_{k+1}] \times \dots \times [N_d],$$

$$L_{k > K} = [N_1] \times \dots \times [N_{k-1}] \times \{K + 1, \dots, N_k\} \times [N_{k+1}] \times \dots \times [N_d].$$

The algorithm — and in particular, our proof of the correctness — is based on the following observation:

Remark 2.19

Let $L = [N_1] \times \dots \times [N_d]$ be a complete lattice and $f : L \rightarrow L$ a monotone function. Then:

- (1) If for some $x \in L$ we have $f(x) \leq x$, then f has a fixed point in $L_{\leq x}$.
- (2) If for some $x \in L$ we have $f(x) \geq x$, then f has a fixed point in $L_{\geq x}$.

Proof. Let $x \in L$ such that $f(x) \leq x$. Then for all $y \in L_{\leq x}$ we have $y \leq x$ and hence $f(y) \leq f(x) \leq x$, which shows that f is a TARSKI instance on $L_{\leq x}$. By Tarski's fixed point Theorem, f has a fixed point in $L_{\leq x}$. The proof for the second point is analogous. \square

Hence, points with these properties seem particularly interesting when searching for fixed points of f . Hence, we want to give them a name:

Definition 2.20 — Progress point.

Let $f : L \rightarrow L$ a TARSKI function. We call a point $x \in L$ a *progress point* if $f(x) \leq x$ or $f(x) \geq x$.

The lattice's smallest vertex and largest vertex are always progress points.

This means that if we have a progress point, we can reduce the area where we need to search for a fixed point. The question now becomes: how do we find such an x ? The algorithm we will present is based on the following observation:

Remark 2.21

Let $f : L \rightarrow L$ on a complete lattice $L = [N_1] \times \dots \times [N_d]$, for a monotone function f , be a TARSKI instance. By fixing some dimension $k \in \{1, \dots, d\}$, we can define the function $f_{k=K} : L_{k=K} \rightarrow L_{k=K}$ as follows:

$$f_{k=K}(x)[i] = \begin{cases} f(x)[i] & \text{if } i \neq k, \\ K & \text{if } i = k. \end{cases} \quad \text{for all } i \in \{1, \dots, d\}$$

Then $f_{k=K}$ is a monotone TARSKI instance on $L_{k=K}$, and if x^* is a fixed point of $f_{k=K}$, then x^* is a progress point of f .

If we can solve a $d - 1$ dimensional TARSKI instance, we can find a point x such that $f(x) \geq x$ or $f(x) \leq x$.

Proof. The monotonicity of $f_{k=K}$ follows directly from the monotonicity of f .

The fact that x^* is a progress point follows from the fact that if x^* is a fixed point of $f_{k=K}$, then $f(x^*)[i] = x^*[i]$, for all $i \neq k$. This means that if $f(x^*)[k] \leq x^*[k]$, then $f(x^*) \leq x^*[k]$ and if $f(x^*)[k] \geq x^*[k]$, then $f(x^*) \geq x^*[k]$. \square

By choosing $K = \lfloor \frac{N_k}{2} \rfloor$ we can find a progress point x such that both $L_{\leq x}$ and $L_{\geq x}$ have at most half the size of L . This means we can reduce the search space by a factor of at least two by solving a $d - 1$ dimensional TARSKI instance. We can solve a d dimensional TARSKI instance by repeatedly solving $d - 1$ dimensional TARSKI instances and reducing the search space size by a factor of at least 2 in each step. This means we can solve a d dimensional TARSKI instance by combining a $d - 1$ dimensional TARSKI solver and a binary search. The $d - 1$ dimensional instances can be solved recursively. We give the recursive algorithm for solving TARSKI instances in Algorithm 2.

A simple analysis shows that this algorithm runs in $\mathcal{O}(\log^d N)$ for $L = [N]^d$. It was conjectured that this is an optimal algorithm for TARSKI [5]. This turned out not to be true, as a better algorithm was developed [22], which mostly relies on a faster way of finding a progress point in the three-dimensional case, which they call the *inner algorithm*. An *outer algorithm* then repeatedly applies the inner algorithm on 3-dimensional instances. Overall this approach achieves a query complexity of $\mathcal{O}(\log^{2\lceil \frac{d}{3} \rceil} N)$ for $L = [N]^d$.

A further advance using more advanced methods were made in [11]. They achieve a query complexity of $\mathcal{O}(\log^{\lceil \frac{d-1}{2} \rceil} N)$ for

[5]: Etessami et al. (2020), *Tarski's Theorem, Supermodular Games, and the Complexity of Equilibria*

[22]: Fearnley et al. (2022), *A Faster Algorithm for Finding Tarski Fixed Points*

[11]: Chen and Li (2022), *Improved Upper Bounds for Finding Tarski Fixed Points*

Algorithm 2: Recursive Algorithm for TARSKI

```

Function RecursiveTarskiSolver( $f: L \rightarrow L, d$ ):
  /* Binary search in the  $d$ -th dimension */
   $l \leftarrow 0, r \leftarrow N_d$ ; /* The search space is  $[l, r]$  */
  while  $r - l > 1$  do
     $m \leftarrow \lfloor \frac{l+r}{2} \rfloor$ ; /* Middle of the interval */
    if  $d = 1$  then
       $x^* \leftarrow m$ 
    else
      /* Solve the  $d-1$  dimensional instance */
       $x^* \leftarrow \text{RecursiveTarskiSolver}(f_{d=m}, d-1)$ ;
      if  $f(x^*)[d] \leq x^*[d]$  then
         $r \leftarrow m$ 
      else
         $l \leftarrow m$ 
  return  $x^*$ 

```

$L = [N]^d$. This is to date the best known bound for solving TARSKI instances.

2.3.3 Lower bounds for TARSKI

The best-known lower bounds for TARSKI are given by [5]. They showed that in the black-box model, where the only way to access the function f is by querying it, solving a d -dimensional TARSKI requires solving at least $\Omega(\log N)$ one-dimensional TARSKI instances, which are as difficult as binary search, hence this means that solving a d -dimensional TARSKI instance requires at least $\Omega(\log^2 N)$ queries. This means that the upper and lower bounds are equal in the two- and three-dimensional cases, but in all other cases, there remains a gap. In particular, the best-known lower bound for solving TARSKI does not depend on the dimension d , which seems somewhat unexpected.

[5]: Etessami et al. (2020), *Tarski's Theorem, Supermodular Games, and the Complexity of Equilibria*

This gives us reason to study TARSKI under the lens of complexity theory, in particular to understand where TARSKI lies in the **TFNP** landscape.

2.3.4 Location of TARSKI in TFNP

Next we summarize where TARSKI lies inside of **TFNP**. It has been shown in [5] that TARSKI lies in **PLS** as we discussed when presenting Algorithm 1. The same paper showed that TARSKI lies \mathbf{P}^{PPAD} . We will provide an alternative proof of this second fact in Chapter 3. Previous work [23] showed that many-to-one reductions and Turing-reduction onto **PPAD** are equivalent. In

[23]: Buss and Johnson (2012), *Propositional proofs and reductions between NP search problems*

particular this means that $\mathbf{P}^{\text{PPAD}} = \text{PPAD}$, and that TARSKI lies in PPAD.

Now that we have established that TARSKI lies inside $\text{PLS} \cap \text{PPAD}$, we want to discuss the structure of $\text{PLS} \cap \text{PPAD}$ and describe recent advances in the study of this class. There have been two surprising advances in the study of $\text{PLS} \cap \text{PPAD}$ in the last few years. The first is that $\text{CLS} = \text{PLS} \cap \text{PPAD}$ [24]. CLS (Continuous Local Search) was first introduced by Daskalakis and Papadimitriou in [16] and can be informally thought of as the class of all problems that can be solved by finding the local optimum of a potential in a discrete space equipped with an adjacency relation. This result shows that the problems in $\text{PLS} \cap \text{PPAD}$ are exactly those that gradient descent algorithms can solve.

[24]: Fearnley et al. (2023), *The Complexity of Gradient Descent: CLS = PPAD \cap PLS*

[16]: Daskalakis and Papadimitriou (2011), *Continuous Local Search*

A further notable collapse is the result $\text{PLS} \cap \text{PPAD} = \text{EOPL}$, which was only recently shown in [12]. This means that TARSKI lies in EOPL. A question that then arises, and which this thesis will attempt to answer, is whether we can construct an explicit reduction of TARSKI to ENDOFPOTENTIALLINE.

[12]: Goos et al. (2022), *Further Collapses in TFNP*

2.3.5 Variants of TARSKI

Before we conclude this chapter, we want to discuss some variants of the TARSKI problem. The first variant we introduce is the *promise* variant of TARSKI, which is defined as follows:

PROMISETARSKI

Input: A boolean circuit $f : [N]^d \rightarrow [N]^d$ such that $x \leq y \implies f(x) \leq f(y)$

Output: A fixpoint x of f .

Notably in this variant of TARSKI we are *promised* that we have a monotone function, and hence we do not allow violations of monotonicity as solutions (as these should not be possible).

Next we introduce two variants of TARSKI which are also promise variants of TARSKI. Instead of promising that the function is monotone, we are promised some properties related to uniqueness of solutions. The first variant is the *unique fixpoint* variant of TARSKI, which is defined as follows:

UNIQUETARSKI

Input: A boolean circuit $f : [N]^d \rightarrow [N]^d$ such that f is monotone and there is a unique fixpoint x

Output: The unique fixpoint x of f .

A very notable recent result, is that the query complexity of **UNIQUETARSKI** is equal to the query complexity of **TARSKI** [25]. We also introduce an even stronger uniqueness variant of **TARSKInext**. This time the promise is that when fixing a subset of coordinates, the induced function has a unique fixpoint, i.e. $\tilde{f} = f_{d_1=K_1, \dots, d_k=K_k}$ has a unique fixpoint for any choice of $d_1, \dots, d_k \in \{1, \dots, d\}$ and $K_1, \dots, K_k \in [N]$.

[25]: Chen et al. (2023), *Reducing Tarski to Unique Tarski (In the Black-Box Model)*

SUPERUNIQUETARSKI

Input: A boolean circuit $f : [N]^d \rightarrow [N]^d$ such that f is monotone and when fixing a subset of coordinates the induce function \tilde{f} has a unique fixpoint.

Output: The unique fixpoint x of f .

The problem **SUPERUNIQUETARSKI** is a very strong promise variant of **TARSKI**, it implies that Algorithm 2 find a unique fixpoint in every step.

Reducing TARSKI to PPAD

3

This chapter explores the membership of TARSKI in the complexity class **PPAD**. We begin by presenting a high-level overview of an established proof of the reduction of this problem to BROUWER [5]. We subsequently introduce a novel problem, TARSKI*, which facilitates a divide-and-conquer approach to solving TARSKI by leveraging the structure of the function f . This new formulation allows us to provide an alternative proof of TARSKI's membership in **PPAD** using *Sperner's Lemma* instead of the traditional *Brouwer's Fixed Point Theorem*. This approach simplifies the proof and sets the stage for further reduction of TARSKI* to **EOPL** in the subsequent chapter.

3.1 Presentation of the known reduction of TARSKI to PPAD

We want to give a high-level presentation of the proof of TARSKI membership in **PPAD** from [5], which will help us motivate the introduction of TARSKI* and the subsequent use of *Sperner's Lemma*. The proof given by Etessami et al. relies on *Brouwer's fixed point theorem*, which we introduce below.

Theorem 3.1 — Brouwer's fixed point theorem.

Let $K \subset \mathbb{R}^d$ be a compact, convex set. Then every continuous function $f : K \rightarrow K$ has a fixed point $x^* \in K$, i.e. $f(x^*) = x^*$.

The original proof can be found in [26], and a more straightforward proof relying on SPERNER'S LEMMA can be found in [27]. This Theorem gives rise to a total search problem which we call BROUWER:

BROUWER

Input: A continuous function $f : K \rightarrow K$.

Output: A fixed point $x^* \in K$ such that $f(x^*) = x^*$.

The problem BROUWER was first introduced and shown to be **PPAD**-complete in [28], meaning that it suffices to reduce TARSKI to BROUWER in order to show that TARSKI is in **PPAD**. We will reduce TARSKI to, at most polynomially, many instances of BROUWER, which will allow us to show that TARSKI is in \mathbf{P}^{PPAD} . Overall, we will construct a Turing reduction of TARSKI to BROUWER, which suffice as **PPAD** is closed under Turing reductions [23].

3.1	Known reduction to PPAD	23
3.2	Introducing TARSKI* . .	24
3.3	Sperner's Lemma . . .	26
3.3.1	on Simplices	27
3.3.2	on Lattices	28
3.3.3	SPERNER to ENDOFFLINE reduction	29
3.4	Reducing TARSKI* to SPERNER	30

[5]: Etessami et al. (2020), *Tarski's Theorem, Supermodular Games, and the Complexity of Equilibria*

[5]: Etessami et al. (2020), *Tarski's Theorem, Supermodular Games, and the Complexity of Equilibria*

[26]: Brouwer (1911), *Über Abbildung von Mannigfaltigkeiten*

[27]: Aigner and Ziegler (2018), *Proofs from THE BOOK*

We leave out the technical detail of how this function is given using boolean circuits and how precise the output needs to be, as it is irrelevant for this high-level presentation.

[28]: Papadimitriou (1994), *On the complexity of the parity argument and other inefficient proofs of existence*

[23]: Buss and Johnson (2012), *Propositional proofs and reductions between NP search problems*

The reduction extends the discrete function f to a function $\tilde{f} : [0, 2^n - 1]^d \rightarrow [0, 2^n - 1]^d$, such that \tilde{f} interpolates the lattice function f , is continuous and piecewise linear between lattice points, and hence continuous. The authors achieve this by using a simplicial decomposition of each cell of the lattice. Now we have an instance of BROUWER, and hence, we can find a fixed point x^* of \tilde{f} . Of course, this fixed point does not need to be *integral*. The critical insight is that we can use this fixed point to reduce the search area for an integral fixed point by at least half or find a violation of monotonicity. In particular, either there is a fixed point in both $\{x \in [2^n]^d : x \geq x^*\}$ and $\{x \in [2^n]^d : x \leq x^*\}$, or there is a violation of monotonicity in the cell containing x^* . We can repeat this procedure, always halving the search area, which allows us to solve a TARSKI instance using at most $\mathcal{O}(d \cdot n)$ calls to BROUWER.

We call a point *integral* if it belongs to the original lattice.

3.2 Introducing TARSKI*

In the previous Section, we have seen that TARSKI can be reduced to a polynomial number of BROUWER instances. We want to study a single such reduction to give an alternative proof that TARSKI is in PPAD. In order to do this, we introduce a new problem, TARSKI*. This problem can be thought of as a subproblem towards solving TARSKI. A standard strategy to solve TARSKI is to use a *divide-and-conquer* strategy, for instance, used in [5], and presented previously in Algorithm 2, of Subsection 2.3.2. We want to construct a problem that allows us to divide the TARSKI problem into two smaller problems, where solving the smaller of the two leads to a solution.

[5]: Etessami et al. (2020), *Tarski's Theorem, Supermodular Games, and the Complexity of Equilibria*

For the sake of generality and in order to achieve more precise proofs in the following, we introduce the problem on the integer lattice $L = N_1 \times \dots \times N_d$, such that $N_i \leq 2^n$ for all $i \in \{1, \dots, d\}$. We propose the following problem:

TARSKI*

Input: A boolean circuit $f : L \rightarrow L$.

Output: Either:

- Two points $x, y \in L$ such that $\|x - y\|_\infty \leq 1$, $x \leq f(x)$ and $y \geq f(y)$, or;
- A violation of monotonicity: Two points $x, y \in L$ such that $x \leq y$ and $f(x) \not\leq f(y)$.

We want to show that TARSKI* is, in a sense, a subproblem of TARSKI.

Claim 3.2

An instance of TARSKI can be solved using $\mathcal{O}(d \cdot n)$ calls to TARSKI* and up to $\mathcal{O}(d)$ additional function evaluations.

Proof. We will show that we can use a single call of TARSKI* to either find a violation of monotonicity, a fixpoint, or an instance of TARSKI which has at most half as many points and must contain a solution. Let x, y be the two points which a Turing machine solving TARSKI* on a function f outputs. We proceed by case distinction:

Case 1: We are done if $f(x) = x$ or $f(y) = y$ because we have found a fixpoint.

Case 2.1: If $x < y$ and $f(x) \not\leq f(y)$, we have a violation of monotonicity, which solves the given TARSKI instance.

Case 2.2: If $x < y$ and $f(x) \leq f(y)$, we claim that we can solve the TARSKI instance in $\mathcal{O}(\|x - y\|_1)$ additional function calls. Notice that we have $\|x - y\|_\infty \leq 1$. Now notice that because $f(x) > x$ (if not, see case 1), there is at least one dimension $i \in \{1, \dots, d\}$ such that $f(x)[i] > x[i]$. Also notice that in this dimension i if $f(y)[i] < y[i]$, then because $|x[i] - y[i]| \leq \|x - y\|_\infty \leq 1$, we would have a violation of the monotonicity of f in this dimension. Therefore, we must have $f(y)[i] = y[i]$. The same argument shows that if in any dimension j we have $f(y)[j] < y[j]$, then $f(x)[j] = x[j]$. Therefore, we know that because there must be at least one such dimension i and j , we have:

$$\|f(x) - f(y)\|_\infty \leq \|x - y\|_\infty \leq 1 \text{ and } \|f(x) - f(y)\|_1 \leq \|x - y\|_1 - 2$$

Hence, we can now repeat the same argumentation with $f(x)$ and $f(y)$, and we can do this at most $\mathcal{O}(\|x - y\|_1)$ times until we find a violation of monotonicity or a fixpoint. Because $\|x - y\|_1 \leq d$, this will take at most $\mathcal{O}(d)$ additional steps.

Case 3: If $x \not\leq y$, then we can partition the set of lattice points into two sets S_x and S_y , as follows:

$$S_x = \{z \in L : z \geq x\} \quad \text{and} \quad S_y = \{z \in L : z \leq y\}.$$

These two sets are disjoint: if there was a $z \in S_x \cap S_y$, then $x \leq z \leq y$, which would imply $x \leq y$, which is a contradiction. We will show that S_x must contain a solution to the TARSKI instance. If for some $z \in S_x$, we have $f(z) \notin S_x$, then we have $f(z) \not\leq f(x)$, which means that z and x form a violation of monotonicity. This means that S_x forms a new valid instance of TARSKI. By the same argumentation, S_y also forms a valid instance of TARSKI and hence, it suffices to solve the smaller of the two instances recursively. In particular, because they are disjoint, one of the instances S_x or S_y contains less than half of the lattice points of L , and hence we can solve the instance in $\mathcal{O}(\log 2^{dn}) = \mathcal{O}(d \cdot n)$ calls of TARSKI*.

□

Now that we know that TARSKI* is a good stepping stone towards solving TARSKI, we want to investigate why TARSKI* lies in PPAD.

3.3 Sperner's Lemma

The preceding discussion hinges on the assumption that TARSKI* is a total problem, implying that every instance of the problem is guaranteed a solution. This Section will substantiate this claim, establishing Tarskistar's classification within TFNP. Rather than employing *Brouwer's fixed point Theorem* — a cornerstone of continuous topology — we pivot to its discrete analog, *Sperner's Lemma*, a foundational result in combinatorial topology. This approach is particularly apt for two main reasons:

- We are working on a discrete lattice, so it seems more natural to use a discrete tool.
- Papadimitriou proved that BROUWER is PPAD-complete by reducing BROUWER to SPERNER [28]. Hence, by reducing to BROUWER, we introduce continuity into the problem, which is unnecessary, as it gets removed again behind the scenes.

[28]: Papadimitriou (1994), *On the complexity of the parity argument and other inefficient proofs of existence*

We aim to apply *Sperner's Lemma* on the integer lattice. Using this tool is not directly possible, as *Sperner's Lemma* is defined on a simplicial decomposition of a simplex. Hence, we will first introduce *Sperner's Lemma* for simplices and then show how it can be adapted to work on an integer lattice.

3.3.1 Sperner's Lemma for Simplices

Before we introduce the Lemma itself, we want to define the setting of the result. We consider a d -dimensional simplex with vertices v_0, v_1, \dots, v_d . We now consider a *simplicial subdivision* of this simplex, meaning that we partition the simplex into smaller simplices. We give an example of such a partition in Figure 3.1 in the 3-dimensional case.

By d dimensional simplex we mean the convex Hull of these $d+1$ points in \mathbb{R}^d




Figure 3.1: Setup for SPERNER'S LEMMA in the 3-dimensional case. The large simplex spanned by v_0, v_1, v_2, v_3 is subdivided into smaller simplices.

Now we introduce a coloring c of the vertices of this subdivision with colors $\{0, 1, \dots, d\}$. We want to enforce that the vertices v_i of the large simplex are colored with color i and that the vertices on a subsimplex $\{v_{i_0}, \dots, v_{i_k}\}$ are colored with colors i_0, \dots, i_k . We give an example of such a coloring in 2 dimensions in Figure 3.2.

We now introduce Sperner's Lemma, which was first proven in [29], and for which a more modern proof can be found in [27].

Theorem 3.3 — Sperner's Lemma.

Suppose a d -dimensional simplex with vertices v_0, \dots, v_d is subdivided into smaller simplices. Now color every vertex with a color $\{0, \dots, d\}$ such that v_i is colored i , and the vertices on a subsimplex $\{v_{i_0}, \dots, v_{i_k}\}$ are colored with colors i_0, \dots, i_k . Then, there is a subsimplex with vertices of every color.

[29]: Sperner (1928), *Neuer beweis für die invarianz der dimensionszahl und des gebietes*

[27]: Aigner and Ziegler (2018), *Proofs from THE BOOK*

We give an example of a 2-dimensional simplex, subdivided into smaller simplices, and colored according to *Sperner's Lemma* in Figure 3.2.




Figure 3.2: Example of SPERNER'S LEMMA in the two-dimensional case, with three colors: orange (0), purple (1), and blue (2). The subsimplex spanned by v_0 and v_1 only contains blue and purple vertices, the subsimplex spanned by v_1 and v_2 contains only purple and blue vertices, and the subsimplex spanned by v_0 and v_2 contains only orange and blue vertices. *Sperner's Lemma* implies that there must be a subsimplex (colored in green) containing all colors.

3.3.2 Sperner's Lemma for an integer lattice

In the previous Section, we introduced *Sperner's Lemma* for an integer lattice. This enables us to find a region of a colored lattice that contains all colors under certain conditions. Instead of a subsimplex, we look for a *cell*¹ of the lattice, which contains all colors.

For this, we proceed as follows: We take the d -dimensional lattice $L = [N_1] \times \dots \times [N_d]$, we subdivide each cell into simplices². We set $v_0 = (0, \dots, 0)$, $v_1 = (N_1 - 1, 0, \dots, 0)$, \dots , $v_d = (0, \dots, 0, N_d - 1)$. We give an example of such a subdivision in the 3-dimensional case in Figure 3.3. Notice that we can deform the lattice and obtain an equivalent simplex and a simplicial decomposition of this simplex.

This means that under the appropriate conditions — which we will detail next — we can apply *Sperner's Lemma* to the lattice. Assume that we color all vertices of the lattice with colors $\{0, \dots, d\}$, such that v_i is colored i , and every vertex x with $x[i] = 0$, is *not* colored i for $i \in \{1, \dots, d\}$. Then, we can apply *Sperner's Lemma* to this simplicial decomposition of the lattice, and we will find a simplex that contains all colors. Of course, because every subsimplex is included in exactly one cell by construction, there must be a cell that contains all colors. This motivates the definition of the total problem SPERNER, which was introduced in [28]. We introduce the problem for a general lattice $L =$

1: By cell, we mean a unit hypercube of the integer lattice

2: How this we do this is not relevant in this chapter but will be discussed in the next chapter.

[28]: Papadimitriou (1994), *On the complexity of the parity argument and other inefficient proofs of existence*




Figure 3.3: Example of the simplicial decomposition of a lattice in the three-dimensional case on the left, and the equivalent simplicial decomposition on the right of a simplex v_0, v_1, v_2, v_3 .

$N_1 \times \dots \times N_d$, such that $N_i \leq 2^n$.

SPERNER

Input: A coloring $c : L \rightarrow \{0, \dots, d\}$ of the vertices of L , such that for every $i \in \{0, \dots, d\}$ the vertices $\{x \in L : x[i] = 0\}$ are not colored i .

Output: A cell C such that for all $i \in \{0, \dots, d\}$ there is a vertex $x \in C$ such that $c(x) = i$.

3.3.3 Reducing SPERNER to ENDOFLINE

In order for a reduction of TARSKI* onto SPERNER to show membership in **PPAD**, we need to show that SPERNER is **PPAD**-Complete. A sketch of the proof was first given in [28]. We will give a complete proof, as the construction we use will become relevant in the following chapters.

[28]: Papadimitriou (1994), *On the complexity of the parity argument and other inefficient proofs of existence*

Theorem 3.4

SPERNER is **PPAD**-complete.

Proof. To prove that SPERNER is **PPAD**-complete it suffices to show that SPERNER can be reduced to ENDOFLINE and that ENDOFLINE can be reduced to SPERNER, because by definition ENDOFLINE is **PPAD**-Complete.

We start by showing that ENDOFLINE reduces onto SPERNER. Take an ENDOFLINE instance $P, S : [N] \rightarrow [N]$, such that 0 is the designated source. Now we construct a two dimensional SPERNER instance on a lattice $L = N^2$ as follows

□

Next, we will use this problem to show that TARSKI* is a total search problem and hence lies in **PPAD**.




Figure 3.4: Reduction of two dimensional SPERNER to ENDOFLINE. The circuit S is given by the full arrows and S' by the dashed arrows. The solutions are colored green. The designated source is colored blue.

3.4 Reducing TARSKI* to SPERNER

For us to be able to use SPERNER's Lemma on our TARSKI* instances, we need to define a coloring of the vertices of L . We propose the following coloring $c : L \rightarrow \{0, \dots, d\}$:

$$c(x) = \begin{cases} 0 & \text{if } x \leq f(x) \\ 1 & \text{else if } x[1] > f(x)[1] \\ \vdots & \\ d & \text{else if } x[d] > f(x)[d] \end{cases}$$

A vertex colored 0 indicates that the function points *weakly forwards* in all dimensions, a vertex colored i for $i \geq 1$ indicates that the function points *backwards* in at least the i -th dimension.

We give an example of the coloring of a Tarski instance in Figure 3.5. We now need two results. First, we need to show that a cell with all colors always exists, allowing us to show that TARSKI* is a total search problem. Second, we need to show that finding a cell with all colors yields a solution to TARSKI* in polynomial time.




Figure 3.5: Coloring of a TARSKI* instance on a 2-dimensional lattice. The vertices colored 0 indicate that the function points weakly forwards in all dimensions, the vertices colored 1 indicate that the function points backwards in the first dimension, and the vertices colored 2 indicate that the function points backwards in the second dimension and not in the first.

Claim 3.5

For any TARSKI* instance with vertices colored as above, there is always a cell with all colors.

Proof. This claim follows directly from SPERNER'S Lemma and the coloring we have defined. There can never be a vertex colored i with $x[i] = 0$ because this would imply that $f(x)[i] < x[i]$, which is a contradiction to the construction of the function. Hence, by dividing each cell of the lattice into simplices, we can apply SPERNER'S Lemma to show that a cell with all colors always exists. The vertices we use as the vertices of the large simplex are $\{(0, \dots, 0), (2^n - 1, 0, \dots, 0), \dots, (0, \dots, 2^n - 1)\}$. \square

Claim 3.6

Finding a cell with all colors yields a solution to TARSKI*, in $\mathcal{O}(d)$ additional steps.

Proof. Assume we have found a simplex, with vertices colored $0, \dots, d$. Let us denote x_i the vertex colored i , for $i \in \{0, \dots, d\}$. Notice that all of these vertices are by construction contained in some cell (hypercube of length 1); let $\mathbf{0}$ be the smallest vertex of this hypercube and $\mathbf{1}$ the largest. In particular, this means

that for all i , we have:

$$\mathbf{0} \leq x_i \leq \mathbf{1} \quad \text{and} \quad f(x_i)[i] < x_i[i] \quad \text{for } i > 0$$

We now proceed by case distinction:

Case 1: If x_0 is a fixed point, then $x = y = x_0$ is a solution to TARSKI*.

Case 2: If $x_0 \neq f(x_0)$ and $x_0 = \mathbf{0}$. Then there is an i such that $f(x_0)[i] > x_0[i]$, which means that $f(x_0)[i] - x_0[i] \geq 1$. At the same time we must have $f(x_i)[i] < x_i[i]$ and $x_0[i] - x_i[i] \leq 0$ because $x_0 = \mathbf{0}$, and hence $x_i[i] - f(x_i)[i] \geq 1$. Now we get:

$$\begin{aligned} f(x_0)[i] - f(x_i)[i] &= \underbrace{f(x_0)[i] - x_0[i]}_{\geq 1} + \underbrace{x_0[i] - x_i[i]}_{\geq 0} + \underbrace{x_i[i] - f(x_i)[i]}_{\geq 1} \\ f(x_0)[i] - f(x_i)[i] &\geq 2 \end{aligned}$$

This implies that $f(x_0) \not\leq f(x_i)$, and hence x_0, x_i are two points witnessing a violation of monotonicity of f , which form a solution to TARSKI*.

Case 3: If $x_0 \neq f(x_0)$ and $x_0 \neq \mathbf{0}$. We claim that either $f(\mathbf{0}) \leq \mathbf{0}$, or we have a violation of monotonicity. Assume for the sake of contradiction that there is an i such that $f(\mathbf{0})[i] > \mathbf{0}[i]$. Then we must have $f(x_i)[i] < x_i[i]$ hence we get: $f(\mathbf{0})[i] \not\leq f(x_i)[i]$, which is a violation of monotonicity. This means that either we can return $y = x_0$ and $x = \mathbf{0}$ as a solution to TARSKI*, or x_i and $\mathbf{0}$ as a violation of monotonicity.

This shows we can solve a TARSKI* instance in $\mathcal{O}(d)$ additional steps. \square

This shows that TARSKI* is a total search problem and can be reduced to SPERNER. Hence, TARSKI* lies in PPAD, and by using that $\mathbf{P}^{\text{PPAD}} = \text{PPAD}$, we have shown that TARSKI lies in PPAD, without relying on BROUWER.

In the previous chapter, we demonstrated how one can prove the membership of TARSKI in **PPAD** through a reduction to SPERNER. We now demonstrate that the same approach yields a reduction to ENDOFPOTENTIALLINE, which lies within **EOPL**. This will necessitate a more meticulous examination of the structure of a TARSKI instance and the induced colouring of the lattice points. In order to achieve our objective, we must first construct a specific simplicial decomposition of the lattice. We do this with the intention of obtaining certain useful properties. Ultimately, our goal is to demonstrate that for a monotone TARSKI instance, the associated ENDOFLINE instance does not contain any cycles. This will prove sufficient to establish a reduction to ENDOFPOTENTIALLINE.

4.1 Warmup: No cycles in two-dimensional TARSKI*-instances

The aim of this chapter is to prove that there are no cycles in the ENDOFLINE instances that result from the reduction of monotone TARSKI*-instances onto ENDOFLINE we presented in Subsection 3.3.3. We will do this by studying the SPERNER instances, and showing that the sequence of simplices that form the lines of the ENDOFLINE-instances, cannot form a cycle if the underlying function f is monotone.

Proving this will be a somewhat involved in the general d -dimensional case. To motivate the definitions, and the ideas we will use in the following sections, we want to start by presenting a proof for the much more simple two-dimensional case.

Proposition 4.1

The ENDOFLINE-instance that results from reducing the two-dimensional SPERNER-instance generated by a two-dimensional monotone function $f : L \rightarrow L$, has no cycles.

Proof. The idea of the proof will be to show that any sequence of simplices that are generated by walking through edges colored $0 - 1$, can only cross an axis parallel to the x -axis, once.

Let $L_{d_1=K}$ for $K \in [N]$ be such an axis given by:

$$L_{d_1=K} = \{x \in L : x[1] = K\}$$

4.1	No cycles in two dimensions	33
4.2	Freudenthal's Simplicial Decomposition	35
4.3	Sequences of simplices	39
4.4	Super-unique TARSKI instances	40
4.5	Orientating the simplices	40
4.5.1	Orienting a simplex	41
4.5.2	Orienting a simplicial complex	43
4.5.3	Colored orientations	46
4.6	Properties of colored of oriented simplicial sequences	47
4.6.1	General properties of the coloring	47
4.6.2	Properties of sequences of simplices	47
4.7	No cycles in the ENDOFLINE instance	50
4.8	Discussing the reduction to EOPL	51

Now let us assume that we have a path of simplices that crosses $L_{d_1=K}$ twice. Then there must be two edges colored $0-1$ on $L_{d_1=K}$. Now let us call these two edges (x_1, x_2) and (y_1, y_2) , such that $x_1 < x_2$ and $y_1 < y_2$. We want to discuss how these edges can be colored. If $c(x_1) = 0$ and $c(x_2) = 1$, we get a contradiction because this leads to $f(x_1)[1] > f(x_2)[1]$. Hence this cannot occur. A similar argument for y_1, y_2 yields that $c(x_1) = c(y_1) = 1$ and $c(x_2) = c(y_2) = 0$. For convenience a sketch of the situation is given in Figure 4.1.

Now when walking along edges colored $0-1$, we notice that the edge we traverse should always be oriented in the same way. For instance the vertex on our left should be colored 0 and the vertex on our right colored 1. Without loss of generality assume that we crossed the edge $x_1 - x_2$ first, then the vertices $y_1 - y_2$ should be colored opposite, to allow us to traverse $L_{d_1=K}$ in the reverse direction. But this is not the case which means that we traverse $L_{d_1=K}$ at most once. This is true for any $K \in [N]$ which guarantees that a cycle cannot exist. \square




Figure 4.1: Sketch of the setting for the two dimensional proof

Let us briefly discuss what the key ingredients for this proof were and what the steps are that we will take to generalize them to the d -dimensional case.

On a high level we want to show that one cannot cross a given hyperplane of the lattice more than once when going through rainbow-faces colored $\{0, \dots, d-1\}$. This turns out not to be true but we will be able to show a weaker result, which still prohibits cycles. In two dimensions we did this by showing that when we traverse such a hyperplane we had restrictions on how the edges could be oriented. We will generalize this notion of *orientation* for the colored faces we are working with. We will then achieve a contradiction by showing that these faces must be oriented the same way if they are part of a path of simplices that form a line in the ENDOFLINE reduction and obtain a contradiction from this.

In order to this we will need to discuss in detail how we subdivide the cells of the lattice into simplices, we will do this by introducing *Freudenthal's simplicial decomposition*. Once we have the simplices, we will discuss *sequences of simplices* which are the objects that will turn into the lines of the ENDOFLINE instances. Then we will need to define the notion of *orientation* for colored simplices. We will then need to show that the monotonicity of f prohibits certain simplices from existing, and use this to obtain a contradiction to the existence of cycles.

4.2 Choosing a simplicial decomposition of the lattice — Freudenthal's Simplicial Decomposition

In the previous chapter, we left the choice of a specific simplicial decomposition of the lattice open, as it did not contribute to our reduction. In this chapter, we aim to be more precise in our approach by selecting a specific simplicial decomposition that will enable us to derive structural results. We begin by outlining the desired properties of our simplicial decomposition. The most fundamental property is that every simplex of the decomposition must be contained within a single cell of the lattice. This implies that we can limit our inquiry to the identification of a simplicial decomposition of a single d -dimensional hypercube of side-length 1. Additionally, it is important to note that our objective does not entail the introduction of any new vertices; instead, we seek a decomposition of the hypercube that can be expressed as a set of subsets of the hypercube's vertices. Finally, we wish for the vertices of a given simplex to be totally ordered with respect to the partial order defined in Section 2.3. This will allow us to argue that two vertices, inside a given simplex, are always comparable, and thus their images through f must also be comparable, which will be useful.

Such a decomposition exists, and is known in the literature as *Freudenthal's simplicial decomposition* [30]. We will introduce it in a combinatorial way here, and refer the reader to the original paper for a geometric construction of the same decomposition.

[30]: Freudenthal (1942), *Simplizialzerlegungen von Beschränkter Flachheit*

Definition 4.2 — Freudenthal's Simplicial Decomposition.

Consider a unit hypercube $[0, 1]^d$ in \mathbb{R}^d and consider S_d the group of all permutations of the dimensions of the hypercube $\{1, \dots, d\}$. For every permutation $\pi \in S_d$, define the simplex S_π as the convex hull of the vertices:

$$\begin{aligned} v_0 &= (0, 0, \dots, 0) \\ v_1 &= v_0 + e_{\pi(1)} \\ v_2 &= v_1 + e_{\pi(2)} \\ &\vdots \\ v_d &= v_{d-1} + e_{\pi(d)} = (1, 1, \dots, 1) \end{aligned}$$

Here we will use the notation e_i to denote the i -th unit vector in \mathbb{R}^d .

The set of such simplexes $\mathcal{S} = \{S_\pi : \pi \in S_d\}$ is Freudenthal's simplicial decomposition of the hypercube $[0, 1]^d$.

We want to begin by arguing why this decomposition is well-defined. We begin by showing that every point of the hypercube is contained in at least one simplex of \mathcal{S} .

Lemma 4.3

Let $x = (x[1], \dots, x[d]) \in [0, 1]^d$, let $\pi \in S^d$ be the permutation such that $x[\pi(1)] \leq x[\pi(2)] \leq \dots \leq x[\pi(d)]$. Then $x \in S_\pi$.

Proof. We want to show that x is a convex combination of the vertices of S_π . We define the following sequence of real numbers:

$$\begin{aligned}\lambda_0 &= x[\pi(1)] \\ \lambda_1 &= x[\pi(2)] - x[\pi(1)] \\ \lambda_2 &= x[\pi(3)] - x[\pi(2)] \\ &\vdots \\ \lambda_{d-1} &= x[\pi(d)] - x[\pi(d-1)] \\ \lambda_d &= 1 - x[\pi(d)]\end{aligned}$$

Notice that we have $\lambda_i \geq 0$ for all i and $\sum_{i=0}^d \lambda_i = 1$, by telescoping the sum. We can now write x as a convex combination of the vertices of S_π as follows by noticing that $v_i = \sum_{j=0}^i e_{\pi(j)}$:

$$\begin{aligned}\sum_{i=0}^d \lambda_i v_i &= \sum_{i=0}^d \lambda_i \left(\sum_{j=0}^i e_{\pi(j)} \right) = \sum_{i=0}^d \sum_{j=1}^i \lambda_i e_{\pi(j)} \\ &= \sum_{j=1}^d \sum_{i=0}^j \lambda_i e_{\pi(j)} = \sum_{j=1}^d e_{\pi(j)} \sum_{i=0}^j \lambda_i = \sum_{j=1}^d e_{\pi(j)} x[\pi(j)] = x\end{aligned}$$

This shows that x is a convex combination of the vertices of S_π , and thus $x \in S_\pi$. \square

Next we discuss why this really forms a partition of the hypercube. Of course a given point x can be contained in multiple simplexes, but we want to show that this does not happen apart from on the boundary of the simplices.

Lemma 4.4

Let $S_\pi \in \mathcal{S}$ be a simplex. Then the *interior* of S_π is:

$$\text{int}(S_\pi) = \{x \in [0, 1]^d : 0 < x[\pi(1)] < x[\pi(2)] < \dots < x[\pi(d)] < 1\}$$

Proof. The same proof as for lemma 4.3, holds with the added constraint that all $\lambda_i > 0$, this then shows that these points are in the interior of the simplex. \square

These two Lemmata together show that we have a well-defined simplicial decomposition of the hypercube. We can now use this decomposition to prove some structural results about the lattice points of a TARSKI instance. We start by showing that this simplicial decomposition has the desired properties.

Lemma 4.5

Let $S_\pi \in \mathcal{S}$ be a simplex. Then the vertices of S_π are totally ordered with respect to the partial order defined in Section 2.3. In particular we claim that:

$$v_0 < v_1 < v_2 < \dots < v_d$$

Proof. Because this relation is transitive it suffice to show that $v_i < v_{i+1}$ for all $i \in \{0, \dots, d-1\}$. This follows immediately from the construction of the v_i as we have $v_i[j] = v_{i+1}[j]$ for all $j \neq \pi(i+1)$ and $v_i[\pi(i+1)] = v_{i+1}[\pi(i+1)] - 1$. \square

This directly implies the following corollary.

Corollary 4.6

For two vertices x, y of any simplex $S \in \mathcal{S}$, if for any $i \in \{1, \dots, d\}$ we have $x[i] < y[i]$, then $x < y$. In particular $x \not\leq y$ is equivalent to $x > y$.

Notice that this is not the case for any two points in the hypercube, as the partial order is not a total order. This is why choosing a simplicial decomposition with this property will be crucial in the following sections. Next we want to introduce a new notation which will allow us describe these simplices more succinctly. Assume that a permutation π of the dimensions, induces a simplex S_π , with vertices v_0, \dots, v_d , as defined in Definition 4.2. Then we will denote the d -dimensional simplex S_π as:

$$v_0 \xrightarrow{\pi(1)} v_1 \xrightarrow{\pi(2)} v_2 \xrightarrow{\pi(3)} \dots \xrightarrow{\pi(d)} v_d$$

This notation means that we obtain v_i by moving by one unit-length in the direction $\pi(i)$ from v_{i-1} . We already briefly discussed how the faces of a given simplex are given. We will also describe how to describe these faces in our notation. We will denote the face of S_π obtained by removing the vertex v_i as:

$$v_0 \xrightarrow{\pi(1)} v_1 \xrightarrow{\pi(2)} \dots v_{i-1} \xrightarrow{\pi(i), \pi(i+1)} v_{i+1} \xrightarrow{\pi(i+2)} \dots \xrightarrow{\pi(d)} v_d$$

We can remark the following about the faces of a simplex.

Remark 4.7

For a given $d - 1$ dimensional simplex F in \mathcal{S} we have that:

(1) If F is of the form:

$$F : v_0 \xrightarrow{\pi(1)} \cdots v_{i-1} \xrightarrow{\pi(i), \pi(i+1)} v_{i+1} \xrightarrow{\pi(i+2)} \cdots \xrightarrow{\pi(d)} v_d$$

Then F is a face of exactly two simplices S_1 and S_2 :

$$S_1 : v_0 \xrightarrow{\pi(1)} \cdots v_{i-1} \xrightarrow{\pi(i+1)} w_i \xrightarrow{\pi(i)} v_i \xrightarrow{\pi(i+2)} \cdots \xrightarrow{\pi(d)} v_d$$

$$S_2 : v_0 \xrightarrow{\pi(1)} \cdots v_{i-1} \xrightarrow{\pi(i)} w'_i \xrightarrow{\pi(i+1)} v_i \xrightarrow{\pi(i+2)} \cdots \xrightarrow{\pi(d)} v_d$$

(2) If F is of the form:

$$F : v_0 \xrightarrow{\pi(1)} v_1 \xrightarrow{\pi(2)} \cdots \xrightarrow{\pi(d-1)} v_{d-1}$$

Then F is a face of exactly two simplices S_1 and S_2 :

$$S_1 : v_0 \xrightarrow{\pi(1)} v_1 \xrightarrow{\pi(2)} \cdots \xrightarrow{\pi(d-1)} v_{d-1} \xrightarrow{\pi(d)} w_d$$

$$S_2 : w_0 \xrightarrow{\pi(d)} v_0 \xrightarrow{\pi(1)} v_1 \xrightarrow{\pi(2)} \cdots \xrightarrow{\pi(d-1)} v_{d-1}$$

Notice that the case (1) is the case where the face is inside the cell, and the case (2) is the case where the face is on the border of the cell.

We discuss what simplices of the decomposition neighbour each other. We claim that a given simplex has $d - 1$ neighboring simplices inside a given cell, and two neighboring simplices in neighboring cells. More precisely we have the following lemma.

Lemma 4.8 — Neighboring Simplices.

Let $S_\pi \in \mathcal{S}$ be a simplex:

$$v_0 \xrightarrow{\pi(1)} v_1 \xrightarrow{\pi(2)} v_2 \xrightarrow{\pi(3)} \cdots \xrightarrow{\pi(d)} v_d$$

Then the following simplices are neighbors of S_π :

- $v_0 \xrightarrow{\pi(2)} v_1 \xrightarrow{\pi(1)} \cdots v_{i-1} \xrightarrow{\pi(i+1)} w_i \xrightarrow{\pi(i+1)} v_i \xrightarrow{\pi(i+2)} \cdots \xrightarrow{\pi(d)} v_d$, for all $i \in \{1, \dots, d-1\}$, where w_i is the vertex obtained by moving one unit in the direction $\pi(i+1)$ from v_{i-1} .
- $w_d \xrightarrow{\pi(d)} v_0 \xrightarrow{\pi(1)} v_1 \xrightarrow{\pi(2)} \cdots \xrightarrow{\pi(d-1)} v_{d-1}$, where w_d is the vertex obtained by moving one unit in the direction $-\pi(d)$ from v_0 .

► $v_2 \xrightarrow{\pi(2)} \cdots \xrightarrow{\pi(d-1)} v_{d-1} \xrightarrow{\pi(d)} v_d \xrightarrow{\pi(1)} w_1$, where w_1 is the vertex obtained by moving one unit in the direction $\pi(1)$ from v_d .

Proof. The proof follows directly by enumerating the faces of S_π and using remark 4.7. \square

4.3 Sequences of simplices

In this section we introduce and study *sequences of simplices*. They will be useful as we will argue that paths in the ENDOFLINE instance are sequences of simplices, and latter that the colored orientation of the simplicial decomposition will prevent these paths from forming cycles. We start by defining what we mean by a sequence of simplices.

Definition 4.9 — Sequence of simplices.

A *sequence of simplices*, or *simplicial sequence* is a sequence $(S_i)_{i=1}^k$ of d -dimensional simplices $S_i \in \mathcal{S}$ such that:

- (1) $S_{i+1} \not\subset \{S_1, \dots, S_i\}$ for all $i \in \{1, \dots, k-1\}$.
- (2) S_i and S_{i+1} share a $d-1$ -dimensional face F_i for all $i \in \{1, \dots, k-1\}$.

We want to also introduce formally the notion of *transition faces*, which will represent the face we are *walking through*.

Definition 4.10 — Transition faces.

Let $(S_i)_{i=1}^k$ be a simplicial sequence. We call the sequence $(F_i)_{i=1}^{k-1}$, of $d-1$ -dimensional faces given by $F_i = S_i \cap S_{i+1}$, the *transition sequence*. We call the individual F_i *transition faces*.

Now of course we need to discuss how we can add colors. In the following $C \subset \{0, \dots, d\}$, will be a subset of colors. We will use this subset of colors to define restrictions on how sequences of simplices we study should be colored.

Definition 4.11 — Rainbow face.

Let $C \subset \{0, \dots, d\}$ be a subset of colors, and F a face of a simplicial complex \mathcal{S} . We say that F is a *rainbow face* for colors C , if all vertices of F are colored with colors from C and all colors in C appear in F .

Let us now restrict these general definitions to the context that we will be studying we will fix $C = \{0, \dots, d-1\}$. We will

only be studying sequences where the F_i 's are rainbow faces i.e. faces colored with exactly the colors $C = \{0, \dots, d-1\}$. We will call a sequence of simplices with this property a *rainbow simplicial sequence*. We will only be studying valid sequences of simplices that have this property, as these are the sequences of simplices that lines in the ENDOFLINE instances. One property is still missing: the sequences of simplices that reduce to the lines in the ENDOFLINE instances are *maximal* in the sense that they can not be prolonged. This motivates the following definition.

Definition 4.12 — Maximal sequence.

A *maximal sequence of simplices for colors C* is a valid sequence $(S_i)_{i=1}^k$ of simplices $S_i \in \mathcal{S}$ for colors C such that:

- (1) There is no simplex $S_{k+1} \in \mathcal{S}$ such that $(S_i)_{i=1}^{k+1}$ is a valid sequence.
- (2) There is no simplex $S_0 \in \mathcal{S}$ such that $(S_i)_{i=0}^k$ is a valid sequence.

Intuitively we say that a sequence is maximal if we cannot make it longer by adding simplices at the beginning or end.

A special type of line that we can obtain in the ENDOFLINE instances are cycles. These are the result of reducing a special class of simplicial sequences, we will also call *cycles*.

Definition 4.13 — Cycle.

A *cycle of simplices for colors C* is a valid maximal sequence $(S_i)_{i=1}^k$ of simplices $S_i \in \mathcal{S}$ for colors C such that $S_1 \cap S_k$ is a $d-1$ dimensional rainbow face for colors C .

Notice that maximal rainbow simplicial sequences are exactly the sequences of simplices which yield the lines in the ENDOFLINE instance which we reduce SPERNER instances to. We can now discuss how we orient these simplices.

4.4 A side note on super-unique TARSKI instances

TODO.

4.5 Orientation of a the simplicial decomposition

In this section we discuss how to orient the simplicial decomposition of the lattice, we defined in the previous section. This will be important as we will argue in the next section, that the existence of a cycle would contradict the orientation of the

simplicial decomposition. We start by defining what we mean by an orientation of a simplex and then discuss how to extend this to a general simplicial complex.

4.5.1 Orientation of a simplex

Definition 4.14 — Orientation of a simplex.

An *orientation* of a simplex S spanned by the vertices v_0, \dots, v_d is a choice of a permutation of the vertices $[v_{\pi(0)}, \dots, v_{\pi(d)}]$.

Notice that this leaves us with $d!$ possible orientations of a simplex. Our notion of orientability should only lead to two possible classes of orientations, as an orientation of a 1-simplex is simply a choice of direction, and an orientation of a 2-simplex is a choice of a cyclic order of the vertices. Hence we want to define when two orientations are equivalent.

Definition 4.15 — Equivalent orientations.

Two orientations π and σ of a simplex S are *equivalent* if they differ by an even permutation. That is if $\sigma = \pi \circ \tau$ for some permutation τ with an even number of inversions.

In particular we give a more explicit definition of the equivalence of orientations of a 2-simplex, by relying on a total order \preceq of the vertices. We then get the following useful lemma:

Lemma 4.16

Two orientations σ, τ of a simplex S are equivalent if and only if $\text{sgn}(\sigma) = \text{sgn}(\tau)$, with respect to the total order \preceq .

For a lattice this can be achieved by defining \preceq to be the lexicographic order of the vertices.

We would like to define the *opposite orientation* of a simplex, which should be an orientation which has the opposite sign with respect to the total order \preceq . This can be achieved by setting:

$$-[v_0, v_1, v_2, \dots, v_d] = [v_1, v_0, v_2, \dots, v_d]$$

We then have that the opposite orientation is not equivalent to the original orientation. This way we have a representative of both equivalence classes.

This means that we now have two equivalence classes of orientations for any simplex. We want to discuss how an orientation of a simplex extends to the faces of this simplex next. Notice that the faces of a simplex are themselves simplices, and thus have an orientation. Let $[v_0, \dots, v_d]$ be an orientation of a simplex S . Now notice that every face can be obtained by removing one of the vertices v_j of S . Hence for every face F ,

the permutation $[v_0, \dots, \hat{v}_j, \dots, v_d]$ is an orientation of F . But the orientation $-[v_0, \dots, \hat{v}_j, \dots, v_d]$ is also a valid orientation of F . For reasons which will become apparent later we define the induced orientation of a face as follows:

We use the notation \hat{v}_j to denote that v_j is missing.

Definition 4.17 — Induced orientation of a face.

Let $\sigma = [v_0, \dots, v_d]$ be an orientation of a simplex S . The *induced orientation* of a face F of S , which is obtained by removing the vertex v_j from the vertex, is the orientation:

$$\sigma_j = (-1)^j \cdot [v_0, \dots, \hat{v}_j, \dots, v_d]$$

We claim that the induced orientations of faces, yields a consistent orientation of the simplex, that is that for every $d-2$ -simplex E which is a face of two $d-1$ -simplices S_1 and S_2 , the induced orientations of E in S_1 and S_2 are opposite.

Claim 4.18

Let F_1 and F_2 be two $d-1$ -simplices in S which share a common face E . Then the induced orientations of E in S_1 and S_2 are opposite.

Proof. Let $[v_0, \dots, v_d]$ be an orientation of S . The face E is obtained by removing two vertices v_i, v_j from S . Without loss of generality assume that F_1 is obtained by removing v_i from S and F_2 is obtained by removing v_j from S . Then the induced orientations S_1 and S_2 are:

$$\begin{aligned} S_1 : & \quad (-1)^i \cdot [v_0, \dots, \hat{v}_i, \dots, v_d] \\ S_2 : & \quad (-1)^j \cdot [v_0, \dots, \hat{v}_j, \dots, v_d] \end{aligned}$$

Now without loss of generality assume that $i < j$, then we have that the induced orientations of E in S_1 and S_2 are:

$$\begin{aligned} E \text{ in } S_1 : & \quad (-1)^i \cdot (-1)^{j-1} \cdot [v_0, \dots, \hat{v}_i, \dots, \hat{v}_j, \dots, v_{d-1}] \\ E \text{ in } S_2 : & \quad (-1)^j \cdot (-1)^i \cdot [v_0, \dots, \hat{v}_i, \dots, \hat{v}_j, \dots, v_{d-1}] \end{aligned}$$

This shows that the induced orientations of E in S_1 and S_2 are opposite. \square

We give an example of the orientation of a 3-simplex and its faces in Figure 4.2. We can now discuss how we can extend this notion to a general simplicial complex.




Figure 4.2: Example of the orientation of a 3-simplex on the left, and the induced orientation of the faces on the right.

4.5.2 Orientation of a simplicial complex

A simplicial complex can be thought of as a collection of simplices which are glued together on their face. Our goal is now to extend this notion of orientation to these simplicial complexes. Formally we define a simplicial complex as follows [31]:

Definition 4.19 — Simplicial complex.

A simplicial complex \mathcal{K} in \mathbb{R}^d is a collection of simplices such that:

- (1) Every face of a simplex in \mathcal{K} is also in \mathcal{K} .
- (2) The intersection of any two simplices in \mathcal{K} is a face of both simplices.

The lattice points which we are interested in, together with Freudenthal's simplicial decomposition of each cell, form a simplicial complex. We now want to define an orientation of a simplicial complex. Of course such an orientation relies on an orientation of each simplex, and we want to make sure that these orientations are in some sense “compatible” on the faces of the simplicial complex. We will define this notion in the following definition.

Definition 4.20 — Orientation of a simplicial complex.

An *orientation* of a simplicial complex \mathcal{K} is a choice of an orientation of every d -simplex in \mathcal{K} , such that for every intersection of two simplices $S_1, S_2 \in \mathcal{K}$, the induced orientation of the face $F = S_1 \cap S_2$ in S_1 and S_2 are opposite.

If such an orientation exists, we say that the simplicial complex is *orientable*.

[31]: Munkres (2018), *Elements of algebraic topology*

We now claim that the simplicial complex formed by the lattice points and Freudenthal's simplicial decomposition is orientable. This will be crucial in the next section, where we will argue that the existence of a cycle in the ENDOFLINE instance would contradict the orientation of the simplicial complex. In particular this shows that a Mobius Strip or the higher dimensional equivalents do not exist in our simplicial complex.

Claim 4.21

The simplicial complex formed by the lattice points and Freudenthal's simplicial decomposition is orientable.




Figure 4.3: Example of the orientation of a Freudenthal's simplicial complex in 2 dimensions.

Proof. We will give an orientation of every d simplex, and then show that the induced orientation of the faces of the simplicial complex are opposite. Let $\pi \in S^d$ be a permutation of the dimensions, and $v_0 \in L$ a vertex of the lattice. We then obtain a simplex $S_\pi \in \mathcal{S}$ as previously described:

$$v_0 \xrightarrow{\pi(1)} v_1 \xrightarrow{\pi(2)} v_2 \xrightarrow{\pi(3)} \dots \xrightarrow{\pi(d)} v_d$$

We now orient S_π using the permutation:

$$\sigma = \text{sgn}(\pi) \cdot [v_0, \dots, v_d]$$

First we notice that for all $d - 2$ simplices, two neighboring $d - 1$ -simplices are contained in exactly one d simplex of the decomposition, and hence the orientation is consistent, as discussed in claim 4.18.

Now let us look at a common face F of two d -simplices S_1 and S_2 . We proceed by case distinction:

Case 1: Assume that S_1 and S_2 are in the same cell, then F is of the form:

$$F : v_0 \xrightarrow{\pi(1)} \cdots v_{i-1} \xrightarrow{\pi(i), \pi(i+1)} v_{i+1} \xrightarrow{\pi(i+1)} v_i \xrightarrow{\pi(i+2)} \cdots \xrightarrow{\pi(d)} v_d$$

And we have that S_1 and S_2 are of the form:

$$\begin{aligned} S_1 : v_0 &\xrightarrow{\pi(1)} \cdots v_{i-1} \xrightarrow{\pi(i+1)} w_i \xrightarrow{\pi(i)} v_i \xrightarrow{\pi(i+2)} \cdots \xrightarrow{\pi(d)} v_d \\ S_2 : v_0 &\xrightarrow{\pi(1)} \cdots v_{i-1} \xrightarrow{\pi(i)} w'_i \xrightarrow{\pi(i+1)} v_i \xrightarrow{\pi(i+2)} \cdots \xrightarrow{\pi(d)} v_d \end{aligned}$$

We immediately notice that $\text{sgn}(S_1) = -\text{sgn}(S_2)$. We remove a vertex w_i, w'_i of the same rank in S_1 and S_2 in order to obtain F . Hence the induced orientation of F in S_1 and S_2 are opposite.

By abuse of notation we will denote by $\text{sgn}(S_1)$ the sign of the permutation inducing S_1 .

Case 2: Next assume that S_1 and S_2 are in neighboring cells, then as dicussed in Remark 4.7 F is of the form:

$$F : v_0 \xrightarrow{\pi(1)} v_1 \xrightarrow{\pi(2)} \cdots \xrightarrow{\pi(d-1)} v_{d-1}$$

And we have that S_1 and S_2 are of the form:

$$\begin{aligned} S_1 : v_0 &\xrightarrow{\pi(1)} v_1 \xrightarrow{\pi(2)} \cdots \xrightarrow{\pi(d-1)} v_{d-1} \xrightarrow{\pi(d)} w_d \\ S_2 : w_0 &\xrightarrow{\pi(d)} v_0 \xrightarrow{\pi(1)} v_1 \xrightarrow{\pi(2)} \cdots \xrightarrow{\pi(d-1)} v_{d-1} \end{aligned}$$

We once again must proceed by case distinction.

Case 2.1: If d is even, then: $\text{sgn}(S_1) = -\text{sgn}(S_2)$, and we remove a vertex of rank d in S_1 and of rank 0 in S_2 to obtain F . We have $(-1)^d = (-1)^0 = 1$ and hence the induced orientation of F in S_1 and S_2 are opposite.

Case 2.2: If d is odd, then $\text{sgn}(S_1) = \text{sgn}(S_2)$, and we remove a vertex of rank d in S_1 and of rank 0 in S_2 to obtain F . We have $(-1)^d = -1$ and $(-1)^0 = 1$ and hence the induced orientation of F in S_1 and S_2 are opposite.

This shows that the simplicial complex formed by the lattice points and Freudenthal's simplicial decomposition is orientable. \square

We give an example of such an orientation in Figure 4.3. Now that we have shown that the Freudenthal decomposition is orientable, we want to talk about how it can be extended to colored simplices in the next section.

4.5.3 Orienting colored simplices

In the previous sections we discussed how one could orient simplices. We now want to discuss the orientation of colored faces. We are only interested in rainbow-faces, i.e. $d - 1$ -dimensional faces, colored with all colors $\{0, \dots, d - 1\}$, as these are the faces that we traverse when reducing SPERNER to ENDOFLINE. We should obtain two orientations for every face, depending on what direction we traverse it in.

Definition 4.22 — Orientation of colored faces.

Let F be a face of a simplex S . Let the vertices of F be labelled v_1, \dots, v_d , such that $v_1 < \dots < v_d$. Then there is a permutation $\gamma \in S_d$ such that:

$$c(v_{\gamma(1)}) < c(v_{\gamma(2)}) < \dots < c(v_{\gamma(d)})$$

Then with σ_F the induced orientation of F in S , the orientation of the colored face F is:

$$\text{orient}_S(F) = \text{sgn}(\gamma) \cdot \text{sgn}(\sigma_F)$$

Now we should check that our definition is sound. We want that the orientation of a face F is opposite when traversing the face from opposite sides. This is the content of the following lemma.

Lemma 4.23 — Well-definedness of colored orientation.

Let S_1 and S_2 be two d -dimensional simplices which share a $d - 1$ -dimensional rainbow face F . Then:

$$\text{orient}_{S_1}(F) = -\text{orient}_{S_2}(F).$$

Proof. Because the simplicial complex we work with is orientable, for $\text{indorient}_{S_1}(F)$ the induced orientation of F in S_1 and $\text{indorient}_{S_2}(F)$ the induced orientation of F in S_2 , we have:

$$\text{sgn}(\text{indorient}_{S_1}(F)) = -\text{sgn}(\text{indorient}_{S_2}(F))$$

This immediately implies the desired result. \square

4.6 Properties of colored of oriented simplicial sequences

4.6.1 General properties of the coloring

For this section we assume that we are working on a integer lattice L , and that for a function $f : L \rightarrow L$, the points have been colored $c : L \rightarrow \{0, \dots, d\}$ as in Section 3.4. Now we are ready to present a first observation, which will be a helpful stepping stone for more advanced results.

Lemma 4.24

Assume that f is monotone and that we have $x_i, x_j \in L$, $c(x_i) = i$ and $c(x_j) = j$ for $i, j \in \{1, \dots, d\}$ and $x_i[i] = x_j[i]$, then either:

- (1) $i \geq j$ or
- (2) $i < j$ and $x_i \not\leq x_j$

Proof. Assume that $i < j$ and $x_i \geq x_j$. We must then have $f(x_j)[i] \geq x_j[i] = x_i[i] > f(x_i)[i]$. Now by monotonicity of f we must have $f(x_i) \geq f(x_j)$, which is not possible if $f(x_j)[i] > f(x_i)[i]$. Hence we must have $x_i \not\leq x_j$. This shows that the lemma holds. \square

For vertices of a given simplex we get the following corollary.

Corollary 4.25

Assume that f is monotone and that we have $x_i, x_j \in S$, for some simplex $S \in \mathcal{S}$. Further assume that $c(x_i) = i$ and $c(x_j) = j$ for $i, j \in \{1, \dots, d\}$ with $i < j$ and that $x_i[i] = x_j[i]$, then $x_i < x_j$.

Proof. $x_i \leq x_j$, follows immediately. Because x_i and x_j are colored differently, they can not be equal which shows the strict inequality. \square

Recall that the coloring was given by:

$$c(x) = \begin{cases} 0 & \text{if } x \leq f(x) \\ 1 & \text{else if } x[1] > f(x)[1] \\ \vdots & \\ d & \text{else if } x[d] > f(x)[d] \end{cases}$$

Notice that if we assume that x_i and x_j are in the same simplex of the simplicial decomposition, then the condition $x_i \not\leq x_j$ is equivalent to $x_i < x_j$.

4.6.2 Properties of sequences of simplices

Now we want to work with sequences of simplices, and show that the coloring of the vertices of these simplices have some nice properties. We start by defining what we mean by a sequence of simplices. Let $C = \{0, \dots, d-1\}$ be the subset of colors, and let $(S_i)_{i=1}^k$ be a valid simplicial sequence.

Lemma 4.26

Let S_i, F_i and x_j be as above. For any $i \in \{1, \dots, k-1\}$ there is exactly one $j \in C$ such that we have $x_i^j \neq x_{i+1}^j$.

Proof. F_i and F_{i+1} are two faces of the same d dimensional simplex, and thus they share exactly $d-1$ vertices. This means that there is exactly one vertex x which is in F_i but not in F_{i+1} , and exactly one vertex y which is in F_{i+1} but not in F_i . This means that there is exactly one j such that $x_i^j = x$ and $x_{i+1}^j = y$. \square

Lemma 4.27 — Orientation of transition faces.

Let $(S_i)_{i=1}^k$ be a valid rainbow simplicial sequence. Then $(\text{orient}_{S_i}(F_i))_{i=1}^{k-1}$ is constant.

This means that our definition of orientation of colored faces, makes sense: when we walk through these faces, we always walk through faces which are oriented in the same way.

Proof. It suffices to show that for all $i \in \{1, \dots, k-2\}$:

$$\text{orient}_{S_i}(F_i) = \text{orient}_{S_{i+1}}(F_{i+1})$$

Fix i , and notice that F_i and F_{i+1} are both faces of S_{i+1} . Let us label the vertices of S_i $v_0 < v_1 < \dots < v_d$. Let us denote k and j such that F_i is obtained by removing v_k from S_{i+1} and F_{i+1} is obtained by removing v_j from S_{i+1} . By using Claim 4.18 and noticing that v_k and v_j have the same color we immediately get:

$$-\text{orient}_{S_{i+1}}(F_i) = \text{orient}_{S_{i+1}}(F_{i+1})$$

Now by using Lemma 4.23 we get:

$$\text{orient}_{S_i}(F_i) = -\text{orient}_{S_{i+1}}(F_i) = \text{orient}_{S_{i+1}}(F_{i+1})$$

This is the desired result and concludes the proof. \square

Before we start we want to make an observation on how the dimension d plays together with the orientation of the simplicial complex.

Lemma 4.28

Let $l \in \{1, \dots, d-1\}$ be a dimension. Let S be a d -simplex with colors $C = \{0, \dots, d-1\}$ in the colored simplicial complex, such that S is of the form:

$$(S) : \quad v_0 \xrightarrow{l} v_1 \rightarrow \dots \rightarrow v_d$$

and assume that the face F spanned by v_1, \dots, v_d is a rainbow face. Then we must have for the colors:

$$(F) : \quad c(v_1) \rightarrow \dots \xrightarrow{d} \dots 0 \rightarrow \dots \rightarrow c(v_d)$$

Proof. Every color $c \in \{0, \dots, d-1\}$ appears exactly once in the face F . If the color $c \neq 0$ appear after 0, then by Corollary 4.25 we must have that we move in dimension c between 0 and c :

$$(F) : \quad c(v_1) \rightarrow \dots 0 \rightarrow \dots \xrightarrow{c} \dots \rightarrow c$$

Because we have this for every color $c_i \neq 0$, which appears after 0 in F we must have:

$$(F) : \quad c(v_1) \rightarrow \dots 0 \xrightarrow{c_1} c_1 \xrightarrow{c_2} c_2 \dots \xrightarrow{c_k} c_k$$

Now it is clear that because no vertex is colored with d , we must have that the change in dimension d occurs before the vertex colored 0 appears. This shows that we must have:

$$(F) : \quad c(v_1) \rightarrow \dots \xrightarrow{d} \dots 0 \rightarrow \dots \rightarrow c(v_d)$$

This shows the Lemma. \square

Lemma 4.29 — Ordering of the vertices in transition faces.

Let $(S_i)_{i=1}^k$ be a valid rainbow simplicial sequence. For all i such that F_i is a face between cells then assume that we are moving in dimension $l \in \{1, \dots, d-1\}$, that is l is not a dimension of F_i . Then for all colors $c \in \{k+1, \dots, d-1\} \cup \{0\}$, we have that c appears after l in F_i . This means that F_i is of the form:

$$(F_i) : \quad c(v_1) \rightarrow \dots \rightarrow l \rightarrow \dots \rightarrow c \rightarrow \dots \rightarrow c(v_d)$$

A face between cell is a face of type (2) in Remark 4.7.

Proof. Assume for the sake of contradiction that for $c \in \{k+1, \dots, d-1\} \cup \{0\}$ we have that c appears before l in F_i . Then by Corollary 4.25 we must have that we move in dimension l between l and c . But the dimension l is not a dimension of F_i . This is a contradiction and shows the Lemma. \square

4.7 No cycles in the ENDOFLINE instance

Remark 4.30

Assume that we have a valid colored simplicial sequence which crosses a hyperplane obtained by fixing one dimension $H = L_{k=K}$, for a dimension $k \in \{1, \dots, d\}$ and $K \in [N]$, twice. Let F_i and F_j be the transition faces which cross the hyperplane. Then:

$$\text{orient}_{S_i}(F_i) = -\text{orient}_{S_{j+1}}(F_j)$$

This means that when looking at the hyperplane from any side, F_i and F_j have opposite orientations.

Proof. This is a reformulation of the more general Lemma 4.27. \square

Now we are ready for the proof of our main result. We will start by proving that there are no cycles in the three dimensional case, and then extend this to the general case.

Theorem 4.31 — No cycles in three-dimensional TARSKI*.

There are no cycles in the ENDOFLINE instance which we reduce three dimensional TARSKI* instances to.

Proof. Assume for the sake of contradiction that we have a cycle in the ENDOFLINE instance. Let $(S_i)_{i=1}^k$ be a cycle of simplices for colors $C = \{0, 1, 2\}$. We will show that this leads to a contradiction.

Consider the faces L_i which are the faces spanned by vertices colored not 0 in S_i , and the faces M_i which are the faces spanned by vertices not colored 1 in S_i . Formally:

$$\begin{aligned} L_i &= \{v \in S_i \mid c(v) \neq 0\} \\ M_i &= \{v \in S_i \mid c(v) \neq 1\} \end{aligned}$$

Notice that these L_i 's and M_i 's are always either 2-dimensional or 1-dimensional. Now for the sake of simplicity assume that we remove all 1-dimensional faces from these sequences, and only consider the 2-dimensional faces. Notice that this is in itself a valid oriented simplicial sequence, and in particular a cycle.

Now if we look at the transition faces $(Q_i)_i$ of $(L_i)_i$, and $(R_i)_i$ of $(M_i)_i$, then we have that by Lemma 4.27 that the orientations of the sequence $(Q_i)_i$ is constant and the orientation of the sequence $(R_i)_i$ is constant. We can now proceed by case distinction

Case 1: If we move in dimension 1 then, let H be a hyperplane obtained by fixing dimension 1 which we cross twice. Then let Q_i and Q_j be the transition faces which cross the hyperplane. Then by the remark above we have that: $\text{orient}_{S_i}(Q_i) = \text{orient}_{S_j}(Q_j)$, which means that for either Q_i or Q_j , we have that the vertex colored 1 is larger than the vertex colored 2. This is not possible because we do not move in dimension 1 between these two vertices. This contradicts monotonicity by Corollary 4.25.

Case 2: If we move in dimensions 2 then, let H be a hyperplane obtained by fixing dimension 2 which we cross twice. Then let M_i and M_j be the transition faces which cross the hyperplane. Then by the remark above we have that: $\text{orient}_{S_i}(M_i) = \text{orient}_{S_j}(M_j)$, which means that for either M_i or M_j , we have that the vertex colored 2 is larger than the vertex colored 0. This is not possible because we do not move in dimension 2 between these two vertices. This contradicts monotonicity by Corollary 4.25

Because we will always move in at least dimension 1 or 2 this concludes the proof. \square

4.8 Discussing the reduction of TARSKI* to ENDOFPOTENTIALLINE

We have now shown that there are no cycles in the ENDOFLINE instance which arise from monotone TARSKI* instances. This does not directly imply a direction onto ENDOFPOTENTIALLINE. Intuitively an ENDOFLINE instance without cycles should be equivalent to an ENDOFPOTENTIALLINE instance. But in ENDOFPOTENTIALLINE the non-existence of a cycle is given by a potential which grows monotonically along the lines.

The challenge in our case is to define such a potential. We briefly want to give some intuition for why this is not easy. A first naive attempt, would be to define the potential as something similar to the sum of the coordinates of a vertex of the simplex. This will not work in our case though, because it can very well happen that the path through the simplicial complex is not monotone with respect to the sum of the coordinates, in particular we can cross a hyperplane multiple times, which would not be possible if the sum of the coordinates was a valid potential.

Recall that this is exactly the same potential as we used to reduce TARSKI onto LOCALOPT.

This means that we need to be more creative in our definition of the potential. The idea is to decide on how we orient our potential locally. We start by giving every cell of the simplicial complex coordinates, these will be the coordinates of the smallest vertex of cell. Now look at the colors of the vertices of the cell.

APPENDIX

Bibliography

- [1] Christos H. Papadimitriou. 'On the complexity of the parity argument and other inefficient proofs of existence'. In: *Journal of Computer and System Sciences* 48.3 (June 1994), pp. 498–532. doi: [10.1016/S0022-0000\(05\)80063-7](#). (Visited on 03/05/2024) (cited on pages 2, 13).
- [2] David S. Johnson, Christos H. Papadimitriou, and Mihalis Yannakakis. 'How easy is local search?' In: *Journal of Computer and System Sciences* 37.1 (Aug. 1988), pp. 79–100. doi: [10.1016/0022-0000\(88\)90046-3](#). (Visited on 03/06/2024) (cited on pages 2, 12).
- [3] John Fearnley et al. *End of Potential Line*. Apr. 18, 2018. URL: <http://arxiv.org/abs/1804.03450> (visited on 03/02/2024) (cited on pages 2, 14).
- [4] Alfred Tarski. 'A lattice-theoretical fixpoint theorem and its applications.' In: *Pacific Journal of Mathematics* 5.2 (Jan. 1, 1955), pp. 285–309 (cited on pages 2, 15).
- [5] Kousha Etessami et al. 'Tarski's Theorem, Supermodular Games, and the Complexity of Equilibria'. In: (2020). In collab. with Thomas Vidick. doi: [10.4230/LIPICS.ITCS.2020.18](#). (Visited on 02/24/2024) (cited on pages 2, 3, 15, 17, 19, 20, 23, 24).
- [6] Donald M. Topkis. 'Equilibrium Points in Nonzero-Sum n -Person Submodular Games'. In: *SIAM Journal on Control and Optimization* 17.6 (Nov. 1979), pp. 773–787. doi: [10.1137/0317054](#). (Visited on 07/31/2024) (cited on page 3).
- [7] Paul Milgrom and John Roberts. 'Rationalizability, Learning, and Equilibrium in Games with Strategic Complementarities'. In: *Econometrica* 58.6 (Nov. 1990), p. 1255. doi: [10.2307/2938316](#). (Visited on 07/31/2024) (cited on page 3).
- [8] Anne Condon. 'The complexity of stochastic games'. In: *Information and Computation* 96.2 (Feb. 1992), pp. 203–224. doi: [10.1016/0890-5401\(92\)90048-K](#). (Visited on 07/31/2024) (cited on page 3).
- [9] Bernd Gärtner, Sebastian Haslebacher, and Hung P. Hoang. 'A subexponential algorithm for ARRIVAL'. In: (2021). In collab. with Bansal, Nikhil, Merelli, Emanuela, and Worrell, James. Artwork Size: 14 p. Medium: application/pdf Publisher: ETH Zurich, 14 p. doi: [10.3929/ETHZ-B-000507056](#). (Visited on 07/30/2024) (cited on page 3).
- [10] Chuangyin Dang, Qi Qi, and Yinyu Ye. *Computations and Complexities of Tarski's Fixed Points and Supermodular Games*. May 19, 2020. URL: <http://arxiv.org/abs/2005.09836> (visited on 07/21/2024) (cited on pages 3, 18).
- [11] Xi Chen and Yuhao Li. 'Improved Upper Bounds for Finding Tarski Fixed Points'. In: *Proceedings of the 23rd ACM Conference on Economics and Computation*. EC '22: The 23rd ACM Conference on Economics and Computation. Boulder CO USA: ACM, July 12, 2022, pp. 1108–1118. doi: [10.1145/3490486.3538297](#). (Visited on 07/30/2024) (cited on pages 3, 19).
- [12] Mika Goos et al. 'Further Collapses in TFNP'. In: (2022) (cited on pages 4, 21).
- [13] Sanjeev Arora and Boaz Barak. *Computational complexity : a modern approach*. Cambridge: Cambridge University Press, 2009 (cited on page 5).

- [14] Nimrod Megiddo and Christos H. Papadimitriou. 'On total functions, existence theorems and computational complexity'. In: *Theoretical Computer Science* 81.2 (Apr. 1991), pp. 317–324. DOI: [10.1016/0304-3975\(91\)90200-L](#). (Visited on 03/05/2024) (cited on pages 6, 12).
- [15] Constantinos Daskalakis, Paul W. Goldberg, and Christos H. Papadimitriou. 'The Complexity of Computing a Nash Equilibrium'. In: *SIAM Journal on Computing* 39.1 (Jan. 2009), pp. 195–259. DOI: [10.1137/070699652](#). (Visited on 03/11/2024) (cited on pages 6, 13).
- [16] Constantinos Daskalakis and Christos Papadimitriou. 'Continuous Local Search'. In: *Proceedings of the Twenty-Second Annual ACM-SIAM Symposium on Discrete Algorithms*. Twenty-Second Annual ACM-SIAM Symposium on Discrete Algorithms. Society for Industrial and Applied Mathematics, Jan. 23, 2011, pp. 790–804. DOI: [10.1137/1.9781611973082.62](#). (Visited on 03/12/2024) (cited on pages 7, 21).
- [17] Alexandros Hollender. 'Structural Results for Total Search Complexity Classes with Applications to Game Theory and Optimisation'. PhD thesis. Oxford University, 2021 (cited on page 7).
- [18] Raymond Greenlaw and H. James Hoover. 'Chapter 9 - Circuit Complexity'. In: *Fundamentals of the Theory of Computation: Principles and Practice*. Ed. by Raymond Greenlaw and H. James Hoover. Oxford: Morgan Kaufmann, Jan. 1, 1998, pp. 241–257. DOI: [10.1016/B978-1-55860-547-3.50013-3](#) (cited on pages 8, 9).
- [19] Christos H. Papadimitriou. *Computational complexity*. Reading (Mass): Addison-Wesley, 1994 (cited on page 12).
- [20] Michael Sipser. 'On relativization and the existence of complete sets'. In: *Automata, Languages and Programming*. Ed. by Mogens Nielsen and Erik Meineche Schmidt. Vol. 140. Series Title: Lecture Notes in Computer Science. Berlin, Heidelberg: Springer Berlin Heidelberg, 1982, pp. 523–531. DOI: [10.1007/BFb0012797](#). (Visited on 07/30/2024) (cited on page 12).
- [21] Paul W. Goldberg and Alexandros Hollender. 'The Hairy Ball problem is PPAD-complete'. In: *Journal of Computer and System Sciences* 122 (Dec. 2021), pp. 34–62. DOI: [10.1016/j.jcss.2021.05.004](#). (Visited on 03/10/2024) (cited on page 13).
- [22] John Fearnley, Dömötör Pálvölgyi, and Rahul Savani. 'A Faster Algorithm for Finding Tarski Fixed Points'. In: *ACM Transactions on Algorithms* 18.3 (July 31, 2022), pp. 1–23. DOI: [10.1145/3524044](#). (Visited on 07/21/2024) (cited on page 19).
- [23] Samuel R. Buss and Alan S. Johnson. 'Propositional proofs and reductions between NP search problems'. In: *Annals of Pure and Applied Logic* 163.9 (Sept. 2012), pp. 1163–1182. DOI: [10.1016/j.apal.2012.01.015](#). (Visited on 02/24/2024) (cited on pages 20, 23).
- [24] John Fearnley et al. 'The Complexity of Gradient Descent: $\text{CLS} = \text{PPAD} \cap \text{PLS}$ '. In: *Journal of the ACM* 70.1 (Feb. 28, 2023), pp. 1–74. DOI: [10.1145/3568163](#). (Visited on 07/21/2024) (cited on page 21).
- [25] Xi Chen, Yuhao Li, and Mihalis Yannakakis. 'Reducing Tarski to Unique Tarski (In the Black-Box Model)'. In: *LIPICs, Volume 264, CCC 2023* 264 (2023). In collab. with Amnon Ta-Shma. Artwork Size: 23 pages, 1766105 bytes ISBN: 9783959772822 Medium: application/pdf Publisher: Schloss Dagstuhl – Leibniz-Zentrum für Informatik, 21:1–21:23. DOI: [10.4230/LIPICs.CCC.2023.21](#). (Visited on 07/30/2024) (cited on page 22).

- [26] L. E. J. Brouwer. 'Über Abbildung von Mannigfaltigkeiten'. In: *Mathematische Annalen* 71.1 (Mar. 1911), pp. 97–115. doi: [10 . 1007 / BF01456931](https://doi.org/10.1007/BF01456931). (Visited on 05/04/2024) (cited on page 23).
- [27] Martin Aigner and Günter M. Ziegler. *Proofs from THE BOOK*. 6th ed. 2018. Berlin, Heidelberg: Springer Berlin Heidelberg : Imprint: Springer, 2018. 1 p. (cited on pages 23, 27).
- [28] Christos H. Papadimitriou. 'On the complexity of the parity argument and other inefficient proofs of existence'. In: *Journal of Computer and System Sciences* 48.3 (June 1994), pp. 498–532. doi: [10 . 1016 / S0022 - 0000 \(05 \) 80063 - 7](https://doi.org/10.1016/S0022-0000(05)80063-7). (Visited on 03/22/2024) (cited on pages 23, 26, 28, 29).
- [29] E. Sperner. 'Neuer beweis für die invarianz der dimensionszahl und des gebietes'. In: *Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg* 6.1 (Dec. 1928), pp. 265–272. doi: [10 . 1007 / BF02940617](https://doi.org/10.1007/BF02940617). (Visited on 04/18/2024) (cited on page 27).
- [30] Hans Freudenthal. 'Simplizialzerlegungen von Beschränkter Flachheit'. In: *The Annals of Mathematics* 43.3 (July 1942), p. 580. doi: [10 . 2307 / 1968813](https://doi.org/10.2307/1968813). (Visited on 03/21/2024) (cited on page 35).
- [31] James Raymond Munkres. *Elements of algebraic topology*. The advanced book program. Boca Raton London New York: CRC Press, 2018. 454 pp. (cited on page 43).

Alphabetical Index

- EOPL, 14
- PPAD, 14
- SPERNER, 29
- TARSKI*, 24
- AND-gates, 10
- Boolean circuit, 9
- Brouwer, 23
- Brouwer's fixed point theorem, 23
- cell, 28
- Computed function of a boolean circuit, 10
- Cycle, 40
- decision problems, 5
- Depth of a boolean circuit, 11
- End of Potential Line, 14
- End-of-Line (EoL), 13
- equivalent, 41
- Equivalent orientations, 41
- Freudenthal's Simplicial Decomposition, 35
- Function **NP (FNP)**, 6
- Gate, 9
- gates, 9
- Induced orientation of a face, 42
- inputs gates, 9
- instance, 5
- integral, 24
- interior, 36
- language, 5
- Localopt, 12
- Many-to-one Reduction, 7
- Maximal sequence, 40
- monotone, 15
- Monotone function, 15
- Neighboring Simplices, 38
- No cycles in three-dimensional TARSKI*, 50
- non-standard source, 14
- NOT-gates, 10
- opposite orientation, 41
- OR-gates, 10
- oracle, 7
- order preserving, 15
- Ordering of the vertices in transition faces, 49
- orientable, 43
- orientation, 41, 43
- Orientation of a simplex, 41
- Orientation of a simplicial complex, 43
- Orientation of colored faces, 46
- Orientation of transition faces, 48
- output gates, 9
- Polynomial Local Search (**PLS**), 13
- polynomially balanced, 6
- Progress point, 18
- promise problems, 7
- PromiseTarski, 21
- Rainbow face, 39
- rainbow simplicial sequence, 40
- Search Problem, 5
- search problem, 5
- search problems, 5
- semantic, 12
- Sequence of simplices, 39
- Simplicial complex, 43
- simplicial complex, 43
- simplicial sequence, 39
- simplicial subdivision, 27
- Simplyfying TARSKI, 16
- Size of a boolean circuit, 11
- solution, 5
- Sperner's Lemma, 27
- SuperUniqueTarski, 22
- Tarski, 15
- Tarski's fixed point Theorem, 15
- Total Function **NP (TFNP)**, 6
- total search problem, 6
- Total search problems, 6
- Transition faces, 39
- transition sequence, 39
- Turing Reduction, 7
- type, 9
- UniqueTarski, 21
- unit vector, 35
- violations, 8
- Well-definedness of colored orientation, 46
- wires, 9