

Statistical Inference for Data Science

Dr. Anja Mühlemann

01. September 2022



Question on Day 2

Day 3

Hypothesis Testing

Today's Topics

- Hypotheses
- p-values
- Error type
- Frequently used tests

Inferential Statistics

Inferential Statistics

With a certain degree of certainty, one would like to draw conclusions from empirical data, even if the data are subject to error or incomplete.

3 main techniques

- **Parameter estimates:** Calculation estimate for unknown parameter of underlying probability distribution
- **Confidence intervals:** Calculation of a region within which unknown parameter should lie with certain degree of certainty
- **Tests:** Tests are intended to prove that a certain effect, e.g. the effect of a vaccine, is indeed present.

Tests

- Method for deciding on the correctness of hypotheses under uncertainty

e.g., new medication is better than the old one

2 Hypotheses:


- **Working hypothesis (H1):** Motivation of the study
e.g., the new medication is **better** than the old one
- **Null hypothesis (H0):** Opposite of H1
e.g. the new medication is **not better** than the old one

Goal: reject the null hypothesis with some degree of certainty


Tests II

- Statistical test rely on a **test statistic**, for which distribution under the test assumptions and H_0 is known.
- We calculate the value of the test statistic for the sample at hand (\hat{T})
- And check whether this value is **probable** for the distribution under H_0 .
- To this end the **p-value** is calculated
- If the **p-value < 1 - desired degree of certainty**, we reject H_0
- Otherwise, we cannot reject H_0 , which does **not** necessarily imply that H_1 holds

Tests III



Typically $\alpha = 5\%$



Unlikely observations under H_0

Errors

- **Type 1:** Wrongly reject the null hypothesis due to a fluctuation (**false positive**) $\alpha = 0.05 \rightarrow$ make sure to only make 5% false positive errors
- **Type 2:** Wrongly keep the null hypothesis by interpreting a real effect as a fluctuation (**false negative**) innocent

		Reality	
		True	False
Measured or Perceived	True	Correct 😊	Type 1 error False Positive
	False	Type 2 error False Negative	Correct 😊

Prison example

Innocent person set free	Innocent person jailed
Guilty person set free	Guilty person jailed

Types of Tests


- **One group:** the mean monthly income is larger than 5000.-
- **Two groups:** the mean income of men is larger than that of women
- **\geq Three groups:** effect of tea on weight loss (green, black, none)


\geq Two groups:


- **Paired:** dependent, repeated measurements on same individual, **e.g.** blood pressure before and after surgery
- **Unpaired:** independent, from separate individuals, **e.g.** blood pressure after medication 1 vs. blood pressure after medication 2

Normality

- Many test assume that the sample comes from a normal distribution
- Thus, we need to check whether this is fulfilled before performing such a test
- Shapiro-Wilk test, Shapiro-Francia test, Q-Q-Plot, ...







Exercise

- 3 Slides to be uploaded to ILIAS today
 - 1 slide: Question that the test tries to answer, assumptions on data, other details
 - 1 slide: example from “real live” (if possible)
 - 1 slide: your conclusion from the Notebook on this test
- Will be presented at tomorrow’s discussion session

Nr	Test
1	One-sample t-test
2	One-sample Wilcoxon SR test
3	Paired t-test
4	Paired Wilcoxon SR test
5	Unpaired t-test
6	Unpaired t-test with Welsch’s correction
7	Mann-Withney U test
8	One-way ANOVA