Chair of Software Modeling and Verification (Computer Science 2) | RWTH AACHEN UNIVERSITY

Bachelor Thesis at the
Software Modeling and Verification Group (MOVES)

Expected Visiting Times in Interval Markov Chains


by
Jesper Nilsson

This written work was submitted to
Lehrstuhl für Informatik 2, RWTH Aachen University.

Submission Date: July 28, 2026

First examiner: Joost-Pieter Katoen
Second examiner: insert text
Thesis advisor: Hannah Mertens

Abstract



Hourglass structure of a good abstract. Start broad (research area and problem), funnel down to the specific contribution, then widen again to state results and implications.

Write a self-contained summary of **roughly 150–250 words** following the hourglass structure above. Cover the following five points in order:

1. **Research area.** Name the field (e.g., strategy synthesis, probabilistic program verification, operational semantics) and state in one sentence what the field studies and why it matters.
2. **Specific problem.** State precisely which problem this thesis addresses and why it is challenging or not yet fully solved by existing approaches.
3. **Approach.** Describe the key idea of your contribution in one or two sentences. Avoid mathematical notation and abbreviations, and use easily understandable language.
4. **Theoretical results.** Name the main property proven (soundness, completeness, an equivalence result, or a complexity bound) and state it informally.

5. **Implementation and evaluation (if applicable).** Name the tool and the benchmark set and give the headline result: how many more examples are solved compared to the baseline and what is the runtime overhead.
6. **Implications.** Explain how your results solved the issue or filled the gap you pointed out beforehand.

Acknowledgements

I want to express my most profound appreciation to my supervisor!

Contents

1 Introduction **1**

2 Preliminaries **3**
2.1 Notation 3
2.2 Matheteil 3

3 Markov Chains **9**
3.1 Expected Visiting Times for DTMCs 14
3.1.1 Relating Expected Visiting Times and Quantitative Reachability 18
3.2 Consideration for limiting the state size 20

4 *Interval Markov Chains* **21**
4.1 Expected Visiting Times in IDTMCs 25
4.1.1 Relating Expected Visiting Times and Quantitative Reachability 25
4.1.2 Relation between the semantics 25
4.2 Calculating Expected Visiting Times of IDTMCs 26
4.2.1 UMC semantic 27
4.2.2 IMDP semantic 34
4.2.3 Special Case: Acyclic IDTMCs 37
4.2.4 Value Iteration 38
4.3 Exkurs: aIMCs 38
4.4 Stationary Distributions in IDTMCs 38

5 Conclusion **45**

A Appendix **47**

Chapter 1

Introduction

The introduction motivates the topic and provides an overview of the thesis. It can be seen as a longer version of the abstract that follows the hourglass structure again, but with emphasis on the top part of the hourglass. However, you can now use citations and explain everything in more detail. It should be accessible to any computer scientist without prior knowledge of the specific area.

The background and motivation part should cover the following points:

1. Name the research area and describe informally what objects are studied (e.g., markov automata, probabilistic programs, POMDPs) and what property is of interest (e.g., reachability, complexity, safety).

2. Explain why the property matters.

3. Give a small, self-contained example that illustrates the problem informally.

4. Explain why answering the question automatically is non-trivial and what makes the problem hard in general.

The open-problem part should contain:

1. A summary of what is already known in the area (prior results, existing automated techniques, tool support).

2. A precise identification of the gap: e.g., no automated procedure exists, existing techniques are unsound or incomplete for a relevant subclass, or no efficient algorithm is known.

3. A clear statement of what this thesis aims to achieve.

Related Work

Discuss the most relevant prior work (either here, or in a seperate chapter depending on the scope of existing work). For each cited work, address the following:

1. What does the work do? Describe the approach in one to three sentences and do not just list references.
2. How does it relate to this thesis? Does it solve a special case, use a different technique, or target a different formalism?
3. What limitation motivates your own contribution?

Contribution

List one item per concrete contribution. Each item should:

1. Identify what is new (a technique, a criterion, a framework, an implementation, or an experimental evaluation).
2. State the property proved, if applicable (soundness, completeness, both, or a complexity bound).
3. Reference the theorem, or section in which the contribution appears.

Structure

End the introduction by outlining the remaining chapters. Give at least one sentence per chapter describing its content. Do not simply write "Chapter X presents the main results."

Example: "The remainder of this thesis is organized as follows. Chapter 2 introduces the formal background required throughout this thesis, including ... Then, **??** presents ... **??** describes the implementation and the experimental evaluation. Finally, Chapter 5 summarizes the results and discusses directions for future work."

Chapter 2

Preliminaries

2.1 Notation

We denote the entry in the i-th row and j-th column of a matrix $\mathcal{T}$, as $(\mathcal{T})_{ij}$. We will use the notation $\underline{n} = \{1, 2, \ldots, n\}$ with $n \in \mathbb{N}$ for sets. For vectors v, we denote the i-th entry as $v(i)$ Cardinality of set S as $|S|$. $\bigcirc$ $\diamond$

$\exists!$ to denote: precisely one.

We use *Iverson Brackets* $[\![B]\!]$ to denote the function that maps a Boolean expression B to 1 or 0 such that

$$[\![B]\!] = \begin{cases} 1 & \text{if B is true,} \\ 0 & \text{otherwise.} \end{cases}$$

2.2 Matheteil

Linear Order Measurable set (used in def. 3.10)

Definition 2.1 (Convergence of Geometric Series)
The geometric series $\sum_{k=0}^{\infty} q^k$ converges for all $q \in \mathbb{R}$ with $|q| < 1$, where

$$\sum_{k=0}^{\infty} q^k = \frac{1}{1-q}\,, \quad \text{for } |q| < 1$$

Definition 2.2 (Gershgorin circle theorem)
Let $\mathcal{T} \in \mathbb{C}^{n \times n}$ be a square complex matrix, then for every eigenvalue λ of $\mathcal{T}$ it is

true that

$$|\lambda - (\mathcal{T})_{ii}| \leq \sum_{j=1 \wedge i \neq j}^{n} |(\mathcal{T})_{ij}| \ ,$$

for any $i \in \underline{n}$ and where $|\cdot|$ denotes the absolute value, i.e. magnitude.
In other words, all eigenvalues of $\mathcal{T}$ lie on a disk in the complex plane centered at $(\mathcal{T})_{ii}$ with radius $\sum_{j=1 \wedge i \neq j}^{n} |(\mathcal{T})_{ij}|$.

Matrix invertible if 1 not eigenvalue

Definition 2.3 (Row-Stochastic Matrix)
A square matrix $\mathcal{T} \in \mathbb{R}_{\geq 0}^{n \times n}$ is row-stochastic, if and only if $\sum_{j=0}^{n} (\mathcal{T})_{ij} = 1$ for all rows $i \in \underline{n}$.

When considering each row as a function $f_i(j) = (\mathcal{T})_{ij}$, these functions are discrete probability distributions.

Definition 2.4 ((Strictly) Row-Substochastic Matrix)
A square matrix $\mathcal{T} \in \mathbb{R}_{\geq 0}^{n \times n}$ is row-substochastic, if and only if $\sum_{j=0}^{n} (\mathcal{T})_{ij} \leq 1$ for all rows $i \in \underline{n}$.
$\mathcal{T}$ is strictly row-substochastic if and only if ther exists at least one row i' with $\sum_{j=0}^{n} (\mathcal{T})_{i'j} < 1$

Theorem 2.5
If a matrix $\mathcal{T}$ is strictly row-substochastic, then it has no eigenvalue $|\lambda| \geq 1$.

Proof 1
Let us assume that $|\lambda| > 1$ is an eigenvalue of $\mathcal{T}$. As $(\mathcal{T})_{ij} \in \mathbb{R}_{\geq 0}$, using def. 2.2

$$|\lambda - (\mathcal{T})_{ii}| = |\lambda| - (\mathcal{T})_{ii} \leq \sum_{j=1 \wedge j \neq i}^{n} |(\mathcal{T})_{ij}|$$

$$\iff 1 < |\lambda| \leq (\mathcal{T})_{ii} + \sum_{j=1 \wedge j \neq i}^{n} (\mathcal{T})_{ij} \ .$$

However, as $\mathcal{T}$ is strictly row-substochastic, we know that $(\mathcal{T})_{ii} + \sum_{j=1 \wedge j \neq i}^{n} (\mathcal{T})_{ij} < 1 < |\lambda|$ for all $i \in \underline{n}$. Thus, any strictly row-substochastic matrix $\mathcal{T}$ has no eigenvalues $|\lambda| > 1$. □

Definition 2.6 (Jordan normal form)
Every square matrix $A \in \mathbb{C}^{n\times n}$ is similar to a block diagonal matrix $J \in \mathbb{C}^{n\times n}$, also called the Jordan normal form of A, meaning $A = PJP^{-1}$ for an invertible matrix $P \in \mathbb{C}^{n\times n}$. The blocks on the diagonal of the Jordan matrix are called Jordan blocks J_i and have the form

$$J_i = \begin{bmatrix} \lambda_i & 1 & 0 & \dots & 0 \\ 0 & \lambda_i & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & 1 & 0 \\ 0 & \ddots & 0 & \lambda_i & 1 \\ 0 & 0 & \dots & 0 & \lambda_i \end{bmatrix},$$

where λ_i is the i-th eigenvalue of A.

Theorem 2.7 (Convergence of strictly row-substochastic matrices)
If a matrix $\mathcal{T}$ is strictly row-substochastic, then it is convergent, i.e. $\lim_{k\to\infty} \mathcal{T}^k = 0$

Proof 2
Given a strictly row-substochastic matrix $\mathcal{T}$, consider its Jordan normal form $\mathcal{T} = PJP^{-1}$, see def. 2.6. Thus, $\lim_{k\to\infty} \mathcal{T}^k = P \lim_{k\to\infty} J^k P^{-1}$. Because $|\lambda| < 1$ for all eigenvalues of $\mathcal{T}$, $\lim_{k\to\infty} J^k = 0$. Thus, $\lim_{k\to\infty} \mathcal{T}^k = 0$, i.e. $\mathcal{T}$ is convergent.□

Definition 2.8 (Matrix inversion by the Neumann Series)
Given a matrix $\mathcal{T} \in \mathbb{R}^{n\times n}$ with $\lim_{k\to\infty} \mathcal{T}^k = 0$, i.e. $\mathcal{T}$ is convergent, the Neumann series $\sum_{k=0}^{\infty} \mathcal{T}^k$ converges against $(\boldsymbol{I} - \mathcal{T})^{-1}$.

Definition 2.9 (Irreducible and Reducible Matrix)
A non-negative square matrix $\mathcal{T} \in \mathbb{R}^{n\times n}_{\geq 0} > 0$ with $n \geq 2$ is irreducible if and only if $\sum_{k=0}^{n-1} \mathcal{T}^k > 0$. If $\mathcal{T}$ is not irreducible, it is reducible.

Definition 2.10 (Permutation Matrix)
A binary square matrix $P \in \{0,1\}^{n\times n}$ is a permutation matrix if and only if

(i) $(P)_{ij} = 1 = (P)_{i'j}$ with $i \neq i'$

(ii) $(P)_{ij} = 1 = (P)_{ij'}$ with $j \neq j'$

$(P^{-1} = P^{\mathsf{T}})$

In other words exactly one entry of P is 1 per row and column.

Definition 2.11 (Upper and Lower Triangular Matrix)
An upper/lower triangular matrix is a square matrix $\mathcal{T}_{up}, \mathcal{T}_{lo} \in \mathbb{R}^{n \times n}$, where all entries below/above the diagonal are 0.

$$\mathcal{T}_{up} = \begin{bmatrix} a_{11} & \dots & a_{1n} \\ & \ddots & \vdots \\ 0 & & a_{nn} \end{bmatrix} \quad \mathcal{T}_{lo} = \begin{bmatrix} a_{11} & & 0 \\ \vdots & \ddots & \\ a_{n1} & \dots & a_{nn} \end{bmatrix}$$

Definition 2.12 (LU-decomposition)
Let $\mathcal{T}$ be a square matrix, then (and only then) there exists an upper triangular matrix U and a lower triangular matrix L, such that

$$\mathcal{T} = L \cdot U \ .$$

By convention, L is made unitriangular, meaning the entries on its diagnonal are equal to 1. L & U invertible: all triangular matrices inv. (if all $a_{ii} \neq 0$)

Blocks $\mathcal{T}_{ii}$ communicating groups/classes (in graph) + $_{ij}$ communicating between groups i&j??

Definition 2.13
Invertible matrices have full rank

Expected Values

Definition 2.14 (Discrete Probability Distribution)
A function P is a *discrete probability distribution*, if $P : X \to [0, 1]$ where X is a countable set and $\sum_{x \in X} P(x) = 1$.

Definition 2.15 (Conditional Probability)
$Pr(A \mid B) = \frac{Pr(A \cap B)}{Pr(B)}$

Definition 2.16 (Law of Total Probability)
Let $B = \{B_1, B_2, \dots\}$ be a countable set of mutually exclusive and collectively exhaustive events, then for any event A,

$$Pr(A) = \sum_{i=1}^{|B|} Pr(A \mid B_i) \cdot Pr(B_i)$$

Definition 2.17 (Monotone Convergence Theorem)
...

Chapter 3

Markov Chains

In this section, we will introduce definitions and theorems for Markov Chains which we will later require for section on Interval Markov Chains. The notation is mostly based on [BK08], [CSH08], [BDLM21] and [MKQW25].

Definition 3.1 (DTMC)
A discrete-time Markov chain (DTMC) is a tuple $\mathcal{D} = (S, \mathbf{P}, \iota_{init})$, where

(i) S is a countable, nonempty set of states,

(ii) $\mathbf{P} : S \times S \to [0, 1]$ is the transition probability function, with $\sum_{s' \in S} \mathbf{P}(s, s') = 1$ for all states $s \in S$, and

(iii) $\iota_{init} : S \to [0, 1]$ is the initial distribution, such that $\sum_{s \in S} \iota_{init}(s) = 1$.

We assume in this thesis, that the DTMCs are finite, i.e., that the set of states S is finite. Furthermore, we assume an arbitrary but fixed linear order $\prec$ on the set S, such that the states $s \in S$ can be indexed. Thus, $s_i \prec s_j$ if $s, s' \in S$ and $i < j$. Due to the conditions (i) and (ii) provided in def. 3.1, the transition probability function $\mathbf{P}$ and the initial distribution ι_{init} are discrete probability distributions.

??(notion of a current state)

Moreover, if $\mathbf{P}(s, s') > 0$ for two given $s, s' \in S$, then it is possible to move from s to s' in one step. We refer to these states s' as direct successors of s (and to s as a direct predecessor of all states s'). Additionally, we define a neighbour set N, with $s' \in N(s)$, if and only if s' is a direct successor of s. For convenience, we define the set $S^{\mathcal{D}}_{in} = \{s \mid \iota^{\mathcal{D}}_{init}(s) > 0\}$.

Definition 3.2 (Transition Matrix)
The transition matrix $\mathcal{T}^{\mathcal{D}}$ of a DTMC $\mathcal{D}$ is a square matrix with size $|S| \times |S|$,

such that $(\mathcal{T}^{\mathcal{D}})_{ij} = \mathbf{P}(s_i, s_j)$.

Notice that due to the condition, given in (ii) of def. 3.1, that $\sum_{s_j \in S} \mathbf{P}(s_i, s_j) = 1$ for all states $s \in S$ and thus by defintion $\sum_{j=0}^{|S|} (\mathcal{T})_{ij} = 1$ for all rows $i \in \underline{n}$. In other words, the transition matrix $\mathcal{T}$ of a DTMC $\mathcal{D}$ is row-stochastic, see def. 2.3.

Theorem 3.3 (k-step Transition Probability)
The k-th power of the transition matrix $(\mathcal{T}^k)_{ij}$ characterizes the probability of transitioning from state i to j in precisely k steps, i.e., $Pr(\pi[k] = s_j \mid \pi[0] = s_i)$.

We prove this theorem after introducing further concepts/definitions in proof 3.

Definition 3.4 (Path of a DTMC)
An *infinite path* π in a DTMC $\mathcal{D}$ is a sequence $\pi = (s_0, s_1, \dots)$ of states with $s_0, s_1, \dots \in S$, such that $\iota_{init}(\pi[0]) > 0$ and $\mathbf{P}(\pi[k], \pi[k+1]) > 0$, where $\pi[k] \in S$ denotes the k-th element of the sequence/path for all $k \in \mathbb{N}_0$, i.e. $\pi[k] = s_k$. A *finite path* is a finite prefix of an *infinite path*.

Furthermore, we define the length $|\pi|$ of a finite path $\pi = (s_0, \dots, s_k) \in Paths^{\mathcal{D}}_{fin}$ to be equal to k and for an infinite path $\pi \in Paths^{\mathcal{D}}_{inf}$ to be $|\pi| = \infty$. The sets of all infinite and finite sets are denoted as $Paths^{\mathcal{D}}_{inf}$ and $Paths^{\mathcal{D}}_{fin}$, respectively.

Further, we construct the probability space of a DTMC as follows.

Definition 3.5 (Probability Space of a DTMC)
The *probability space* of a DTMC $\mathcal{D}$ is defined as $(Paths^{\mathcal{D}}_{inf}, \sum_{Paths^{\mathcal{D}}_{inf}}, Pr^{\mathcal{D}})$, where

- $\sum_{Paths^{\mathcal{D}}_{inf}}$ denotes the smallest σ-algebra containing the cylinder sets $Cyl(\pi')$ for all finite paths $\pi' \in Paths^{\mathcal{D}}_{fin}$ with

$$Cyl(\pi') = \{\pi \in Paths^{\mathcal{D}}_{inf} \mid \forall k \in \underline{|\pi'|} : \pi[k] = \pi'[k]\}$$

- $Pr^{\mathcal{D}}$ is the unique probability measure on $(Paths^{\mathcal{D}}_{inf}, \sum_{Paths^{\mathcal{D}}_{inf}})$ satisfying

$$Pr^{\mathcal{D}}(Cyl(\pi')) = \iota^{\mathcal{D}}_{init}(\pi'[0]) \cdot \prod_{k=0}^{|\pi'|-1} \mathbf{P}^{\mathcal{D}}(\pi'[k], \pi'[k+1])$$

  for all finite paths $\pi' \in Paths^{\mathcal{D}}_{fin}$

Additionally, we will use the notation $Pr_s(\pi)$ to denote the probability measure on the measurable space induced by fixing $s \in S$ as the unique initial state.

Definition 3.6 (Path Fragment)
Given a path π in a DTMC, a path fragment π' is an infix of π. In other words, a path fragment is a path with an arbitrary value for $\iota_{init}(\pi'[0])$.

Definition 3.7 (Until Operator)
Let $R, T \subseteq S^{\mathcal{D}}$ be two sets of states of a DTMC $\mathcal{D}$. For these two sets, we define the "until" Operator as follows,

$$R \cup T = \{\pi \in Paths^{\mathcal{D}}_{inf} \mid \exists k \in \mathbb{N} : \pi[k] \in T \wedge \forall i < k : \pi[i] \in R\} \ .$$

In other words, the set $R \cup T$ contains all paths that only visit states in R, until a state in T is visited.

With the "until" operator, we define the reachability probability.

Definition 3.8 (Reachability Probability)
The reachability probability of a set $T \subseteq S^{\mathcal{D}}$ in a DTMC $\mathcal{D}$ is given by $Pr(\Diamond T)$, where $\Diamond T = S \cup T$. The operator $\Diamond$ is also called the "eventually" operator.

As $S^{\mathcal{D}}$ defines all states in $\mathcal{D}$, this simply defines the probability of visiting a state $t \in T$ at an arbitrary step in the future, i.e. if $Pr(\Diamond T) > 0$ at least one state $t \in T$ is reachable. In case $T = \{t\}$ is a singleton set, we omit the braces and denote $\Diamond t$ instead of $\Diamond\{t\}$ (and $S \cup t$ instead of $S \cup T$). Lastly, we denote $\Diamond_t T = \{\pi \in \Diamond T \mid \pi[0] = t\}$

$\sum_{Paths^{\mathcal{D}}_{inf}}$ DEFINIEREN (?)

Definition 3.9 (Characterizing Reachability Probability for Path Fragments)
For some set of paths $\Diamond S$ and an intial state t.

$$Pr^{\mathcal{D}}_t(\Diamond S) = \sum_{\pi \in \Diamond_t S} \prod_{k=0}^{|\pi|-1} \mathbf{P}^{\mathcal{D}}(\pi[k], \pi[k+1])$$

Now that we have defined the probability space for DTMCs, we can prove theorem 3.3.

Proof 3
We will prove this theorem by induction.

Notice, that $Pr(\pi[k] = s_j \mid \pi[0] = s_i)$ for some $s_i, s_j \in S$ is precisely $Pr_{s_i}(\{\pi \in \Diamond_{s_i} s_j \mid |\pi| = k\})$. By defintion, see def. 3.2, $(\mathcal{T})_{ij} = \mathbf{P}(s_i, s_j) = Pr_{s_i}(\{\pi \in \Diamond_{s_i} s_j \mid |\pi| = 1\})$, thus the theorem holds for $k = 1$. Now, assume that the theorem holds for any $k \in \underline{n}$. As $(\mathcal{T}^{k+1})_{ij} = (\mathcal{T}^k \cdot \mathcal{T})_{ij} = \sum_{m=1}^{n} (\mathcal{T}^k)_{im} \cdot (\mathcal{T})_{mj}$. Using our induction hypothesis, $(\mathcal{T}^k)_{im}$ characterizes the probability of transition from state

i to m in k steps, for all $m \in \underline{n}$

We use the LTL unary prefix operator $\bigcirc$ to denote the temporal modality "next", meaning

Definition 3.10 (Next Step Probability)
Let $\mathcal{D}$ be a DTMC and let $E \in \sum_{Paths^{\mathcal{D}}_{inf}}$ be a (countable) set of infinite paths in $\mathcal{D}$. We now define the "next step" operator, as

$$\bigcirc E = \{\pi \in Paths^{\mathcal{D}}_{inf} \mid (\pi[1], \pi[2], \dots) \in E\} \ ,$$

i.e., a path satisfies $\bigcirc E$ if and only if after removing the first state $\pi[0]$, the remaining path belongs to E. Moreover, the next step probability is defined as $Pr(\bigcirc E) = \sum_{s' \in N(s)} \mathbf{P}(s, s') \cdot Pr_{s'}(E)$.

In this thesis, we will use the "next step" operator $\bigcirc$ in combination with the "eventually" operator $\Diamond$. Specifically, we use them to denote the return probability $Pr_s(\bigcirc\Diamond s)$, i.e., the probability to reach s from all of its successors.

Definition 3.11 ((Bottom) Strongly Connected Component)
A set of states $C \subseteq S^{\mathcal{D}}$ of a DTMC $\mathcal{D}$ is strongly connected if and only if for every pair of states $s, t \in S$, there exists a path $\pi \in Paths^{\mathcal{D}}_{inf}$, such that $\pi[0] = s$ and $\pi[k]$ for some $k \in \mathbb{N}$. If C is strongly connected and no proper subset, i.e. $C' \subset C$, is strongly connected, then C is called a strongly connected component (SCC).

Furthermore, an SCC C is called a bottom stronlgy connected component (BSCC) if there exists no $k \in \mathbb{N}$ and no path $(\pi[0], \dots, \pi[k], \dots) \in Paths^{\mathcal{D}}_{inf}$ such that $\pi[0] \in C$ and $\pi[k] \notin C$.

In other words, once the model enters a BSCC, it cannot leave it again. Notice, that any grouping

SCC (minimal amount of states of set such that still strongly connec) (Acyclic DTMC)

The states can be classified ...

Definition 3.12 (Transient and Recurrent states)
A state $s \in S$ is called recurrent if and only if $Pr_s(\bigcirc\Diamond s) = 1$. All states that are not recurrent are called transient.

In other words, a state is recurrent precisely when we are guaranteed to reach it from all of its direct successors, meaning once we reach a recurrent state, we are bound to return to it. (Thus reaching it infinitely often) Using these definitions, we can split the states S of a DTMC into two sets with S_{tr} containing all of the transient states and S_{re} containing all of the recurrent states. Furthermore, we define $S_{tr \cap in} \subseteq S_{in}$ with $S_{tr \cap in} = S_{in} \cap S_{tr}$. Additionally, we define an absorbing state.

Definition 3.13 (Absorbing state)
A states s of a DTMC $\mathcal{D}$ is refered to as an absorbing state if and only if $\mathbf{P}^{\mathcal{D}}(s,s) = 1$.

This means, once the model enters an absorbing state, it cannot leave it. Notice, that absorbing states are a special type of recurrent states, as the only direct successor of an absorbing state $s \in S$ is s itself and therefore trivially $Pr_s(\Diamond s) = 1$.

Notice further, that intuitively the Markov Chain almost surely reaches a recurrent state in any DTMC.

Theorem 3.14 (Limit Behaviour)
For any DTMC $\mathcal{D}$, $s \in S$ with $\iota_{init}(s) > 0$ it holds that $Pr(\Diamond S_{re}) = 1$

Definition 3.15 (Acyclic DTMC)
A DTMC $\mathcal{D}$ is called acyclic if and only if each SCC is either a BSSC or a singleton set $\{s\}$ with $\mathbf{P}(s,s) = 0$

Definition 3.16 (Stopping Time)
A random variable $T : Paths^{\mathcal{D}}_{inf} \to \mathbb{N} \cup \{\infty\}$ is called a *stopping time* if, for every $n \geq \mathbb{N}$,

Definition 3.17 (Strong Markov Property)

Definition 3.18 (Canonical Form)
For a DTMC $\mathcal{D}$, its transition matrix $\mathcal{T}$ is said to be in canonical form if and only if

$$\mathcal{T} = \begin{bmatrix} Q & R \\ 0 & S \end{bmatrix}$$

where

(i) $Q = (\mathbf{P}(s,t))_{s,t \in S_{tr}}$ contains the transition probabilities between the transient states

(ii) $R = (\mathbf{P}(s,t))_{s \in S_{tr}, t \in S_{re}}$ contains the transition probabilities between transient and recurrent state pairs

(iii) $S = (\mathbf{P}(s,t))_{s,t \in S_{re}}$ contains the transition probabilities between the transient states

For absorbing DTMCs $S = \boldsymbol{I}$. As mentioned in **??**, this is precisely possible for

(Reward Structure)

3.1 Expected Visiting Times for DTMCs

Intuitively, the notion of EVTs describes how often we expect the model to be in a given state. To formally introduce EVTs, we fix a DTMC $\mathcal{D}$ and start by introducing *visiting times* along paths.

Definition 3.19 (Visiting Times of Infinite Paths)
The function $vt_s : Paths^{\mathcal{D}}_{inf} \to \mathbb{R}$ describes the *visiting time* of a state $s \in S$ along infinite paths in $\mathcal{D}$, where for $\pi \in Paths^{\mathcal{D}}_{inf}$

$$vt_s(\pi) = \sum_{k=0}^{\infty} [\![\pi[k] = s]\!]$$

Borel Measurability

$\to$ Thus, the Lebesgue integral of the non-negative function vt_s is well-defined for all states $s \in S^{\mathcal{D}}$. With this function, we can now define the EVT of a state s as the expected value of vt_s over all possible paths.

Definition 3.20 (Expected Visiting Times in DTMCs)
For a state $s \in S$, the *expected visiting time* of s is characterized as

$$\mathbb{E}^{\mathcal{D}}[vt_s] = \int_{\pi \in Paths^{\mathcal{D}}_{inf}} vt_s(\pi) dPr^{\mathcal{D}}(\pi) \ .$$

Additionally, we denote the *expected visiting time* that is obtained by fixing a certain state $t \in S$ as the only initial state as $\mathbb{E}^{\mathcal{D}}_t[vt_s]$. It is defined as

$$\mathbb{E}^{\mathcal{D}}_t[vt_s] = \int_{\pi \in Paths^{\mathcal{D}}_{inf} \wedge \pi[0]=t} vt_s(\pi) dPr^{\mathcal{D}}(\pi) \ .$$

As we defined the probability measure of paths π through a product containing the factor $\iota^{\mathcal{D}}_{init}(\pi[0])$ (see definition def. 3.5), it is nonzero only for paths that begin in a state s with $\iota^{\mathcal{D}}_{init}(s) > 0$. Thus the value $\mathbb{E}^{\mathcal{D}}[vt_s]$ gives the number of times the state s is expected to be visited from all paths in $\mathcal{D}$ that start in one of the initial states. Whenever it is clear from the context, we will omit the DTMC $\mathcal{D}$ from $\mathbb{E}^{\mathcal{D}}[vt_s]$ and $\mathbb{E}^{\mathcal{D}}_t[vt_s]$.

Using the aforementioned classification of the states of $\mathcal{D}$ into transient and recurrent states in def. 3.12, we can already give a characterization for the latter.

Lemma 3.21 (EVTs of Recurrent States)
For all recurrent states $s \in S_{re}$, it holds that

$$\mathbb{E}[vt_s] = \begin{cases} \infty & \text{if s is reachable,} \\ 0 & \text{else.} \end{cases}$$

For a proof, see [MKQW25].

Thus, we will focus on finding the non-trivial expected visiting times of the transient states.

We, will now see how the expected visiting times of the transient states can be calculated through a linear equation system. For this, we will first define the *fundamental matrix* which was introduced by Kemeny and Snell [KS76].

Definition 3.22 (Fundamental Matrix)
For a DTMC $\mathcal{D}$ with the matrix $Q = (\mathbf{P}^{\mathcal{D}}(s,t))_{s,t \in S_{tr}}$ containing the transition probabilities between the transient states (defined in def. 3.18), the fundamental matrix F is defined as,

$$F = (I - Q)^{-1} .$$

Theorem 3.23
The matrix Q, see def. 3.18, is strictly row-substochastic.

Proof 4
First, notice that there always needs to exist at least one recurrent state. We will prove this through a proof by contradiction. Assume that a Markov Chain $\mathcal{D}$ consists only of transient states. According to def. 3.12 this means, that $Pr_s(\bigcirc\diamond s) < 1$ for all states $s \in S$. Thus there exists a path $\pi \in Paths^{\mathcal{D}}_{inf}$ for all $s \in S$ such that for some $k \in \mathbb{N}$ $Pr_{\pi[k]}(\bigcirc\diamond s) = 0$, i.e. after step k it is impossible to return to state s. However, as the state $\pi[k]$, by our assumption, is transient as well, there need to exist $\pi' \in Paths^{\mathcal{D}}_{inf}$ and $k' \in \mathbb{N}$ such that $Pr_{\pi'[k']}(\bigcirc\diamond\pi[k]) = 0$. By repeating this argument (at most $|S|$ times), we arrive at an contradiction, where we necessarily transition to a state that is no longer reachable. Thus the statement is proven. Next, notice that there always needs to exist a $s \in S_{tr}$ and $s' \in S_{re}$ such that $\mathbf{P}(s,s') > 0$, i.e. there exists at least one transient state that as an edge to a recurrent state. By assumption all states in any given DTMC $\mathcal{D}$ are reachable and thus also a transient state $s \in S_{tr}$. Using the same argument as for the first step, we can conclude that a recurrent state $s' \in S_{re}$ must be reachable from s, i.e. $Pr_s(\bigcirc\diamond) = 1$. Thus there necessarily exists an edge from transient to recurrent. Through condition (ii) of def. 3.1, we know that $\sum_{j=0}^{n}(Q)_{ij} \leq 1$ for

all $i \in |S_{tr}|$, i.e. Q is row-substochastic. As at least one transient state s has an edge $\mathbf{P}(s, s') > 0$, with $s' \in S_{re}$, it follows that $\sum_{j=0}^{n}(Q)_{sj} < 1$,i.e. Q is strictly row-substochastic. □

As Q is strictly row-substochastic, we can thus conclude using theorem 2.7, that Q is also convergent. Therefore, the Neumann series of Q converges with $\sum_{k=0}^{\infty} Q^k = (\boldsymbol{I} - Q)^{-1}$, i.e. the fundamental matrix (see def. 3.22) is well-defined.

Furthermore, in [KS76] Kemeny and Snell prove that each entry $(F)_{ij} = \mathbb{E}_i[vt_j]$. From this equation, it intuitively follows that transient states are only visited a finite amount of times. With this characterization of EVTs for transient states, we can prove the following lemma.

Lemma 3.24 (Reachability of Transient States)
Transient states are reachable only from other transient states. Moreover, it follows from this that if a state is reachable from a recurrent state, it must also be recurrent

We provide a short proof by contradiction

Proof 5
Assume that a transient state s_{tr} is reachable from a recurrent state s_{re}, i.e. $Pr_{s_{re}}(\bigcirc\diamond s_{tr}) > 0$. As $Pr_{s_{re}}(\bigcirc\diamond s_{re}) = 1$, see def. 3.12, necessarily s_{re} is reachable from s_{tr} as well and crucially $Pr_{s_{tr}}(\bigcirc\diamond s_{re}) = 1$. We now view each visit of s_{re} as an indepenent trial, due to the strong Markov property def. 3.17. We can then describe the probability of not reaching s_{tr} as $\lim_{N\to\infty}(1 - Pr_{s_{re}}(\bigcirc\diamond s_{tr}))^N = 0$, as s_{re} is visited infinitely often, see lemma 3.21. Thus, we are guarenteed to visit s_{tr}, when starting in s_{re}. As $Pr_{s_{tr}}(\bigcirc\diamond s_{re}) = 1$, this process repeats each time s_{tr} is visited. Therefore, s_{tr} is visited infinitely often as well, i.e. $\mathbb{E}[vt_{s_{tr}}] = \infty$. This contradicts the characterization of the EVTs of transient states, proven by [KS76]. Thus, we conclude that transient states cannot be reachable from recurrent states, i.e. can only be reached from other transient states. □

Moreover, we can use this result to define a general characterization for the EVTs of transient states using the initial distribution.

Theorem 3.25 (Calculating EVTs using the Fundamental Matrix)
Given a DTMC $\mathcal{D}$ and the corresponding matrix Q, we can define

$$\mathbb{E}[vt_s] = \sum_{t \in S_{tr} \cap in} (F)_{ts} \cdot \iota_{init}(t)$$

Proof 6
As defined in def. 3.20,

$$\mathbb{E}^{\mathcal{D}}[vt_s] = \int_{\pi \in Paths^{\mathcal{D}}_{inf}} vt_s(\pi) dPr^{\mathcal{D}}(\pi) \ .$$

We now substitute the visit time function $vt_s(\pi)$, using def. 3.19.

$$\begin{aligned}\mathbb{E}^{\mathcal{D}}[vt_s] &= \int_{\pi \in Paths^{\mathcal{D}}_{inf}} \sum_{k=0}^{\infty} [\![\pi[k] = s]\!] \, dPr^{\mathcal{D}}(\pi) \\ &= \sum_{k=0}^{\infty} \int_{\pi \in Paths^{\mathcal{D}}_{inf}} [\![\pi[k] = s]\!] \, dPr^{\mathcal{D}}(\pi), \quad \text{(by def. 2.17 and since } vt_s \text{ is non-negative).}\end{aligned}$$

Since $\int_{\pi \in Paths^{\mathcal{D}}_{inf}} [\![\pi[k] = s]\!] dPr^{\mathcal{D}}(\pi) = Pr^{\mathcal{D}}(\pi[k] = s)$, we can thus conclude that

$$\mathbb{E}^{\mathcal{D}}[vt_s] = \sum_{k=0}^{\infty} Pr^{\mathcal{D}}(\pi[k] = s) \ .$$

Using the *Law of Total Probability*, we can characterize the probability $Pr^{\mathcal{D}}(\pi[k] = s)$ by conditioning it on the possible initial states. Thereby, $Pr^{\mathcal{D}}(\pi[k] = s) = \sum_{t \in S_{in}} Pr^{\mathcal{D}}(\pi[k] = s \mid \pi[0] = t) \cdot Pr^{\mathcal{D}}(\pi[0] = t)$, where $Pr^{\mathcal{D}}(\pi[0] = t) = \iota_{init}(t)$. We now substitute $Pr^{\mathcal{D}}(\pi[k] = s)$ with this term to get

$$\begin{aligned}\mathbb{E}^{\mathcal{D}}[vt_s] &= \sum_{k=0}^{\infty} \sum_{t \in S_{in}} Pr^{\mathcal{D}}(\pi[k] = s \mid \pi[0] = t) \, \iota_{init}(t) \\ &= \sum_{k=0}^{\infty} \sum_{t \in S_{tr \cap in}} Pr^{\mathcal{D}}(\pi[k] = s \mid \pi[0] = t) \, \iota_{init}(t), \qquad \text{(by lemma 3.24)} \\ &= \sum_{t \in S_{tr \cap in}} \left(\sum_{k=0}^{\infty} Pr^{\mathcal{D}}(\pi[k] = s \mid \pi[0] = t) \right) \iota_{init}(t), \qquad \text{(all terms are non-negative)} \\ &= \sum_{t \in S_{tr \cap in}} \left(\sum_{k=0}^{\infty} Pr^{\mathcal{D}}(\pi[k] = s \mid \pi[0] = t) \right) \cdot \iota_{init}(t)\end{aligned}$$

Using theorem 3.3, we can now substitute $Pr^{\mathcal{D}}(\pi[k] = s \mid \pi[0] = t) = (Q^k)_{ts}$. Notice that this is true, as crucially s can only be reached from paths that contain

exclusively transient states, see lemma 3.24. Thus, $\sum_{k=0}^{\infty} Pr^{\mathcal{D}}(\pi[k] = s \mid \pi[0] = t) = \sum_{k=0}^{\infty}(Q^k)_{ts} = (F)_{ts}$. Finally, we can derive the final equation

$$\mathbb{E}[vt_s] = \sum_{t \in S_{tr} \cap in} (F)_{ts} \cdot \iota_{init}(t) \ .$$

□

Let us now define $\tau_{init} = (\iota_{init}(s_1), \ldots, \iota_{init}(s_n))^{\mathsf{T}}$ for all $s \in S_{tr}$, so that we can rewrite theorem 3.25 to $\mathbb{E}[vt_s] = (F^{\mathsf{T}})_s \cdot \tau_{init}$, where $(F^{\mathsf{T}})_s$ denotes the row vector of F^{T} corresponding to the state s. As introduced in def. 3.22, $(F)_s = ((I-Q)^{-1})_s$, from which we can conclude that $\mathbb{E}[vt_s] = ((I - Q^{\mathsf{T}})^{-1})_s \cdot \tau_{init}$ or $\mathcal{E} = (I - Q^{\mathsf{T}})^{-1} \cdot \tau_{init}$, where $\mathcal{E} = (\mathbb{E}[vt_{s_1}], \ldots, \mathbb{E}[vt_{s_n}])^{\mathsf{T}}$. However, as the calculation of the inverse of a high-dimensional matrix becomes infeasible for large models, we will take a look at how the EVTs can be calculated without the fundamental matrix.

For this, we rewrite the equation $\mathcal{E} = (I - Q^{\mathsf{T}})^{-1} \cdot \tau_{init}$ to $\mathcal{E} = \tau_{init} + Q^{\mathsf{T}} \cdot \mathcal{E}$. For a single state $s \in S_{tr}$, we can thus calculate the respective EVT as follows,

$$\mathbb{E}[vt_s] = \iota_{init}(s) + \sum_{t \in S_{tr}} \mathbf{P}(t, s) \cdot \mathbb{E}[vt_t] \ . \tag{3.1}$$

3.1.1 Relating Expected Visiting Times and Quantitative Reachability

Lemma 3.26
Given a DTMC $\mathcal{D}$ and its fundamental matrix F, we can define

$$(F)_{ts} = Pr_t(\Diamond s) \cdot (F)_{ss}$$

for all $s \neq t \in S_{tr}$

Trivially, all paths that reach s from t at least once visit s. Thus, intuitively, the *expected visiting time* of s starting in t is simply the product of the probability of reaching s and the *expected visiting time* of s starting in s.

Lemma 3.27
Given a DTMC $\mathcal{D}$ and its fundamental matrix F, we can define

$$(F)_{ss} = \frac{1}{1 - Pr_s(\bigcirc \Diamond s)}$$

for all $s \in S_{tr}$, where $Pr_s(\bigcirc \Diamond s) = \sum_{t \in N(s)} \mathbf{P}(s, t) \cdot Pr_t(\Diamond s)$

(The fraction is well-defined as $Pr_s(\bigcirc \Diamond s) < 1$, see def. 3.12)

Proof 7
As the only initial state is s, it is necessarily visited at least once. After that the probability that s will be visited at least one more time is $Pr_s(\bigcirc\Diamond s)$ and the probability that it will not, is accordingly $(1 - Pr_s(\bigcirc\Diamond s))$. Moreover, the probability of visiting s exactly k times is precisely the probability of returning $k-1$ times and not return again afterwards. Thus, $Pr_s(N_s = k) = Pr_s(\bigcirc\Diamond s)^{k-1}(1 - Pr_s(\bigcirc\Diamond s))$ for $k \in \mathbb{N}$, where N_s is a random variable representing the total number of time s is visited (Pr_s means starting in s). Intuitively,

$$
\begin{aligned}
(F)_{ss} &= \sum_{k=1}^{\infty} k \cdot Pr_s(N_s = k) \\
&= \sum_{k=1}^{\infty} k \cdot Pr_s(\bigcirc\Diamond s)^{k-1}(1 - Pr_s(\bigcirc\Diamond s)) \\
&= (1 - Pr_s(\bigcirc\Diamond s)) \sum_{k=1}^{\infty} k \cdot Pr_s(\bigcirc\Diamond s)^{k-1} \\
&= (1 - Pr_s(\bigcirc\Diamond s)) \cdot \frac{1}{(1 - Pr_s(\bigcirc\Diamond s))^2} = \frac{1}{1 - Pr_s(\bigcirc\Diamond s)} \qquad \text{(by geometric series.)}
\end{aligned}
$$

□

Finally, using theorem 3.25,

$$
\begin{aligned}
\mathbb{E}[vt_s] &= \sum_{t \in S_{tr} \cap in} (F)_{ts} \cdot \iota_{init}(t) \\
&= \sum_{t \in S_{tr} \cap in} Pr_t(\Diamond s) \cdot (F)_{ss} \cdot \iota_{init}(t) \qquad \text{(by lemma 3.26)} \\
&= \sum_{t \in S_{tr} \cap in} \frac{Pr_t(\Diamond s)}{1 - Pr_s(\bigcirc\Diamond s)} \cdot \iota_{init}(t) \qquad \text{(by lemma 3.27)} \qquad (3.2)
\end{aligned}
$$

Notice, that for any given DTMC $\mathcal{D}$ we can construct $\mathcal{D}_{uni} = (S^{\mathcal{D}} \cup \{s_{in}\}, \mathbf{P}^{\mathcal{D}_{uni}}, \iota_{init}^{\mathcal{D}_{uni}})$, where $\mathbf{P}^{\mathcal{D}_{uni}}(s, s') = \mathbf{P}^{\mathcal{D}}(s, s')$, $\mathbf{P}^{\mathcal{D}_{uni}}(s_{in}, s) = \iota_{init}^{\mathcal{D}}(s)$ for all $s, s' \in S^{\mathcal{D}}$ and $\iota_{init}^{\mathcal{D}_{uni}}(s_{in}) = 1$. Essentially, $\mathcal{D}_{uni}$ is a copy of $\mathcal{D}$, where instead of using ι_{init} as an initial distribution, we simply have an initial state s_{in}.

Theorem 3.28
The reachability probabilities and thus also EVTs in $\mathcal{D}$ and the corresponding $\mathcal{D}_{uni}$ are identical.

Proof 8
First, we show that the reachability probabilities are equal (see def. 3.7) $Pr^{\mathcal{D}}(\Diamond T) =$

$Pr^{\mathcal{D}_{uni}}(\Diamond T)$ for all $T \in S^{\mathcal{D}} \backslash \{s_{in}\}$. For this, we define the bijection $\phi : Paths_{inf}^{\mathcal{D}_{uni}} \to Paths_{inf}^{\mathcal{D}}, \pi = (s_{in}, s_1, s_2, \dots) \mapsto (s_1, s_2, \dots)$. This bijection is well defined, as $\mathbf{P}^{\mathcal{D}_{uni}}(s, s') = \mathbf{P}^{\mathcal{D}}(s, s')$ for all $s, s' \in S^{\mathcal{D}}$ and $N(s_{in}) = \{s \mid \iota_{init}^{\mathcal{D}}(s) > 0\}$. Moreover,

$$\begin{aligned}
Pr^{\mathcal{D}_{uni}}(\pi) &= \underbrace{\iota_{init}^{\mathcal{D}_{uni}}(\pi[0])}_{=1} \cdot \prod_{k=0}^{|\pi|-1} \mathbf{P}^{\mathcal{D}_{uni}}(\pi[k], \pi[k+1]) \\
&= \mathbf{P}^{\mathcal{D}_{uni}}(\pi[0], \pi[1]) \cdot \prod_{k=1}^{|\pi|-1} \mathbf{P}^{\mathcal{D}_{uni}}(\pi[k], \pi[k+1]) \\
&= \iota_{init}^{\mathcal{D}}(\phi(\pi)[1]) \cdot \prod_{k=1}^{|\pi|-1} \mathbf{P}^{\mathcal{D}}(\phi(\pi)[k], \phi(\pi)[k+1]) \\
&= \iota_{init}^{\mathcal{D}}(\hat{\pi}[0]) \cdot \prod_{k=0}^{|\hat{\pi}|-1} \mathbf{P}^{\mathcal{D}}(\hat{\pi}[k], \hat{\pi}[k+1]) \\
&= Pr^{\mathcal{D}}(\hat{\pi})
\end{aligned}$$

for all $\pi \in Paths_{inf}^{\mathcal{D}_{uni}}$ with $\phi(\pi) = \hat{\pi}$. Therefore, we can conclude $Pr^{\mathcal{D}_{uni}}(\Diamond T) = Pr^{\mathcal{D}}(\Diamond T)$, as $\Diamond T \subseteq Paths_{inf}^{\mathcal{D}_{uni}}$. Using eq. (3.2), we can then also conclude that the EVTs of $\mathcal{D}$ and $\mathcal{D}_{uni}$ must be equal. □

Thus, the reachability probabilities $Pr_{s_{in}}^{\mathcal{D}_{uni}}(\Diamond s) = \sum_{t \in S_{tr}} Pr_t^{\mathcal{D}}(\Diamond s)$ are equal. Thus, we can simply the previous expression,

$$\mathbb{E}^{\mathcal{D}}[vt_s] = \frac{Pr_{s_{in}}^{\mathcal{D}_{uni}}(\Diamond s)}{1 - Pr_s^{\mathcal{D}_{uni}}(\bigcirc \Diamond s)} \cdot \iota_{init}(s_{in}) (= \frac{Pr^{\mathcal{D}}(\Diamond s)}{1 - Pr_s^{\mathcal{D}}(\bigcirc \Diamond s)} \cdot \iota_{init}(s_{in})) \tag{3.3}$$

Moreover, using lemma 3.26 and the construction $\mathcal{D}_{uni}$, we can calculate the quantitative reachability of a given state with $Pr(\Diamond s) = \frac{(F^{\mathcal{D}_{uni}})_{ts}}{(F)_{ss}}$.

3.2 Consideration for limiting the state size

In sec. 3.1, we concluded that due to lemma 3.21, only the transient states need to be considered for calculating EVTs. As we are interested in the calculation of EVTs, we will only consider DTMCs and IDTMCs with a finite amount of transient states.

Moreover, we will only consider DTMCs (and IDTMCs) where all states are reachable, as if they are not the EVTs of those states are trivially 0 and have no impact on the EVTs of the other states

Chapter 4

Interval Markov Chains

The focus of this thesis are Interval Markov Chains. In the following sections we will mainly discuss how the expected visiting times can be calculated and shortly examine stationary distributions of Interval Markov Chains.

We start by introducing concepts which we will use in the following sections.

Definition 4.1 (Interval Markov Chain)
An interval discrete-time Markov chain (IDTMC) is a tuple $\mathcal{I} = (S, \underline{\mathbf{P}}, \overline{\mathbf{P}}, \underline{\iota}_{init}, \overline{\iota}_{init})$, where

(i) S is a countable, nonempty set of states,

(ii) $\underline{\mathbf{P}} : S \times S \to [0,1]$ is the transition probability function, where $\underline{\mathbf{P}}(s, s')$ gives the lower bound on the transition probability from state s to s',

(iii) $\overline{\mathbf{P}} : S \times S \to [0,1]$ is the transition probability function, where $\overline{\mathbf{P}}(s, s')$ gives the upper bound on the transition probability from state s to s',

(iv) $\underline{\iota}_{init} : S \to [0,1]$ is the upper bound on the initial distribution, and

(v) $\overline{\iota}_{init} : S \to [0,1]$ is the upper bound on the initial distribution.

To resolve the uncertainty caused by the intervals, we need to define semantics to interpret the Interval Markov Chain. In other words we need to resolve an Interval Markov Chain to some set of Markov Chains that correspond to it. We say that all of these DTMCs $\mathcal{D}_i$ satisfy the IDTMC $\mathcal{I}$ and denote it as $\mathcal{D}_i \models \mathcal{I}$. We assume in this thesis that there exists at least one satisfying DTMC for every IDTMC.

In this paper, we will consider two popular semantics: the *once-and-for-all* or *Uncertain Markov Chain (UMC) semantic* and the *IMDP semantic*. We will first define the former, similar to [BDLM21, CSH08, BLW13].




Figure 4.1: IDTMC $\mathcal{I}$

Definition 4.2 (Once-and-for-all/UMC Semantic)
An arbitrary DTMC $\mathcal{D}$ satisfies an IDTMC $\mathcal{I}$ if and only if $\underline{\mathbf{P}}^{\mathcal{I}}(s,s') \leq \mathbf{P}^{\mathcal{D}}(s,s') \leq \overline{\mathbf{P}}^{\mathcal{I}}(s,s')$ and $\underline{\iota}^{\mathcal{I}}_{init}(s) \leq \iota^{\mathcal{D}}_{init}(s) \leq \overline{\iota}^{\mathcal{I}}_{init}(s)$ for all $s, s' \in S^{\mathcal{I}} = S^{\mathcal{D}}$.

We will denote this as $\mathcal{D} \models_{\mathrm{U}} \mathcal{I}$.

In other words, any DTMC whose initial distribution and transition probabilities lie in the intervals defined by the IDTMC is satisfactory. Moreover, notice that the satisfaying DTMCs have the same state size (amount of states) as the IDTMC.

Example 4.3
Consider the IDTMC $\mathcal{I}$ in fig. 4.1. The DTMC $\mathcal{D}_U$ depicted in fig. 4.2 satisfies $\mathcal{I}$ under the UMC semantic, i.e., $\mathcal{D}_U \models_U \mathcal{I}$.




Figure 4.2: DTMC $\mathcal{D}_U$ with $\mathcal{D}_U \models_U \mathcal{I}$

The second semantic which we will define, is the *IMDP semantic*. There are two common definitions for this semantic. We will start by introducing the one that can

be found in [BDL+17] or [BDLM21] and then continue with the other definition found in [SVA06] or [CSH08].

Definition 4.4 (IMDP semantic 1)
An arbitrary DTMC $\mathcal{D} = (T, \mathbf{P}, \iota_{init})$ with $|T| \geq |S|$ (or $S \subseteq T$) satisfies an IDTMC $\mathcal{I} = (S, \underline{\mathbf{P}}, \overline{\mathbf{P}}, \underline{\iota}_{init}, \overline{\iota}_{init})$ if and only if there exists a surjective mapping $\tau : T \to S$ such that

(i) for every $s \in S$ there exists exactly one $t \in T$ with $\tau(t) = s$ such that $\underline{\iota}_{init}(s) \leq \iota_{init}(t) \leq \overline{\iota}_{init}(s)$. Thus, for all other $\tau(t_i) = s$ and $t_i \neq t \implies \iota_{init}(t_i) = 0$.

(ii) $\underline{\mathbf{P}}(\tau(t), \tau(t')) \leq \mathbf{P}(t, t') \leq \overline{\mathbf{P}}(\tau(t), \tau(t'))$ for all $t, t' \in T$.

We will denote this as $\mathcal{D} \models_I \mathcal{I}$.

Another common definition of the *IMDP semantic*

Definition 4.5 (IMDP semantic 2)
We will denote this as $\mathcal{D} \models_I \mathcal{I}$ (as well).

Example 4.6
Consider the IDTMC $\mathcal{I}$ in fig. 4.1. The DTMC $\mathcal{D}_I$ depicted in fig. 4.3 satisfies $\mathcal{I}$ under the UMC semantic, i.e., $\mathcal{D}_I \models_I \mathcal{I}$.




Figure 4.3: DTMC $\mathcal{D}_I$ with $\mathcal{D}_I \models_I \mathcal{I}$

Intuitively, the IMDP semantic allows for a probability distribution to be chosen at every step in each state within the constraints/bounds set by the IDTMC. In contrast to the UMC semantic, where each state in the satisfying DTMCs have a corresponding

state in the IDTMC and thus a fixed probability distribution, while the IMDP semantic allows for multiple states in the satisfying DTMCs with different fixed probability distributions to be mapped to the same corresponding state in the IDTMC. We can thus imagine an IDTMC under the IMDP semantic as a Markov Decision Process.

Definition 4.7 (Markov Descision Process (MDP))
[BK08] A *Markov Decision Process* (MDP) is a tuple $D = (S, Act, A, \mathbf{P}, \iota_{init})$, where

(i) S is a countable set of states

(ii) Act is a set of actions

(iii) $A(s) \subseteq Act$ denotes the applicable actions for all states $s \in S$

(iv) $\mathbf{P} : S \times Act \times S \to [0, 1]$ is the transition probability function such that $\sum_{s' \in S} \mathbf{P}(s, \alpha, s') = 1$ for all $s \in S$ and $\alpha \in A(s)$

(v) $\iota_{init} : S \to [0, 1]$ is the initial distribution such that $\sum_{s \in S} \iota_{init}(s) = 1$

The most intuitive reduction from an IDTMC $\mathcal{I}$ under the IMDP semantic to a MDP $\mathcal{M}$ is to define $S^{\mathcal{M}} = S^{\mathcal{I}}$ and an uncoutable set $\cup_{s \in S} A(s) = Act$ of actions where each action $\alpha \in A(s)$ corresponds to a probability distribution $\underline{\mathbf{P}}^{\mathcal{I}}(s, s') \leq \mathbf{P}^{\mathcal{M}}(s, \alpha, s') \leq \overline{\mathbf{P}}^{\mathcal{I}}(s, s')$ (for $s' \in N(s)$). In other words, the actions $\alpha \in A(s)$ in a given state $s \in S$ correspond to the probability distributions that can be chosen in s in any given step. However, as even the subsets $A(s)$ can be uncoutable, this reduction is trivially not "usable", as no algorithm, such as value or policy iteration, would terminate.

Instead, it is thus argued in [SVA06] and [CSH08], that the local constraints of each state of these MDPs and IDTMCs, see in def. 4.7 $(iv) - (v)$ and def. 4.1 $(ii) - (v)$, can be used to limit the amount of probability distributions that need to be mapped(, as we only need to consider the most extreme "behaviour"). In the theory of linear programming, this is called a basic feasible solution (BFS) .

(Only need to extend in the sense, that I need to say: my ι_{init} construction does not destroy anything: see $\mathcal{D}_{uni}$ (which has same EVTs etc. but no ι_{init}; which is simply 1 for $\iota_{init}(s_i n)$))

(Satisfying DTMC is policy)

In contrast to the UMC semantic, the IMDP semantic allows for satisfying DTMCs with larger state spaces than the IDTMC.

If we want to emphasize that a DTMC $\mathcal{D}$ satisfies the IDTMC $\mathcal{I}$ under both semantics, we write $\mathcal{D} \models \mathcal{I}$. Furthermore, if two DTMCs $\mathcal{D}_1$ and $\mathcal{D}_2$ satisfy a given IDTMC $\mathcal{I}$, we write $\mathcal{D}_1, \mathcal{D}_2 \models \mathcal{I}$ for short.

4.1 Expected Visiting Times in IDTMCs

As, we have discussed , in the previous subsection **??**, IDTMCs can be seen as a collection of DTMCs that satisfy the interval constraints defined by the IDTMC. As only the satisfying DTMCs can be properly interpreted ... When defining EVTs for IDTMCs it is thus useful to define them as bounds for the EVTs of the satisfying DTMCs.

Definition 4.8 (EVTs of IDTMCs)
For a IDTMC $\mathcal{I}$, we define the *upper expected visiting time* (uEVT) $\overline{\mathbb{E}}^{\mathcal{I}}[vt_s]$ and *lower expected visiting time* (lEVT) $\underline{\mathbb{E}}^{\mathcal{I}}[vt_s]$ for all states $s \in S^{\mathcal{I}}$, as follows

(i) $\overline{\mathbb{E}}^{\mathcal{I}}[vt_s] = \mathbb{E}^{\mathcal{D}}[vt_s]$, where $\mathcal{D} \models \mathcal{I}$ and $\mathbb{E}^{\mathcal{D}}[vt_s] \geq \mathbb{E}^{\mathcal{D}'}[vt_s]$ for all $\mathcal{D}' \models \mathcal{I}$.

(ii) $\underline{\mathbb{E}}^{\mathcal{I}}[vt_s] = \mathbb{E}^{\mathcal{D}}[vt_s]$, where $\mathcal{D} \models \mathcal{I}$ and $\mathbb{E}^{\mathcal{D}}[vt_s] \leq \mathbb{E}^{\mathcal{D}'}[vt_s]$ for all $\mathcal{D}' \models \mathcal{I}$.

4.1.1 Relating Expected Visiting Times and Quantitative Reachability

We use the definition of the quantitative reachability property in IDTMCs provided in [BDL+17].

Given a IDTMC $\mathcal{I}$ and a set of states $\Gamma \subseteq S$, then a quantitative reachability property on $\mathcal{I}$ is defined as $Pr^{\mathcal{I}}(\Diamond\Gamma) \sim p$, where $0 \leq p \leq 1$ and $\sim \in \{\leq, <, >, \geq\}$. This quantitative property is satisfied if and only if there exists a DTMC $\mathcal{D} \models \mathcal{I}$ such that $Pr^{\mathcal{D}}(\Diamond\Gamma) \sim p$.

Given the quantitative reachabilities of the DTMCs that satisfy a certain IDTMC $\mathcal{I}$, we can calculate the corresponding lEVTs $\underline{\mathbb{E}}$ and uEVTs $\overline{\mathbb{E}}$ using the results from sec. 3.1.1.

As we concluded in sec. 3.1.1, we can calculate the EVT of a given state s in a DTMC $\mathcal{D}$ via quantitative reachability, using the corresponding DTMC $\mathcal{D}_{uni}$ and the equation $\mathbb{E}^{\mathcal{D}}[vt_s] = \frac{Pr_{s_{in}}^{\mathcal{D}_{uni}}(\Diamond s)}{1 - Pr_s^{\mathcal{D}_{uni}}(\bigcirc\Diamond s)} \cdot \iota_{init}(s_{in})$. Thus, the uEVT and lEVT of a given IDTMC $\mathcal{I}$ are given by the largest and smallest ratio of these two quantitative reachabilities among the satisfying Markov Chains.

However, determining the quantitative reachabilities

4.1.2 Relation between the semantics

Clearly, the IMDP semantic is more expressive than the UMC semantic [BDLM21]. More specifically for any given IDTMC $\mathcal{I}$, a DTMC $\mathcal{D}$ that satisfies $\mathcal{I}$ according to the UMC semantic, i.e. $\mathcal{D} \models_U \mathcal{I}$, does also satisfy $\mathcal{I}$ according to the IMDP semantic, i.e. $\mathcal{D} \models_I \mathcal{I}$ [BDLM21].

Theorem 4.9
For any IDTMC $\mathcal{I}$, if $\mathcal{D} \models_U \mathcal{I}$, then $\mathcal{D} \models_I \mathcal{I}$

For this, we provide a slightly more detailed proof of [BDLM21] (Remark 1).

Proof 9
If $\mathcal{D} \models_U \mathcal{I}$, then we define $T = S$ and $\tau(s) = s$. Then (i) of def. 4.5 is fulfilled as τ is bijective and $\iota_{init}(s) \in [\underline{\iota}_{init}(s), \overline{\iota}_{init}(s)]$ for all $s \in S$, as $\mathcal{D} \models_U \mathcal{I}$. Moreover, (ii) is fulfilled as $\mathbf{P}(s,t) \in [\underline{\mathbf{P}}(s,t), \overline{\mathbf{P}}(s,t)]$ for all $s,t \in S$, as $\mathcal{D} \models_U \mathcal{I}$. Thus, we can conclude that $\mathcal{D} \models_I \mathcal{I}$. □

Importantly, the two semantics have the same maximum and minimum reachability probability for any given (initial) state, according to [BDL+17] and [CHK13]. Moreover, computing the maximum or minimum reachability probability is shown to be P-complete for the IMDP semantic, and thus also for the UMC semantic, in [CHK13].

(Perhaps write about impact: i.e. if algo through reach probs then EVT in PTIME)

4.2 Calculating Expected Visiting Times of IDTMCs

We will now have a look at how the *expected visiting times* for IDTMCs can be calculated

The most intuitive upper and lower bounds for the uEVTs and lEVTs of IDTMCs can be derived by using eq. (3.1) and the upper and lower transition probability functions and initial distributions defined in def. 4.1.

Theorem 4.10 (Upper and Lower Bounds for uEVTs and lEVTs)
For any IDTMC $\mathcal{I}$, the uEVT $\overline{\mathbb{E}}[vt_s]$ is bounded from above by $\mathbb{E}_{upp}[vt_s] = \overline{\iota}_{init}(s) + \sum_{t \in S_{tr}} \overline{\mathbf{P}}(t,s)^\mathsf{T} \cdot \mathbb{E}_{upp}[vt_t]$ and the lEVT $\underline{\mathbb{E}}[vt_s]$ is bounded from below by $\mathbb{E}_{low}[vt_s] = \underline{\iota}_{init}(s) + \sum_{t \in S_{tr}} \underline{\mathbf{P}}(t,s)^\mathsf{T} \cdot \mathbb{E}_{low}[vt_t]$

Proof 10
For any IDTMC $\mathcal{I}$ there exists a DTMC $\mathcal{D}$ that satisfies $\mathcal{I}$, i.e., $\mathcal{D} \models \mathcal{I}$, and $\mathbb{E}^{\mathcal{D}}[vt_s] = \overline{\mathbb{E}}^{\mathcal{I}}[vt_s]$. As $\mathcal{D} \models \mathcal{I}$, it follows that $\mathbf{P}(s,s') \leq \overline{\mathbf{P}}(s,s')$ for all $s,s' \in S$.

Thus, $\mathbb{E}^{\mathcal{D}}[vt_s] = \iota_{init}(s) + \sum_{t \in S_{tr}} \mathbf{P}(t,s)^\mathsf{T} \cdot \mathbb{E}[vt_t] \leq \overline{\iota}_{init}(s) + \sum_{t \in S_{tr}} \overline{\mathbf{P}}(t,s)^\mathsf{T} \cdot \mathbb{E}_{upp}[vt_t] = \mathbb{E}^{\mathcal{I}}_{upp}[vt_s]$.

The identical argument can be made to prove the lower bound for the lEVT. □

While these bounds can be computed as easily as eq. (3.1), i.e., in PTIME, they are not very tight.

Example 4.11
For this take a look at the upper bound of the IDTMC provided in fig. 4.4. Using theorem 4.10, we can calculate that $\mathbb{E}_{upp}[vt_{s_4}] = 2$, as $\mathbb{E}_{upp}[vt_{s_1}] = \mathbb{E}_{upp}[vt_{s_2}] = \mathbb{E}_{upp}[vt_{s_3}] = 1$. However, as the IDTMC is acyclic, see def. 3.15, every state can be visited at most once. Thus, the upper bound $\mathbb{E}_{upp}[vt_{s_4}]$ is clearly not tight. Similarly, $\mathbb{E}_{low}[vt_{s_4}] = 0$, however s_4 needs to be visited at least once. Thus, this lower bound $\mathbb{E}_{low}[vt_{s_4}]$ is not tight either.




Figure 4.4: An example of a DTMC.

In the following, we will therefore take a look at how we can characterize the precise values of the uEVTs and lEVTs.

4.2.1 UMC semantic

In this section we examine how the precise values of the uEVTs and lEVTs can be calculated for IDTMCs under the UMC semantic.

As defined, there can be multiple DTMCs $\mathcal{D}_i$ that satisfy an IDTMC $\mathcal{I}$ with differing transition probability functions $\mathbf{P}^{\mathcal{D}_i}$ and initial distributions $\iota_{init}^{\mathcal{D}_i}$. To determine the uEVT and lEVT for a given state $s \in S^{\mathcal{I}}$, we thus have to find the satisfying DTMCs that maximize and minimze eq. (3.1), respectively. In other words, we need to solve the equations

$$\mathbb{E}[vt_s] = \text{opt}_{\mathcal{D} \models \mathcal{I}} \left(\iota_{init}(s) + \sum_{t \in S_{tr}} \mathbf{P}^{\mathcal{D}}(t, s) \cdot \mathbb{E}[vt_t] \right), \qquad \text{opt} \in \{\min, \max\}.$$

For this, we define two NLPs.

Theorem 4.12 (NLPs for uEVT and lEVT)
For any IDTMC $\mathcal{I}$, the uEVTs and lEVTs of all states $s \in S^{\mathcal{I}}$ can be calculated

as follows

$$
\begin{aligned}
\text{opt} \quad & \iota_{init}(s) + \sum_{t \in S_{tr}} \mathbf{P}^{\mathcal{D}}(t,s) \cdot \mathbb{E}[vt_t] && \text{opt} \in \{\min, \max\} && \\
\text{s.t.} \quad & \sum_{s_j \in N(s)} \mathbf{P}(s_i, s_j) = 1 && \forall s_i \in S_{tr} && (1) \\
& \sum_{s_i \in S} \iota_{init}(s_i) = 1 && && (2) \\
& \underline{\mathbf{P}}^{\mathcal{I}}(s_i, s_j) \leq \mathbf{P}(s_i, s_j) \leq \overline{\mathbf{P}}^{\mathcal{I}}(s_i, s_j) && \forall s_i, s_j \in S_{tr}^{\mathcal{I}} && (3) \\
& \underline{\iota}_{init}^{\mathcal{I}}(s_i) \leq \iota_{init}(s_i) \leq \overline{\iota}_{init}^{\mathcal{I}}(s_i) && \forall s_i \in S_{tr}^{\mathcal{I}} && (4)
\end{aligned}
$$

Values for S_{re} chosen arbitrarily but correctly (bounds)

As the objective function is recursive, we will now briefly prove that the chain (?) is finite. For this, we will use LU-decomposition, see def. 2.12. Thus, we can rewrite

$$
\begin{aligned}
\mathcal{E} &= \iota_{init} + Q^{\mathsf{T}} \cdot \mathcal{E} = \iota_{init} + (LU)^{\mathsf{T}} \cdot \mathcal{E} \\
&= \iota_{init} + U^{\mathsf{T}} \cdot L^{\mathsf{T}} \cdot \mathcal{E}
\end{aligned}
$$

Using the triangular property of U and L, we can now show, that the computation is finite.

Proof 11
Since L^{T} is an upper triangular matrix, we can define $(L^{\mathsf{T}}e)_i = \sum_{i=j}^{n} (L^{\mathsf{T}})_i j \cdot e_j = e_i + \sum_{j=i+1} (L^{\mathsf{T}})_{ij} \cdot e_j$ (making use of $(L_{ii}) = 0$, as L is unitriangular).

And thus since U^{T} is a lower triangular matrix, $e_i = \iota_{init}(s_i) + \sum_{j=1}^{i} (U^{\mathsf{T}})_{ij} \cdot (L^{\mathsf{T}}e)_j = \iota_{init}(s_i) + \sum_{j=1}^{i} (U^{\mathsf{T}})_{ij} \cdot (e_j + \sum_{k=j+1}^{n} (L^{\mathsf{T}})_{kj} \cdot e_k)$. □

In the rest of this thesis, we will use the shorter vector notation $\mathbb{E}[vt_s] = \iota_{init}(s) + (Q)_s \cdot \mathcal{E}$ NOTICE THAT ONLY S_{tr} USED (as only those necessary for calc.) –¿ Others irrelevant

We will now prove the optimality of these NLPs.

Proof 12
We will have a look at the calculation of the uEVTs through ILPs first. First, we will prove "correctness" (??) (Continue with optimality). For this, let x be a given valid (not necessarily optimal) solution of the ILP for the uEVT of a state s for a IDTMC $\mathcal{I}$ and $\mathcal{D}$ a DTMC that satisfies $\mathcal{I}$, i.e. $\mathcal{D} \models \mathcal{I}$, where $\mathbf{P}^{\mathcal{D}}(s, s') = q_{s,s'}$ and $\iota_{init}^{\mathcal{D}}(s) = \iota(s)$ for all $s, s' \in S_{tr}$.

Thus, if x is a solution, then $\sum_{\hat{s} \in N(s')} \mathbf{P}(s', \hat{s}) = 1$ for all $s' \in S_{tr}$.

Moreover, $\underline{\mathbf{P}}^{\mathcal{I}}(s', \hat{s}) \leq \mathbf{P}^{\mathcal{D}}(s, s') \leq \overline{\mathbf{P}}^{\mathcal{I}}(s', \hat{s})$ and $\underline{\iota}^{\mathcal{I}}_{init}(s') \leq \iota^{\mathcal{D}}_{init} \leq \overline{\iota}^{\mathcal{I}}_{init}(s')$ for all transitive states $s', \hat{s} \in S_{tr}$ and by definition also for all recurrent states S_{re}.

...

Definition 4.13 (Trace of a matrix)
For a given matrix M, its trace is defined as $tr(M) = \sum_i (M)_{ii} = \sum_i \lambda_i$, where λ_i denote the eigenvalues of M

Definition 4.14 (Nonzero eigenvalues)
For a given matrix M, if $M \neq 0$, then M has at least one nonzero eigenvalue

As shown before, the EVTs of a DTMC $\mathcal{D}$ can be characterized by $e = (I - Q^{\mathsf{T}})^{-1} \cdot \iota_{init}$. Let us know consider the subclass of this problem, where $Q^3 = 0$ (and $\iota_{init}(s) > 0$ for at least one $s \in S_{tr}$). For this subclass it is necessarily true, that $Q^k = 0$, for all $k \geq 3$. As $\lim_{k\to\infty} Q^k = 0$ for all Q of our problem (as Q is strictly row-substochastic and trivially especially for this subclass), we know that $(I - Q^{\mathsf{T}})^{-1} = (\sum_{k=0}^{\infty} Q^k)^{\mathsf{T}}$. Thus, for our subclass $(I - Q^{\mathsf{T}})^{-1} = (I + Q + Q^2)$ and therefore, $e = (I + Q + (Q^{\mathsf{T}})^2) \cdot \iota_{init} = (I + Q^{\mathsf{T}} + (Q^2)^{\mathsf{T}}) \cdot \iota_{init}$.

Thus, $e = \iota_{init} + \begin{bmatrix} \sum_{j=0}^{n} q_{j1} \cdot \iota_{init}(j) \\ \vdots \\ \sum_{j=0}^{n} q_{jn} \cdot \iota_{init}(j) \end{bmatrix} + \begin{bmatrix} \sum_{j=0}^{n} (\sum_{k=0}^{n} q_{jk} q_{k1}) \iota_{init}(j) \\ \vdots \\ \sum_{j=0}^{n} (\sum_{k=0}^{n} q_{jk} q_{kn}) \iota_{init}(j) \end{bmatrix}$.

Or for a single state:

$$e_t = \iota_{init}(t) + \sum_{j=0}^{n} q_{j\hat{t}} \cdot \iota_{init}(j) + \sum_{j=0}^{n} (\sum_{k=0}^{n} q_{jk} q_{k\hat{t}}) \iota_{init}(j) \ . \tag{4.1}$$

To determine the uEVTs and lEVTs, we are trying to find the maximum and minimum for this equation, that satisfies the constraints of the IDTMC $\mathcal{I}$.
(where $\hat{t}$ gives the index of state t in the ordering)

Definition 4.15 (NP-hard quadratic programming (QP))
A quadratic programming problem with n variables and m constraints can be formulated as:

$$\text{minimize } x^{\mathsf{T}} M x + c^{\mathsf{T}} x + d$$

$$\text{s.t. } Ax \leq b \ ,$$

where $c, x \in \mathbb{R}^n$, $M \in \mathbb{R}^{n \times n}$ and symmetric, $A \in \mathbb{R}^{m \times n}$, $b \in \mathbb{R}^m$, and $d \in \mathbb{R}$.

The quadratic programming problem where M is not positive definite, i.e. M has a negative eigenvalue [PV91], is NP-hard

Thus, we can now rewrite eq. (4.1) to fit such a quadratic programming problem.

For this, we further specify our subclass to only include IDTMCs with a fixed initial distribution for the transient states, i.e. $\underline{\iota}_{init}(s) = \iota_{init}(s) = \overline{\iota}_{init}(s)$ for all $s \in S_{tr}$. We can then define $d = \iota_{init}(t)$ and $x = (q_{00}, q_{01}, \ldots, q_{0n}, q_{10}, \ldots, q_{nn})^\mathsf{T}$. Furthermore, we then accordingly define $M_{p,q} = \frac{1}{2}\iota_{init}(j)$, where $p = j(n+1)+k$ and $q = k(n+1)+\hat{t}$, for all $j, k \in \underline{n}$ and all other entries as zero. Lastly, we define $c_{j(n+1)+\hat{t}} = \iota_{init}(j)$ and all other entries as zero.

We will now continue by briefly showing, that this formulation is correct, i.e. $x^\mathsf{T}Mx + c^\mathsf{T}x + d = \iota_{init}(t) + \sum_{j=0}^{n} q_{j\hat{t}} \cdot \iota_{init}(j) + \sum_{j=0}^{n}(\sum_{k=0}^{n} q_{jk}q_{k\hat{t}})\iota_{init}(j)$. Firstly, trivially $d = \iota_{init}(t)$, (see its definition). Continuing, $c^\mathsf{T}x = \sum_{j=0}^{n} c_{j(n+1)+\hat{t}} \cdot x_{j(n+1)+\hat{t}}$, as all other entries of c are defined as zero. As $c_{j(n+1)+\hat{t}} = \iota_{init}(j)$ and $x_{j(n+1)+\hat{t}} = q_{j\hat{t}}$, thus $c^\mathsf{T}x = \sum_{j=0}^{n} q_{j\hat{t}} \cdot \iota_{init}(j)$. Lastly, $x^\mathsf{T}Mx = \sum_{pq} M_{pq}x_p x_q$. When substituing the nonzero entries of M, where $p = j(n+1)+k$ and $q = k(n+1)+\hat{t}$, we get $\sum_{j,k} M_{j(n+1)+k,k(n+1)+\hat{t}} \cdot x_{j(n+1)+k} \cdot x_{k(n+1)+\hat{t}} + \sum_{j,k} M_{k(n+1)+\hat{t},j(n+1)+k} \cdot x_{k(n+1)+\hat{t}} \cdot x_{j(n+1)+k}$. As M is symmetrical and $M_{pq} = \iota_{init}(j)$ for these nonzero entries, we can simplify to: $\sum_{j,k} \iota_{init}(j) \cdot x_{j(n+1)+k} \cdot x_{k(n+1)+\hat{t}}$. Now, finally substituing back $x_{j(n+1)+k} = q_{jk}$ and $x_{k(n+1)+\hat{t}} = q_{k\hat{t}}$ gets us: $x^\mathsf{T}Mx = \sum_{j,k} \iota_{init}(j) q_{jk}q_{k\hat{t}} = \sum_{j=0}^{n}(\sum_{k=0}^{n} q_{jk}q_{k\hat{t}})\iota_{init}(j)$. This objective is then, similar to theorem 4.12, subject to the constraints $\underline{\mathbf{P}}^{\mathcal{I}}(s', s'') \leq q_{s's''} \leq \overline{\mathbf{P}}^{\mathcal{I}}(s', s'')$ for all $s', s'' \in S^{\mathcal{I}}_{tr}$. The soundness and completeness of this QP are identical to the NLP (and will thus be omitted). Thus, this subclass of the problem is NP-hard if and only if M is not positive definite. For this we will further specify our subclass to only include IDTMCs $\mathcal{I}$ with $\underline{\iota}_{init}(t) = \overline{\iota}_{init}(t) = 0$, where t is the state for which the uEVT and lEVT are calculated. With how we defined M, the entries $(M)_{ii}$ are nonzero if and only if $j = k = \hat{t}$. Therefore, (M_{ii}) is nonzero if and only if $\iota_{init}(t) > 0$ and thus for the IDTMCs of our subclass $tr(M) = 0$. Furthermore, as $M \neq 0$ (trivially as we have defined for our subclass that $\iota_{init}(s)$ for at least one $s \in S_{tr}$), we can use def. 4.14 to reason that M must have at least one nonzero eigenvalue. Moreover, using def. 4.13, we know that $\sum_i \lambda_i = tr(M) = 0$, necessarily M must have a negative eigenvalue. Thus, we can reason that according to [PV91] this subclass belongs to a class of problems that is NP-hard. This would suggest that this problem is rather hard.

Thus after

Definition 4.16 (Intermediate state)
Given a DTMC $\mathcal{D}$, a set of states T, and a (goal) state $t \in S^{\mathcal{D}}$, then $u \in S^{\mathcal{D}}$ is intermediate if and only if there exists path fragment $\tilde{\pi}$ with $|\tilde{\pi}| = n$, see $\ldots$, such that $\tilde{\pi}[0] \in T, \tilde{\pi}[n] = s$ and $\tilde{\pi}[k] = u$ for some $k \in \underline{n}$. In other words there exists a path fragment from some $t \in T$ to s in $\mathcal{D}$ that contains u.

We denote the set of these intermediate states as $I^{\mathcal{D}}_{Ts}$ and I_{Ts} if the DTMC is clear from the context. Moreover, if T is a singleton set, i.e., $T = \{t\}$, then we simply denote the set of intermediate states as I_{ts}.

Lemma 4.17 (Recursive Formula for Reachability Probability)
UMC semantic:

For states $t, s \in S$:

$$Pr_t(\Diamond s) = \sum_{u \in N(t) \cap I_{ts}} \mathbf{P}(t,u) \cdot Pr_u(\Diamond s)$$

Proof 13
According to def. 3.9,

$$\begin{aligned}
Pr_t^{\mathcal{D}}(\Diamond s) &= \sum_{\pi \in \Diamond_t s} \prod_{k=0}^{|\pi|-1} \mathbf{P}^{\mathcal{D}}(\pi[k], \pi[k+1]) \\
&= \sum_{\pi \in \Diamond_t s} \mathbf{P}^{\mathcal{D}}(\pi[0], \pi[1]) \cdot \prod_{k=1}^{|\pi|-1} \mathbf{P}^{\mathcal{D}}(\pi[k], \pi[k+1]) \qquad (\pi[0] = t \text{ and } \pi[1] \in N(t)) \\
&= \sum_{u \in N(t)} \underbrace{\mathbf{P}(t,u)}_{=0, \text{ for all } u \notin I_{ts}^{\mathcal{D}}} \cdot \sum_{\pi \in \Diamond_u s} \prod_{k=1}^{|\pi|-1} \mathbf{P}^{\mathcal{D}}(\pi[k], \pi[k+1]) \\
&= \sum_{u \in N(t) \cap I_{ts}^{\mathcal{D}}} \mathbf{P}(t,u) \cdot \sum_{\pi \in \Diamond_u s} \prod_{k=1}^{|\pi|-1} \mathbf{P}^{\mathcal{D}}(\pi[k], \pi[k+1]) \\
&= \sum_{u \in N(t) \cap I_{ts}^{\mathcal{D}}} \mathbf{P}(t,u) \cdot Pr_u(\Diamond s)
\end{aligned}$$

□

Remark. *Existence of DTMC with optimal reachability.* For every transient state $t \in S_{tr}$ there exists a DTMC $\mathcal{D} \models_U \mathcal{I}$ with $Pr_t^{\mathcal{D}}(\Diamond s) = \sup_{\mathcal{D}_i \models_U \mathcal{I}} Pr_t^{\mathcal{D}_i}(\Diamond s)$. This follows directly from the compactness and continuity of the polytope of DTMCs satisfying $\mathcal{I}$; it is not reproved here, only reused. To shorten the notation we will denote $\dot{Pr}_t^{\mathcal{D}}(\Diamond s) = \sup_{\mathcal{D}_i \models_U \mathcal{I}} Pr_t^{\mathcal{D}_i}(\Diamond s)$

Theorem 4.18
Given an IDTMC $\mathcal{I}$, a transient state $t \in S^{\mathcal{I}}$, a (goal) state $s \in S^{\mathcal{I}}$, and a DTMCs $\mathcal{D}$ with $\mathcal{D} \models \mathcal{I}$. (Positive transition/reachability probability from t to s, i.e. there needs to exist a path fragment from t to s)

$\mathcal{D}$ has the maximum reachability probability from t to s, i.e., $Pr_t^{\mathcal{D}}(\Diamond s) = \sup_{\mathcal{D}_i \models \mathcal{I}} Pr_t^{\mathcal{D}_i}(\Diamond s)$,

if and only for all $u \in I_{ts}^{\mathcal{D}}$:

$$Pr_u^{\mathcal{D}}(\diamond s) = \sup_{\mathcal{D}_i \models \mathcal{I}} Pr_u^{\mathcal{D}_i}(\diamond s)$$

Proof 14
We will start by proving this theorem for $\mathcal{D} \models_I \mathcal{I}$ and then show why $\mathcal{D} \models_U \mathcal{I}$ follows from it.

As shown in [BK08] (Theorem 10.100, pg. 851), for any MDP $\mathcal{M}$ the maximum reachability probability is characterized by the equation

$$Pr_t^{\max}(\diamond s) = \max \left\{ \sum_{u \in N(t)} \mathbf{P}(t, \alpha, u) \cdot Pr_u^{\max}(\diamond s) \mid \alpha \in Act(t) \right\} . \tag{4.2}$$

□

Clearly, the policies which solve this equation have the maximum reachability probability $Pr_u(\diamond s)$ for all $u \in I_{ts}^{\mathcal{D}}$, as shown in [BK08]. As shown (not yet but later), an IDTMC under the IMDP semantic can be mapped to a MDP, where the policy represents a given satisfying DTMC (bzw. nach einer/der üblichen IMDP Semantiken ist eine policy/adversary das satisfying DTMC). Thus, this theorem is proven for the IMDP semantic. (Also importantly, the solution to the set of equations is unique, as shown in [BK08]. Thus, the maximum reachability probabilities of the states are unique)

As mentioned in sec. 4.1.2, [BDL+17] and [CHK13] prove that the maximum (& minimum) reachability probability are identical under the UMC and IMDP semantic. Let $\mathcal{D}_U \models_U \mathcal{I}$ and $\mathcal{D}_I \models_I \mathcal{I}$, be two DTMCs with maximum reachability probability ($Pr_t(\diamond s)$). As discussed in sec. 4.1.2, every DTMC that satisfies an IDTMC under the UMC semantic, also satisfies it under the IMDP semantic. Thus, $\mathcal{D}_U \models_I \mathcal{I}$ and because it has the maximum reachability probability $Pr_t(\diamond s)$, it clearly satisfies eq. (4.2) (i.e. is a solution/policy for it). Thus, this theorem is proven for the UMC semantic.

Corollary 4.19
Given an IDTMC $\mathcal{I}$, two transient state $t_1, t_2 \in S^{\mathcal{I}}$, a (goal) state $s \in S^{\mathcal{I}}$, and two DTMCs $\mathcal{D}_1, \mathcal{D}_2$ with $\mathcal{D}_1, \mathcal{D}_2 \models_U \mathcal{I}$ and $Pr_{t_1}^{\mathcal{D}_1}(\diamond s) = \sup_{\mathcal{D} \models_U \mathcal{I}} Pr_{t_1}^{\mathcal{D}}(\diamond s)$ and $Pr_{t_2}^{\mathcal{D}_2}(\diamond s) = \sup_{\mathcal{D} \models_U \mathcal{I}} Pr_{t_2}^{\mathcal{D}}(\diamond s)$. In other words the DTMCs have a maximum reachability probability for t_1 and t_2, respectively.

For all $u \in I_{t_1 s}^{\mathcal{D}_1} \cap I_{t_2 s}^{\mathcal{D}_2}$, the reachability probabilities $Pr_u^{\mathcal{D}_1}(\diamond s) = Pr_u^{\mathcal{D}_2}(\diamond s)$ are identical.

Proof 15
This theorem follows from theorem 4.18. Using theorem 4.18, we can conclude that $Pr_u^{\mathcal{D}_1}(\Diamond s) = \sup_{\mathcal{D}_i \models_U \mathcal{I}} Pr_u^{\mathcal{D}_i}(\Diamond s)$ for all $u \in I_{t_1 s}^{\mathcal{D}_1}$ and $Pr_u^{\mathcal{D}_2}(\Diamond s) = \sup_{\mathcal{D}_i \models_U \mathcal{I}} Pr_u^{\mathcal{D}_i}(\Diamond s)$ for all $u \in I_{t_2 s}^{\mathcal{D}_2}$. Thus, $Pr_u^{\mathcal{D}_1}(\Diamond s) = Pr_u^{\mathcal{D}_2}(\Diamond s)$ for all $u \in I_{t_1 s}^{\mathcal{D}_1} \cap I_{t_2 s}^{\mathcal{D}_2}$. □

Theorem 4.20 (One-state Extension)
Given a set $T \subset S_{tr}$, a transient state $t \notin T$, and a (goal) state s, assume that $\mathcal{D}_T \models_U \mathcal{I}$ with $\overline{Pr}_u^{\mathcal{I}}(\Diamond s) = Pr_u^{\mathcal{D}_T}(\Diamond s)$ for all $u \in T$ and $\mathcal{D}_t \models_U \mathcal{I}$ with $\overline{Pr}_t^{\mathcal{I}}(\Diamond s) = Pr_t^{\mathcal{D}_t}(\Diamond s)$. Define $\mathcal{D}'$ by

$$\mathbf{P}^{\mathcal{D}_{cmb}}(w,v) = \begin{cases} \mathbf{P}^{\mathcal{D}_t}(w,v) & \text{if } w \in I_{ts}^{\mathcal{D}_t} \backslash I_{Ts}^{\mathcal{D}_T} \\ \mathbf{P}^{\mathcal{D}_T}(w,v) & \text{else.} \end{cases}$$

Then, $\overline{Pr}_u^{\mathcal{D}_{cmb}}(\Diamond s) = Pr_u^{\mathcal{D}_{cmb}}(\Diamond s)$ for all $u \in T \cup \{t\}$, i.e., $\mathcal{D}_{cmb}$ has the maximum reachability probability for all states in $T \cup \{t\}$.

Proof 16
As $\mathbf{P}^{\mathcal{D}_{cmb}}(w,v) = \mathbf{P}^{\mathcal{D}_T}(w,v)$ for all $w \in I_{Ts}^{\mathcal{D}_T}$, clearly $\overline{Pr}_u^{\mathcal{I}}(\Diamond s) = Pr_u^{\mathcal{D}_{cmb}}(\Diamond s)$ for all $u \in T$, see lemma 4.17.

Thus, we need to show that $Pr_t^{\mathcal{D}_{cmb}}(\Diamond s) = \overline{Pr}_t^{\mathcal{I}}(\Diamond s)$. As we know that $Pr_t^{\mathcal{D}_t}(\Diamond s) = \overline{Pr}_t^{\mathcal{I}}(\Diamond s)$, we will prove that $Pr_t^{\mathcal{D}_{cmb}}(\Diamond s) = Pr_t^{\mathcal{D}_t}(\Diamond s)$. Notice, that if $I_{ts}^{\mathcal{D}_t} \cap I_{Ts}^{\mathcal{D}_T} = \emptyset$, then the proof concludes using the same argument as for $u \in T$. Furthermore, if $I_{ts}^{\mathcal{D}_t} \subseteq I_{Ts}^{\mathcal{D}_T}$, then the proof concludes directly using theorem 4.18.

Just use previous argument

This assignment might have conflictions for $u, w \in I_{t_1 s}^{\mathcal{D}_1} \cap I_{t_2 s}^{\mathcal{D}_2}$. For these conflictions we assign the value of $\mathbf{P}^{\mathcal{D}_1}(u,w)$. Clearly, $\mathcal{D}_{cmb}$ has the maximum reachability probability for $Pr_{t_1}(\Diamond s)$. (1) Recall, that $Pr_{t_2}^{\mathcal{D}_{cmb}}(\Diamond s) = \sum_{u \in N(t)} \mathbf{P}^{\mathcal{D}_{cmb}}(t_2,u) \cdot Pr_u^{\mathcal{D}_{cmb}}(\Diamond s)$, see lemma 4.17. For all $u \in I_{t_2 s}^{\mathcal{D}_2} \backslash I_{t_1 s}^{\mathcal{D}_1}$, $\mathbf{P}^{\mathcal{D}_{cmb}}(u,w) = \mathbf{P}^{\mathcal{D}_2}(u,w)$ and for all $u \in I_{t_1 s}^{\mathcal{D}_1} \cap I_{t_2 s}^{\mathcal{D}_2}$, $Pr_u^{\mathcal{D}_{cmb}}(\Diamond s) = Pr_u^{\mathcal{D}_2}(\Diamond s)$. From this, we can conclude that $Pr_{t_2}^{\mathcal{D}_2}(\Diamond s) = Pr_{t_2}^{\mathcal{D}_{cmb}}(\Diamond s)$.

(2) As, $\mathcal{D}_1$ has maximum reachability probability for all states $u \in I_{t_1 s}^{\mathcal{D}_1}$, see theorem 4.18, the states in $I_{t_1 s}^{\mathcal{D}_1} \cap I_{t_2 s}^{\mathcal{D}_2}$ have maximum reachability probability in $\mathcal{D}_{cmb}$. Thus, all states in $I_{t_2 s}^{\mathcal{D}_2}$ have maximum reachability probability in $\mathcal{D}_{cmb}$. As, $\mathbf{P}^{\mathcal{D}_{cmb}}(t_2,u) = \mathbf{P}^{\mathcal{D}_2}(t_2,u)$ for all $u \in I_{t_2 s}^{\mathcal{D}_2}$, we can conclude that $\mathcal{D}_{cmb}$ also has the maximum reachability probability for $Pr_{t_2}(\Diamond s)$.

Is it understandabale enough why this is true?? (Why we do not have to argue for correct choosing of $\mathbf{P}$, i.e., why only max reach prob for shared intermediate states is sufficient)?

Pick one of the two current explanations: (1) or (2) (or combine somehow) □

Crucially, this theorem can be extended to all transient states.

Theorem 4.21
Given an IDTMC $\mathcal{I}$ and a (goal) state $s \in S$ there exists a DTMC $\mathcal{D} \models_U \mathcal{I}$ such that $Pr_t^{\mathcal{D}}(\Diamond s) = \sup_{\mathcal{D}_i \models_U \mathcal{I}} Pr_t^{\mathcal{D}_i}(\Diamond s)$ for all transient states $t \in S_{tr}$.

Proof 17
Consider any given enumeration $t_1, t_2, \ldots$ of S_{tr}. Clearly this enumeration is finite, since S_{tr} is finite, see sec. 3.2.

Construction. Now, let $D^{(1)} \models_U \mathcal{I}$ with $Pr_{t_1}^{\mathcal{D}^{(1)}}(\Diamond s) = \overline{Pr}_{t_1}^{\mathcal{I}}$ □

Konstruktion: jeweils zusammengefasste init states und direct successors von s. –¿ zeigen, dass das die selben EVTs hat (Bijektion für DTMCs). Zeigen das IDTMC Konstruktion nach lemma DTMC mit max EVT hat (beide max reach probs) (vllt reicht es auch zu zeigen dass es für diesen DTMC ein Äquivalent im ursprünglichen IDTMC gibt das selbe EVTs hat (für s halt))

+ Zwei state Konstruktion schnelle Möglichkeit zugrunde liegende Struktur zu berechnen (einfach geaddete states in transitionen übersetzen)

4.2.2 IMDP semantic

In this subsection we will look at how uEVTs and lEVTs can be calculated for the IMDP semantic.

Recall, that we concluded in sec. 4.1.1 that the uEVT and lEVT of a given IDTMC $\mathcal{I}$ can be calculated by maximizing or minimizing the term $\sum_{t \in S_{tr} \cap in} \frac{Pr_t(\Diamond s)}{1 - Pr_s(\bigcirc \Diamond s)} \cdot \iota_{init}(t)$ for all DTMCs $\mathcal{D} \models \mathcal{I}$. Intuitively, we can thus find the lEVTs and uEVTs by maximizing or minimizing the reachability probability from the initial state t to the given state s ($Pr_t(\Diamond s)$) and from s back to itself ($Pr_s(\bigcirc \Diamond s)$). As shown in [CHK13] the problem of finding this supremum or infimum for a given IDTMC is P-complete. Thus, we can derive the two suprema or infima belonging to (at most) two different satisfying DTMCs $\mathcal{D}_{s_{in} \to s}$ and $\mathcal{D}_{s \to s}$ ($\mathcal{D}_{s_{in} \to s}$, $\mathcal{D}_{s \to s} \models_I \mathcal{I}$).

As we have defined the IMDP semantic using adversarial behaviour (?), notice that we can define a policy that choses the probability distribution specified by $\mathcal{D}_{s_{in} \to s}$ until state s is visited for the first time and then choses the probability distribution specified by $\mathcal{D}_{s \to s}$. This policy can be characterized as having two stages where we transition

from the first to the second stage when passing through s for the first time. Notice however, that the stochastic process of the Markov Chain/MDP (?) would no longer be memoryless, i.e. the Markov Property (SEE ...) would break, as the process is not only dependent on the current state but also on what stage the policy is in. To solve this problem, we propose a remapping/reduction (state augmentation).

Construction. Given an IDTMC $\mathcal{I} = (S, \underline{\mathbf{P}}, \overline{\mathbf{P}}, \underline{\iota}_{init}, \overline{\iota}_{init})$, the previously defined $\mathcal{D}_{s_{in}\to s}$, $\mathcal{D}_{s\to s}$ (corresponding to either the suprema or infima) and a state s (for which we want to calculate the EVT), we define $\mathcal{D}^s_{aug}$, with $S^{\mathcal{D}^s_{aug}} = S_{st1} \cup S_{st2}$, where the state set for stage 1 is $S_{st1} = \{s^{(1)} \mid s \in S^{\mathcal{I}}_{tr}\}$ (transient states in $\mathcal{I}$) and the state set for stage 2 is $S_{st2} = \{s^{(2)} \mid s \in S^{\mathcal{I}}\}$. Furthermore, $\mathbf{P}^{\mathcal{D}^s_{aug}}(s_1^{(1)}, s_2^{(1)}) = \mathbf{P}^{\mathcal{D}_{s_{in}\to s}}(s_1, s_2)$ for all $s_1^{(1)} \in S_{st1}$, $s_2^{(1)} \in S_{st1}\backslash\{s\}$ and $\mathbf{P}^{\mathcal{D}^s_{aug}}(s_1^{(1)}, s_2^{(2)}) = \mathbf{P}^{\mathcal{D}_{s_{in}\to s}}(s_1, s_2)$ for $s_1^{(1)} \in S_{st1}$, $s_2^{(2)} \in S_{st2} \cap S^{\mathcal{I}}_{re}$. Similarily, $\mathbf{P}^{\mathcal{D}^s_{aug}}(s_1^{(2)}, s_2^{(2)}) = \mathbf{P}^{\mathcal{D}_{s\to s}}(s_1, s_2)$ for all $s_1^{(2)}, s_2^{(2)} \in S_{st2}$. Moreover, $\mathbf{P}^{\mathcal{D}^s_{aug}}(s^{(1)}, s_i^{(2)}) = \mathbf{P}^{\mathcal{D}_{s\to s}}(s, s_i)$ for $s^{(1)} \in S_{st1}$ and $s_i \in N^{\mathcal{I}}(s)$. Lastly, $\iota^{\mathcal{D}^s_{aug}}_{init}(t^{(1)}) = \iota^{\mathcal{D}_{s_{in}\to s}}_{init}(t)$ for all $t^{(1)} \in S_{st1}$ and $\iota^{\mathcal{D}^s_{aug}}_{init}(t^{(2)}) = \iota^{\mathcal{D}_{s_{in}\to s}}_{init}(t)$ for all $t^{(2)} \in S_{st2} \cap S_{re}$, while $\iota^{\mathcal{D}^s_{aug}}_{init}(t^{(2)}) = 0$ for all $t^{(2)} \in S_{st2}\backslash S_{re}$.




Figure 4.5: target state in red

Example 4.22
Recall the IDTMC $\mathcal{I}$ in fig. 4.1, displayed again in fig. 4.5. fig. 4.6 shows $\mathcal{D}^{s_4}_{aug}$, i.e., the construction of the maximizing DTMC for the EVT of $\mathcal{I}$.

(We will now show that this remapping / characterization of this history-dependent policy is coincides (?) with def. 4.5.)

Lemma 4.23
Given and IDTMC $\mathcal{I}$ and the corresponding $\mathcal{D}^s_{aug}$, it holds that $\mathcal{D}^s_{aug} \models_I \mathcal{I}$




Figure 4.6: $\tau(t_1) = \tau(t_4) = s_1$, $\tau(t_2) = \tau(t_7) = s_2$, $\tau(t_5) = s_3$, $\tau(t_3) = \tau(t_6) = s_4$. Target state in red.

Proof 18
To prove this lemma, we will show that def. 4.5 holds for $\mathcal{D}^s_{aug}$. For this, we will first define a mapping $\tau : S^{\mathcal{D}^s_{aug}} \to S^{\mathcal{I}}$. For this, we need to account for S_{st1} and S_{st2}. We define $\tau(s_i^{(1)}) = s_i$ for all $s_i^{(1)} \in S_{st1} \cap S_{tr}$ and $\tau(s_i^{(2)}) = s$ for all $s_i^{(2)} \in S_{st2}$. Thus, for all $s \in S^{\mathcal{I}}_{tr}$, there exists precisely one $s^{(1)} \in S_{st1}$ with $\tau(s^{(1)}) = s$ and $\iota^{\mathcal{D}_{aug}}_{init}(s^{(1)}) \in [\underline{\iota}^{\mathcal{I}}_{init}(s), \overline{\iota}^{\mathcal{I}}_{init}(s)]$ and $s^{(2)} \in S_{st2}$ with $\tau(s^{(2)}) = 0$. Moreover, for all $s \in S^{\mathcal{I}}_{re}$, there exists precisely one $s^{(2)} \in S_{st2}$ with $\tau(s^{(2)}) = s$ and $\iota^{\mathcal{D}_{aug}}_{init}(s^{(2)}) \in [\underline{\iota}^{\mathcal{I}}_{init}(s), \overline{\iota}^{\mathcal{I}}_{init}(s)]$ and $s^{(1)} \notin S_{st1}$. Therefore, we can conclude that condition (i) of def. 4.5 holds.

Secondly, condition (ii) clearly holds for all $s_i^{(2)} \in S_{st2}$, as all of its transition are defined to be identical to that of $\mathcal{D}_{s\to s}$ and $\mathcal{D}_{s\to s} \models \mathcal{I}$. Thus, it remains only to show that condition (ii) also holds for all states in S_{st1}. For all states $s_1^{(1)} \in S_{st1} \backslash \{s\}$, $\mathbf{P}^{\mathcal{D}^s_{aug}}(s_1^{(1)}, s_2^{(1)}) = \mathbf{P}^{\mathcal{D}_{s_{in}\to s}}(s_1, s_2)$, for all $s_2 \in S^{\mathcal{D}_{s_{in}\to s}}_{tr}$ and $s_2 \in S^{\mathcal{D}_{s_{in}\to s}}_{re}$. Moreover, for $\mathbf{P}^{\mathcal{D}^s_{aug}}(s^{(1)}, s_i^{(1)}) = 0$ and $\mathbf{P}^{\mathcal{D}^s_{aug}}(s^{(1)}, s_i^{(2)}) = \mathbf{P}^{\mathcal{D}_{s\to s}}(s, s_i)$. As, $\mathcal{D}_{s_{in}\to s}$, $\mathcal{D}_{s\to s} \models \mathcal{I}$, we can thus conclude that condition (ii) also holds for all states in S_{st1}. As both conditions of def. 4.5 are fulfilled, we can thus conclude that $\mathcal{D}^s_{aug} \models_I \mathcal{I}$. □

Construction. For a given DTMC $\mathcal{D}$ and a state $s \in S$, consider the DTMC $\mathcal{D}_{abs} = (S^{\mathcal{D}_{abs}}, \mathbf{P}^{\mathcal{D}_{abs}}, \iota^{\mathcal{D}}_{init})$ with the set of states $S^{\mathcal{D}_{abs}} = S^{\mathcal{D}}_{tr} \cup \{s_{re}\}$. Moreover, the probability

function $\mathbf{P}_{abs}$ is identical to $\mathbf{P}^{\mathcal{D}}$, except the transitions $\mathbf{P}_{abs}(s, s') = 0$ for all $s' \in S$ and $\mathbf{P}_{abs}(s, s) = 1$, as well as $\mathbf{P}^{\mathcal{D}}(s_1, s_2) = \mathbf{P}^{\mathcal{D}_{abs}}(s_1, s_{re})$, for all $s_1 \in S^{\mathcal{D}}_{tr}$, $s_2 \in S^{\mathcal{D}}_{re}$ and further $\mathbf{P}^{\mathcal{D}_{abs}}(s_{re}, s_{re}) = 1$. In other words, $\mathcal{D}_{abs}$ is modified only in such a way that s is absorbing and all recurrent states are summarized into one state.

Lemma 4.24
The reachability probabilities $Pr^{\mathcal{D}}(\Diamond s) = Pr^{\mathcal{D}_{abs}}(\Diamond s)$ are equal.

Proof 19
If the set of initial states S_{init} includes s then trivially the reachability probability is 1 for both $\mathcal{D}$ and $\mathcal{D}_{abs}$.

As $Pr(\Diamond s) = Pr(S \backslash \{s\} \cup s)$, we can reason/conclude using def. 3.5 and def. 3.7 that $Pr(\Diamond s) = \iota^{\mathcal{D}}_{init}(\pi[0]) \cdot \prod_{k=0}^{|\pi|-1} \mathbf{P}^{\mathcal{D}}(\pi[k], \pi[k+1])$. As there is no factor $\mathbf{P}(s, s')$ for any $s' \in S$ in this product, as well as no factor of the form $\mathbf{P}(s_1, s_2)$, where $s_1 \in S^{\mathcal{D}}_{re}$ and the probabilty functions $\mathbf{P}$ and $\mathbf{P}_{abs}$ are identical for all transitions except those, we can conclude that lemma 4.24 is true. □

Thus, we can reason that transitions from the state s itself and recurrent states have no influence on the reachability probability $Pr^{\mathcal{D}}(\Diamond s)$.

Lemma 4.25
Let $Pr^{max}(\Diamond s)$ and $Pr^{max}_s(\bigcirc\Diamond s)$ be the maximum reachability probabilities of $\mathcal{I}$. It holds that $Pr^{D^s_{aug}}(\Diamond s) = Pr^{max}(\Diamond s)$ and $Pr_s^{D^s_{aug}}(\bigcirc\Diamond s) = Pr^{max}_s(\bigcirc\Diamond s)$

Proof 20
First, we will show that $Pr^{D^s_{aug}}(\Diamond s) = Pr^{max}(\Diamond s)$. Notice, that for all $s \in S_{st1} \backslash \{s\}$, the probability function $\mathbf{P}^{\mathcal{D}^s_{aug}}$ is identical to $\mathbf{P}^{\mathcal{I}}$, i.e. it is identical for all transient states except s. Using lemma 4.24, we can thus conclude that $Pr^{D^s_{aug}}(\Diamond s) = Pr^{max}(\Diamond s)$.

Next, we will show that $Pr_s^{D^s_{aug}}(\bigcirc\Diamond s) = Pr^{max}_s(\bigcirc\Diamond s)$. Trivial because (basicaly/literaly) the same DTMC. □

4.2.3 Special Case: Acyclic IDTMCs

Definition 4.26 (Acyclic IDTMC)
An IDTMC $\mathcal{I}$ is acyclic if and only if all satisfying DTMCs $\mathcal{D}$ ($\mathcal{D} \models \mathcal{I}$) are acyclic.

– l/uEVT is simply maximum/minimum reachability (which can be calc in P, cite paper).

4.2.4 Value Iteration

Explain why useful & interesting to define Bellman operator usw. (Runtime advantages, etc.)

In [KTDB19], the authors prove

4.3 Exkurs: aIMCs

Definition 4.27 (Augmented Interval Markov Chains)
An augmented Interval Markov Chain (aIMC) $\mathcal{A}$ is a tuple $(S, \underline{\mathbf{P}}, \overline{\mathbf{P}}, \underline{\iota}_{init}, \overline{\iota}_{init}, C)$, where $(S, \underline{\mathbf{P}}, \overline{\mathbf{P}}, \underline{\iota}_{init}, \overline{\iota}_{init})$ is an IDTMC $\mathcal{I}$ and $C \subseteq S^4$ is a set of *edge equality constraints*. Similar to the once-and-for-all semantic for IDTMCs, see **??**, we define, that an arbitrary DTMC $\mathcal{D}$ satisfies an aIMC $\mathcal{A}$ if it satisfies the corresponding IDTMC $\mathcal{I}$ and $\mathbf{P}^{\mathcal{D}}(s_1, s_2) = \mathbf{P}^{\mathcal{D}}(s_3, s_4)$ for all $(s_1, s_2, s_3, s_4) \in C$. [Cho17]

In other words, aIMCs extend the definition of IMCs with an "edge-constraint" which enforces certain transition probabilities to be equal. A restriction of the quantitative reachability problem, which we discussed in sec. 3.1.1 and sec. 4.1.1, is the qualitative reachability problem, where is restricted to $p \in \{0, 1\}$ (see sec. 4.1.1).

In [Cho17], the authors prove that the qualtitative reachability problem for aIMCs is NP-complete. Thus, as trivially the qualitative problem can be reduced to the quantitative problem, we can thus deduce that the quantitative problem for aIMCs is NP-hard. As we have shown in ...

4.4 Stationary Distributions in IDTMCs

As stated in theorem 3.3 the k-th power of the transition matrik $\mathcal{T}^k$ contains the probabilities of transitioning from any state i to any other state j in precisely k steps. Thus, the product $\iota_{init}^{\mathsf{T}} \cdot \mathcal{T}^k$ characterizes the probability distribution of the current state of the given Markov state in step k. We define this product as $\theta_k^{\mathsf{T}} = \iota_{init}^{\mathsf{T}} \cdot \mathcal{T}^k$, where the i-th entry of θ_k characterizes the probability that the current state in step k is i. We can also define θ_k iteratively through $\theta_k^{\mathsf{T}} = \theta_{k-1}^{\mathsf{T}} \cdot \mathcal{T}$

Definition 4.28 (Stationary Distribution)
A distribution θ_k is stationary if and only if $\theta_k^{\mathsf{T}} = \theta_k^{\mathsf{T}} \cdot \mathcal{T}$, i.e. the probability distribution of the current state remains unchanged through future steps. We denote the stationary distribution as θ, where $\theta(s)$ denotes the entry of θ which corresponds to state s.

In contrast to EVTs, when considering the stationary distribution of a DTMC or IDTMC, we are interested in the recurrent, not transient, states.

Theorem 4.29
If θ is stationary, then $\theta(i) = 0$ if the corresponding state s_i is transient and $\theta(i) > 0$ if s_i is recurrent.

We provide a short proof.

Proof 21
Assume that some s is transient and for the stationary distribution $\theta(s) > 0$. As θ is stationary, there exists some $k \in \mathbb{N}$ such that $\theta_{k'}$ is stationary for $k' \geq k$. Thus, there is a nonzero probability that the current state in step $k' \geq k$ is s. However, as s is transient it follows that $Pr_s(\bigcirc\diamond s) < 1$, see def. 3.12. Therefore, there must be some $k'' > k$ such that $\theta_{k''}(i) = 0$. This contradicts the stationary property of θ. Now, assume that some s is recurrent and for the stationary distribution $\theta(s) = 0$. As s is reachable by definition/construction, see sec. 3.2, we can conclude that $Pr(\diamond s) > 0$. Thus, there exists some $k' \in \mathbb{N}$ such that $\theta_{k'}(i) > 0$. As s_i is recurrent, it follows that $Pr_{s_i}(\bigcirc\diamond s_i) = 1$, see again def. 3.12. Thus, there exist infinitely many $k'' \in \mathbb{N}$ with $k'' > k'$ and $\theta_{k''}(i) > 0$. Therefore, θ_k is not stationary. □

Notice, that any maximal set of connected recurrent states, i.e., every state is connected to some other state in the set, in a DTMC, is a BSCC, see def. 3.11. Firstly, a set of connected recurrent states in a DTMC is strongly connected. By definition, see sec. 3.2, all of the recurrent states in the set are reachable. As the return probability $Pr_{s_{re}}(\bigcirc\diamond S_{re})$ is equal to 1 for any recurrent state s_re, see def. 3.12, any state that is a successor to s_{re} must also be a predecessor, i.e., every edge in this set is bidirectional (in a sense). Thus, trivially strongly connected.

→ It is SCC

Clearly also BSCC

(Bridge: recurrent groups are BSCC & BSCC is irreducible DTMC (perhaps prove this shortly)) We denote stationary distribution of a DTMC $\mathcal{D}$ restricted to a BSCC B as $\theta^{\mathcal{D}|_B}$.

As proven in [Kul16], the stationary distribution $\theta^{\mathcal{D}}$ of the recurrent states of a DTMC $\mathcal{D}$ is characterized by $\theta^{\mathcal{D}}(s) = Pr^{\mathcal{D}}(\diamond B) \cdot \theta^{\mathcal{D}|_B}(s)$ for all $s \in S_{re}$. Moreover, in [MKQW25], it is shown that the stationary distribution within a BSCC can be computed through its EVTs. For this they construct a DTMC $\mathcal{B}\,\hat{\uparrow}\hat{v}$ for every BSCC B with any arbitrary state $v \in S^{\mathcal{D}|_B}$ in which all incoming transitions of v are redirected to the fresh absorbing state $\hat{v}$.

Construction. More precisely, $\mathcal{B}\,\hat{\uparrow}\hat{v} = (S^{\mathcal{D}|_B} \cup \{\hat{v}\}, \mathbf{P}^{\mathcal{B}\hat{\uparrow}\hat{v}}, \iota_{init}^{\mathcal{B}\hat{\uparrow}\hat{v}})$, where $\mathbf{P}^{\mathcal{B}\hat{\uparrow}\hat{v}}(\hat{v}, \hat{v}) = 1$, $\mathbf{P}^{\mathcal{B}\hat{\uparrow}\hat{v}}(s, \hat{v}) = \mathbf{P}(s, v)$ and $\mathbf{P}^{\mathcal{B}\hat{\uparrow}\hat{v}}(s, v) = 0$ for all $s \in S^{\mathcal{D}|_B}$, $\mathbf{P}^{\mathcal{B}\hat{\uparrow}\hat{v}}(s, t) = \mathbf{P}(s, t)$ for all $s \in S$, $t \in S \backslash \{v\}$, and $\iota_{init}^{\mathcal{B}\hat{\uparrow}\hat{v}}(v) = 1$.

The stationary distribution within a DTMC is then given by $\theta^{\mathcal{D}|_B} = \frac{\mathbb{E}^{\mathcal{B}\hat{v}}[vt_s]}{\sum_{t\in S}\mathbb{E}^{\mathcal{B}\hat{v}}[vt_t]}$, as proven in [MKQW25].

(Trivially then uppper bound for stationary distribution given by uEVT and lower bound given by lEVT)

Definition 4.30 (Intermediate sets are closed under composition)
For any DTMC $\mathcal{D}$ and states $t, s, u, w \in S^{\mathcal{D}}$: if $u \in I^{\mathcal{D}}_{ts}$ and $w \in I^{\mathcal{D}}_{us}$, then $w \in I^{\mathcal{D}}_{ts} \cup \{t\}$.

Proof 22
Let ρ witness $u \in I^{\mathcal{D}}_{ts}$, i.e. a path fragment $t \to \cdots \to u \to \cdots \to s$ passing through u at some index k, and let σ witness $w \in I^{\mathcal{D}}_{us}$, i.e. a path fragment $u \to \cdots \to w \to \cdots \to s$. Truncating ρ at its first visit to u and appending σ yields a valid path fragment from t to s (transitions still respect $\mathbf{P}^{\mathcal{D}}$) that passes through w. Hence w is intermediate for (t, s), unless $w = t$. □

Corollary 4.31
If $R \subseteq S^{\mathcal{D}}$ contains t and $I^{\mathcal{D}}_{ts}$, then $I^{\mathcal{D}}_{us} \subseteq R \cup \{s\}$ for every $u \in R$. We call such an *R closed (w.r.t. $\mathcal{D}$ and s).*

Lemma 4.32 (Persistence on a closed region)
Let $\mathcal{D} \models_U \mathcal{I}$ and let $R \subseteq S$ be closed w.r.t. $\mathcal{D}$ and s (corollary 4.31) with $Pr^{\mathcal{D}}_u(\Diamond s) = M(u)$ for every $u \in R$. If $\mathcal{D}' \models_U \mathcal{I}$ satisfies $\mathbf{P}^{\mathcal{D}'}(u, \cdot) = \mathbf{P}^{\mathcal{D}}(u, \cdot)$ for every $u \in R$, then $Pr^{\mathcal{D}'}_u(\Diamond s) = M(u)$ for every $u \in R$ as well.

Proof 23
Fix $u \in R$. Every path fragment $\pi \in \Diamond_u s$ (w.r.t. $\mathcal{D}$) has all its intermediate states in $I^{\mathcal{D}}_{us} \subseteq R \cup \{s\}$ (corollary 4.31), so every transition of π leaves from a state in R; since $\mathbf{P}^{\mathcal{D}'} = \mathbf{P}^{\mathcal{D}}$ on R, π has the identical probability under $\mathcal{D}'$, and is a valid (positive-probability) path fragment of $\mathcal{D}'$ too. Summing over these paths, $Pr^{\mathcal{D}}_u(\Diamond s) \leq Pr^{\mathcal{D}'}_u(\Diamond s)$. Since $\mathcal{D}' \models_U \mathcal{I}$, also $Pr^{\mathcal{D}'}_u(\Diamond s) \leq M(u) = Pr^{\mathcal{D}}_u(\Diamond s)$. Hence equality. □

Lemma 4.33 (One-state extension)
Let $R \subseteq S$, $\mathcal{D} \models_U \mathcal{I}$ be closed and optimal on R as in lemma 4.32, and let $t^* \in S_{tr} \setminus R$. Let $\mathcal{D}_{t^*} \models_U \mathcal{I}$ satisfy $Pr^{\mathcal{D}_{t^*}}_{t^*}(\Diamond s) = M(t^*)$ (sec. 4.2.1), and set

$D := I^{\mathcal{D}_{t^*}}_{t^*s} \cup \{t^*\}$. Define $\mathcal{D}'$ by

$$\mathbf{P}^{\mathcal{D}'}(u,\cdot) := \begin{cases} \mathbf{P}^{\mathcal{D}_{t^*}}(u,\cdot) & u \in D \setminus R \\ \mathbf{P}^{\mathcal{D}}(u,\cdot) & \text{otherwise.} \end{cases}$$

(I.e. $\mathcal{D}'$ equals $\mathcal{D}$ except that rows of states in $D \setminus R$ are overwritten by $\mathcal{D}_{t^*}$'s rows.) Then $\mathcal{D}' \models_U \mathcal{I}$, and $Pr^{\mathcal{D}'}_u(\Diamond s) = M(u)$ for every $u \in R \cup \{t^*\}$.

Proof 24
$\mathcal{D}' \models_U \mathcal{I}$ is immediate, since every row of $\mathcal{D}'$ is copied verbatim from a DTMC already satisfying $\mathcal{I}$. Optimality on R follows directly from lemma 4.32, since $\mathbf{P}^{\mathcal{D}'} = \mathbf{P}^{\mathcal{D}}$ on R.

It remains to show $Pr^{\mathcal{D}'}_{t^*}(\Diamond s) = M(t^*)$. Write $B := D \setminus R$ (so $t^* \in B$) and $A := D \cap R$. By construction, $\mathbf{P}^{\mathcal{D}'} = \mathbf{P}^{\mathcal{D}_{t^*}}$ on all of B. Consider the DTMC X obtained by running $\mathcal{D}'$ started at t^*, and Y obtained by running $\mathcal{D}_{t^*}$ started at t^*. Because their transition laws agree on B, we may couple X and Y on the same probability space so that $X_k = Y_k$ for all $k \le \tau$, where $\tau := \min\{k \ge 1 : X_k \notin B\}$ is the first exit time from B (a stopping time for both processes, by the coupling).

At time τ, $X_\tau = Y_\tau$ lands in exactly one of three disjoint sets:

- $\{s\}$;
- $A = D \cap R \subseteq R$ (more generally, landing anywhere in R is possible — this can only happen via a transition out of $B \subseteq D$ under $\mathbf{P}^{\mathcal{D}_{t^*}}$, so in particular $A \subseteq R$ covers this; we simply write $u \in R$ for the landing state);
- some $d \notin R \cup \{s\}$; since τ exits $B = D \setminus R$ and $d \notin R$, also $d \notin D$ (as $D = A \cup B$), i.e. $d \notin I^{\mathcal{D}_{t^*}}_{t^*s} \cup \{t^*, s\}$, so by definition $Pr^{\mathcal{D}_{t^*}}_d(\Diamond s) = 0$.

By the strong Markov property applied to Y under $\mathcal{D}_{t^*}$ at the stopping time τ (standard first-step analysis for DTMCs, cf. the argument underlying theorem 4.18):

$$M(t^*) = Pr^{\mathcal{D}_{t^*}}_{t^*}(\Diamond s) = P(Y_\tau = s)\cdot 1 + \sum_{u \in R} P(Y_\tau = u)\cdot Pr^{\mathcal{D}_{t^*}}_u(\Diamond s) + \underbrace{\sum_{d \notin R \cup \{s\}} P(Y_\tau = d)\cdot 0}_{=0}. \tag{$*$}$$

Applying the same decomposition to X under $\mathcal{D}'$:

$$Pr^{\mathcal{D}'}_{t^*}(\Diamond s) = P(X_\tau = s)\cdot 1 + \sum_{u \in R} P(X_\tau = u)\cdot Pr^{\mathcal{D}'}_u(\Diamond s) + \sum_{d \notin R \cup \{s\}} P(X_\tau = d)\cdot Pr^{\mathcal{D}'}_d(\Diamond s).$$

By the coupling, $P(X_\tau = \cdot) = P(Y_\tau = \cdot)$. By lemma 4.32, $Pr^{\mathcal{D}'}_u(\Diamond s) = M(u) =$

$Pr_u^{\mathcal{D}_{t^*}}(\Diamond s)$ for every $u \in R$ (the last equality since $\mathcal{D}_{t^*}$ is optimal on all of $I_{t^*s}^{\mathcal{D}_{t^*}} \supseteq A$ by theorem 4.18, and both sides equal $M(u)$). The last sum is ≥ 0. Hence, comparing with $(*)$,

$$Pr_{t^*}^{\mathcal{D}'}(\Diamond s) \;\geq\; P(Y_\tau = s) + \sum_{u \in R} P(Y_\tau = u)\, M(u) + 0 \;=\; M(t^*).$$

On the other hand $Pr_{t^*}^{\mathcal{D}'}(\Diamond s) \leq M(t^*)$ trivially, since $\mathcal{D}' \models_U \mathcal{I}$ and $M(t^*)$ is the supremum over all such DTMCs. Hence $Pr_{t^*}^{\mathcal{D}'}(\Diamond s) = M(t^*)$. □

Theorem 4.34
Given an IDTMC $\mathcal{I}$ and a (goal) state $s \in S$ there exists a DTMC $\mathcal{D} \models_U \mathcal{I}$ such that $Pr_t^{\mathcal{D}}(\Diamond s) = \sup_{\mathcal{D}_i \models_U \mathcal{I}} Pr_t^{\mathcal{D}_i}(\Diamond s)$ for all transient states $t \in S_{tr}$.

Proof 25
Since S is countable (def. 4.1(i)), so is S_{tr}; let $t_1, t_2, t_3, \ldots$ enumerate it (if S_{tr} is finite the construction below simply terminates after $|S_{tr}|$ steps and no limit is needed).

Construction. Let $\mathcal{D}^{(1)} \models_U \mathcal{I}$ satisfy $Pr_{t_1}^{\mathcal{D}^{(1)}}(\Diamond s) = M(t_1)$ (sec. 4.2.1) and set $R_1 := I_{t_1 s}^{\mathcal{D}^{(1)}} \cup \{t_1, s\}$; by theorem 4.18, every $u \in R_1$ satisfies $Pr_u^{\mathcal{D}^{(1)}}(\Diamond s) = M(u)$, and R_1 is closed by corollary 4.31.

Inductively, given $\mathcal{D}^{(n)} \models_U \mathcal{I}$ and $R_n \ni s$, closed and optimal (i.e. $Pr_u^{\mathcal{D}^{(n)}}(\Diamond s) = M(u)$ for all $u \in R_n$): if $t_{n+1} \in R_n$, set $\mathcal{D}^{(n+1)} := \mathcal{D}^{(n)}$, $R_{n+1} := R_n$. Otherwise apply lemma 4.33 with $R = R_n$, $\mathcal{D} = \mathcal{D}^{(n)}$, $t^* = t_{n+1}$ to obtain $\mathcal{D}^{(n+1)} \models_U \mathcal{I}$ optimal on $R_n \cup \{t_{n+1}\}$; set $R_{n+1} := R_n \cup I_{t_{n+1}s}^{\mathcal{D}^{(n+1)}} \cup \{t_{n+1}\}$. By theorem 4.18 applied to $\mathcal{D}^{(n+1)}$ (now known optimal at t_{n+1}), every $u \in I_{t_{n+1}s}^{\mathcal{D}^{(n+1)}}$ is optimal in $\mathcal{D}^{(n+1)}$ too, so R_{n+1} is fully optimal; R_{n+1} is closed by applying corollary 4.31 to each generator $t_1, \ldots, t_{n+1}$ in turn.

Crucially, by construction (either branch), $\mathbf{P}^{\mathcal{D}^{(n+1)}}(u, \cdot) = \mathbf{P}^{\mathcal{D}^{(n)}}(u, \cdot)$ for every $u \in R_n$ — rows are only ever overwritten on $D \setminus R_n$, disjoint from R_n. Since $R_n \subseteq R_{n+1}$ for all n, this "freezing" persists: for all $m \geq n$, $\mathbf{P}^{\mathcal{D}^{(m)}}(u, \cdot) = \mathbf{P}^{\mathcal{D}^{(n)}}(u, \cdot)$ for every $u \in R_n$.

Passing to the limit. For a state $u \in S$, let $m(u) := \min\{n : u \in R_n\}$ if it exists. By the freezing property, $\mathbf{P}^{\mathcal{D}^{(n)}}(u, \cdot)$ is constant in n for $n \geq m(u)$; if u never enters any R_n its row is likewise never touched past $\mathcal{D}^{(1)}$. Either way $\lim_{n\to\infty} \mathbf{P}^{\mathcal{D}^{(n)}}(u, \cdot)$ exists for every u; define $\mathcal{D}^{(\infty)}$ by these limiting rows. Since each row is copied from some $\mathcal{D}^{(n)} \models_U \mathcal{I}$, we get $\mathcal{D}^{(\infty)} \models_U \mathcal{I}$.

For each i, rows agree between $\mathcal{D}^{(\infty)}$ and $\mathcal{D}^{(i)}$ on all of R_i (both equal the frozen

value from step i onward), R_i is closed and optimal for $\mathcal{D}^{(i)}$, and $t_i \in R_i$. By lemma 4.32 (with $\mathcal{D} = \mathcal{D}^{(i)}$, $\mathcal{D}' = \mathcal{D}^{(\infty)}$, $R = R_i$),

$$Pr_{t_i}^{\mathcal{D}^{(\infty)}}(\Diamond s) = M(t_i).$$

As $t_1, t_2, \ldots$ enumerate all of S_{tr}, $\mathcal{D}^{(\infty)}$ simultaneously attains $Pr_t^{\mathcal{D}^{(\infty)}}(\Diamond s) = \sup_{\mathcal{D}_i \models_U \mathcal{I}} Pr_t^{\mathcal{D}_i}(\Diamond s)$ for every transient state $t \in S_{tr}$. □

Chapter 5

Conclusion

Appendix A

Appendix

Enter text

Bibliography

[BDL+17] Anicet Bart, Benoit Delahaye, Didier Lime, Eric Monfroy, and Charlotte Truchet. Reachability in Parametric Interval Markov Chains using Constraints, 2017.

[BDLM21] Giovanni Bacci, Benoît Delahaye, Kim G. Larsen, and Anders Mariegaard. *Quantitative Analysis of Interval Markov Chains*, pages 57–77. Springer International Publishing, Cham, 2021.

[BK08] Christal Baier and {Joost P.} Katoen. *Principles of Model Checking*. MIT Press, United States, May 2008.

[BLW13] Michael Benedikt, Rastislav Lenhardt, and James Worrell. LTL Model Checking of Interval Markov Chains. In Nir Piterman and Scott A. Smolka, editors, *Tools and Algorithms for the Construction and Analysis of Systems*, pages 32–46, Berlin, Heidelberg, 2013. Springer Berlin Heidelberg.

[CHK13] Taolue Chen, Tingting Han, and Marta Kwiatkowska. On the complexity of model checking interval-valued discrete time Markov chains. *Information Processing Letters*, 113(7):210–216, 2013.

[Cho17] Ventsislav Chonev. Reachability in Augmented Interval Markov Chains, 2017.

[CSH08] Krishnendu Chatterjee, Koushik Sen, and Thomas A. Henzinger. Model-Checking ω-Regular Properties of Interval Markov Chains. In Roberto Amadio, editor, *Foundations of Software Science and Computational Structures*, pages 302–317, Berlin, Heidelberg, 2008. Springer Berlin Heidelberg.

[KS76] John G. Kemeny and J. Laurie Snell. *Finite Markov Chains*. Undergraduate Texts in Mathematics. Springer, 1976.

[KTDB19] Thomas Krak, Natan T'Joens, and Jasper De Bock. Hitting Times and Probabilities for Imprecise Markov Chains. In Jasper De Bock, Cassio P. de Campos, Gert de Cooman, Erik Quaeghebeur, and Gregory Wheeler,

editors, *Proceedings of the Eleventh International Symposium on Imprecise Probabilities: Theories and Applications*, volume 103 of *Proceedings of Machine Learning Research*, pages 265–275. PMLR, 03–06 Jul 2019.

[Kul16] Vidyadhar Kulkarni. *Modeling and analysis of stochastic systems: Third edition.* 11 2016.

[MKQW25] Hannah Mertens, Joost-Pieter Katoen, Tim Quatmann, and Tobias Winkler. Computing Expected Visiting Times and Stationary Distributions in Markov Chains: Fast and Accurate. *Journal of Automated Reasoning*, 69(3), August 2025.

[PV91] Panos M. Pardalos and Stephen A. Vavasis. Quadratic programming with one negative eigenvalue is NP-hard. *Journal of Global Optimization*, 1:15–22, 1991.

[SVA06] Koushik Sen, Mahesh Viswanathan, and Gul Agha. Model-Checking Markov Chains in the Presence of Uncertainties. In Holger Hermanns and Jens Palsberg, editors, *Tools and Algorithms for the Construction and Analysis of Systems*, pages 394–410, Berlin, Heidelberg, 2006. Springer Berlin Heidelberg.