

Introduction

We present an open-source software package - **SCOTER** - for multiple-event location. The tool implements source-specific station terms (SSST) technique (Richards-Dinger & Shearer, 2000; Lin & Shearer, 2005) to reduce the effect of spatially correlated residuals caused by 3-D velocity structure. By using this technique, a set of P- and/or S-wave station corrections is computed iteratively for each source-receiver path. The calculated station terms are added to the travel times computed from the velocity model to take the unmodelled structure into account. Thus, the travel-time residual misfit function is a function of both the station terms and the hypocenter parameters (Nooshiri et al., 2017).




Figure 1: A cartoon illustrating source-specific station term calculation.

The basic algorithm applied is described by the following steps:

- All the earthquakes are located with the actual arrival times.
- SSST values are calculated for each source-receiver pair as the mean of the residuals from neighboring events located within a sphere of radius r_{max} around the target event.
- The original arrival time picks are corrected by the calculated SSSTs.
- The earthquakes are relocated with these corrected arrival times.
- The above steps are repeated iteratively.

Some Features of the SCOTER Program

- ✓ Supports local, regional, and teleseismic location.
- ✓ Adopted for non-linear, global-search location method implemented in the software package **NonLinLoc** (Lomax et al., 2000).
- ✓ Allows each neighboring event to be directly weighted according to the inter-event distance and residual performance.
- ✓ Uses **Pyrocko** toolbox and library for data handling (e.g. QuakeML data format).
- ✓ Can be run as a command-line tool:

```
# To get further help and a list of available options and subcommands:
$ scoter --help
# To get an example configuration file:
$ scoter init myconfig.sf
# To run SCOTER:
$ scoter go myconfig.sf --steps=A,B,C --parallel=8
```

Code Availability

<https://gitext.gfz-potsdam.de/nooshiri/scoter>

Application to the Seismicity Along the Northern Arctic Ridge




Figure 2:

Calculated source-specific station terms (SSSTs) for regional seismic array SPITS for (a) first P arrivals and (b) first S arrivals. The SSST values are color coded and plotted at the event locations.





Figure 4:

Histograms of the travel-time residuals from (top) single-event and (bottom) the SSST relocations for the station shown in Fig. 2. The scaled median absolute deviation (smad), mean, and sample standard deviation of each distribution are indicated on each panel.

Figure 3:

Relocated seismicity in the Arctic Ridge using (a) a single-event, non-linear algorithm and (b) source-specific station correction terms (SSSTs) implemented in the SCOTER program, and (c) BayesLoc locations presented in Gibbons et al. (2017).




Dataset

A catalog of 614 events with magnitude range of M 2.0-6.0 in the Arctic region since 1998 has been generated using high-quality Pn and Sn readings on stations on both sides of the spreading ridge in addition to teleseismic P phases (Gibbons et al., 2017).

Figure 5:

Frequency of occurrence (heat plots) of travel-time residuals for (top) single-event and (bottom) the SSST locations. The heat plots are normalized in each distance range so that the residual bin with the largest number of hits (the scaling cell) is assigned a value of 1.



Summary & Conclusion

- SCOTER is a publicly available software package that applies the station correction terms to perform precise relocations of earthquake hypocenters.
- SCOTER is run as a command line tool and provides a set of sub-commands to develop inputs (dataset, configuration etc) and export results (hypocenter parameters, travel-time residuals etc).
- Relocation results for the seismicity in the Arctic region demonstrate the method's viability in illuminating key features of the seismicity that are obscured by single-event locations.

References

- Gibbons, S.J., Harris, D.B., Dahl-Jensen, T., Kvaerna, T., Larsen, T.B., Paulsen, B., and Voss, P.H. (2017). Locating seismicity on the Arctic plate boundary using multiple-event techniques and empirical signal processing. *Geophys. J. Int.*, 211:1613-1627.
- Heimann, S., Kriegerowski, M., Isken, M., Cesca, S., Daout, S., Grigoli, F., Juretzek, C., Megies, T., Nooshiri, N., Steinberg, A., and et al. (2017). Pyrocko - an open-source seismology toolbox and library. GFZ Data Services. <http://doi.org/10.5880/GFZ.2.1.2017.001>.
- Lin, G. and Shearer, P. (2005). Tests of relative earthquake location techniques using synthetic data. *J. Geophys. Res.*, 110(B4), B04304.
- Lomax, A., Virieux, J., Volant, P., and Berge-Thierry, C. (2000). Probabilistic earthquake location in 3D and layered models. In Thurber, C. and Rabinowitz, N., editors, *Advances in Seismic Event Location*, volume 18 of *Modern Approaches in Geophysics*, pages 101-134.
- Nooshiri, N., Saul, J., Heimann, S., Tilmann, F., and Dahm, T. (2017). Revision of earthquake hypocentre locations in global bulletin data sets using source-specific station terms. *Geophys. J. Int.*, 208(2):589-602.
- Richards-Dinger, K. B. and Shearer, P. M. (2000). Earthquake locations in southern California obtained using source-specific station terms. *J. Geophys. Res.*, 105(B5):10939-10960.