

DIFFERENTIAL COHOMOLOGY SEMINAR 4 (DRAFT)

TALK BY ALESSANDRO NANTO

In the last talk we learned the definition of a differential cohomology theory, as a sheaf valued in spectra on the site of manifolds. This talk continues our journey through differential cohomology theories, and focuses on the following three topics:

- (1) We want to learn how to construct non-trivial examples out of sheaves valued in chain complexes.
- (2) We want to understand how we can extend classical cohomology operations to the setting of differential cohomology theories.
- (3) We want to introduce suitable analogues of fiber-wise integration.

1. ABELIAN GROUPS, SPECTRA AND THE HEART

Let us start by reviewing the relation between abelian groups, rings and spectra.

Definition 1.1. Let $n \in \mathbb{Z}$ and X be a spectrum, define $\pi_n(X) := \pi_0(\Omega^{\infty+n} X) = \pi_0(X_{-n})$. We call π_n the n -th homotopy group of X .

Remark 1.2. Note that since $X_n \simeq \Omega^2 X_{n+2}$, for any n , the set $\pi_0(X_n)$ underlies the structure of an abelian group.

The category $\mathcal{S}p$ underlies the structure of a symmetric monoidal ∞ -category ([Lur17, Corollary 4.8.2.19]). Following [Lur17], we denote by \otimes the tensor product on $\mathcal{S}p$.

Definition 1.3. A (commutative) algebra object in $\mathcal{S}p$ is called an (\mathbb{E}_∞) -ring spectrum, see [Lur17, Definition 7.1.0.1]. Given a ring spectrum R , denote by Mod_R the corresponding category of left R -module spectra, see [Lur17, Definition 7.1.1.2].

Remark 1.4. The sphere spectrum \mathbb{S} acts as the monoidal unit of $\mathcal{S}p$, therefore it is a \mathbb{E}_∞ -ring spectrum. The category $\text{Mod}_{\mathbb{S}}$ is canonically equivalent to $\mathcal{S}p$.

Definition 1.5. Denote by $\mathcal{S}p_{\geq 0} \subseteq \mathcal{S}p$ the full sub-category generated by *connective spectra*, i.e. spectra X such that $\pi_n(X) \simeq 0$, for all $n < 0$. Denote by $\mathcal{S}p^\heartsuit \subseteq \mathcal{S}p_{\geq 0}$ the *heart of spectra*, i.e. the full sub-category generated by spectra X such that $\pi_n(X) \simeq 0$, for all $n > 0$.

We have the following result relating connective spectra and the heart, which follow immediately.

Lemma 1.6. *Let X be a connective spectrum. The following are equivalent:*

- (1) X is in the heart.
- (2) $\pi_n(\Omega^\infty X) = 0$, for all $n > 0$.
- (3) $\text{Hom}_{\mathcal{S}_*}(S, \Omega^\infty X) \simeq 0$, for all connected, pointed spaces S .
- (4) X is local with respect to the class of maps $\Sigma^\infty S \rightarrow 0$, for every connected pointed space S .

The category $\mathcal{S}p_{\geq 0}$ is presentable and π_0 induces an equivalence between the heart and $\mathcal{A}b$ ([Lur17, Proposition 1.4.3.6]). The heart is a sub-category of local objects of connective spectra, therefore the inclusion $\mathcal{A}b \simeq \mathcal{S}p^\heartsuit \subseteq \mathcal{S}p_{\geq 0}$ is a right adjoint. The category $\mathcal{S}p_{\geq 0}$ is closed under \otimes and, given X, Y connective spectra,

$$(1.7) \quad \pi_0(X \otimes Y) \simeq \pi_0(X) \otimes \pi_0(Y)$$

see [Dav24, Theorem 2.3.28]

Definition 1.8. Given an abelian group A , denote by HA the (unique up to equivalence) spectrum of the heart such that $\pi_0(HA) \simeq A$. We call HA the *Eilenberg-Mac Lane spectrum* of A .

Using [Equation \(1.7\)](#) and the adjunction between H and π_0 , one can prove H , viewed as a functor $\mathcal{A}b \rightarrow \mathcal{S}p$, is lax monoidal. In particular, if R is a commutative ring, then HR is a connective \mathbb{E}_∞ -ring spectrum. On the other hand, if R is a connective \mathbb{E}_∞ -ring spectrum and M a connective module, then $\pi_0(M)$ is a $\pi_0(R)$ -module.

Definition 1.9. Given a commutative ring R , denote by $\mathrm{Ch}(R) = \mathrm{Ch}(\mathrm{Mod}_R)$ the ordinary category of unbounded chain complexes. Let $\mathcal{D}(R)$ be the ∞ -localization of $\mathrm{Ch}(R)$ at the class of quasi-isomorphisms.

Similar to the heart of spectra, given an \mathbb{E}_∞ -ring spectrum R , denote by $\mathrm{Mod}_R^\heartsuit \subseteq \mathrm{Mod}_R$ the full subcategory generated by R -modules such that the underlying spectrum belongs to the heart of spectra.

Theorem 1.10 (Stable Dold-Kan Correspondence). *Let R be a commutative ring.*

- (1) $\mathrm{Mod}_R \simeq \mathrm{Mod}_{HR}^\heartsuit$ via taking Eilenberg-Mac Lane spectra.
- (2) The equivalence in (1) extends to an equivalence $H : \mathcal{D}(R) \simeq \mathrm{Mod}_{HR}$ of symmetric monoidal ∞ -categories.

Proof. (1) is [[Lur17](#), Proposition 7.1.1.13], while (2) is [[Lur17](#), Theorem 7.1.2.13]. □

An interesting consequence of [Theorem 1.10](#) is the following:

Corollary 1.11. *Given $F \in \mathcal{D}(R)$, then $\pi_n(HF) \simeq H_n(F)$, for all $n \in \mathbb{Z}$.*

Proof.

$$\begin{aligned}
 \pi_n(HF) &= \pi_0(\Omega^{\infty+n} HF) \\
 &\stackrel{\textcircled{1}}{\simeq} \pi_0(\mathrm{Hom}_{\mathcal{S}p}(\Sigma^n \mathcal{S}, HF)) \\
 &\stackrel{\textcircled{2}}{\simeq} \pi_0(\mathrm{Hom}_{\mathrm{Mod}_{HR}}(\Sigma^n HR, HF)) \\
 &\stackrel{\textcircled{3}}{\simeq} \pi_0(\mathrm{Hom}_{\mathcal{D}(R)}(R[n], F)) \\
 &\stackrel{\textcircled{4}}{\simeq} H_n(F)
 \end{aligned}$$

- ① The functor $\Omega^{\infty+n}$ is corepresented by the shifted sphere spectrum $\Sigma^n \mathcal{S}$.
- ② The forgetful functor $\mathrm{Mod}_{HR} \rightarrow \mathrm{Mod}_{\mathcal{S}} \simeq \mathcal{S}p$ is right adjoint to tensoring with HR and $HR \otimes (\Sigma^n \mathcal{S}) \simeq \Sigma^n HR$.
- ③ [Theorem 1.10](#)
- ④ π_0 of the mapping space $\mathrm{Hom}_{\mathcal{D}(R)}(R[n], F)$ is equivalent to the mapping space $R[n] \rightarrow F$ in the *ordinary* derived category of R , i.e. homotopy classes of maps $R[n] \rightarrow F$, which correspond exactly to classes in $H_n(F)$. □

2. LOCALLY CONSTANT SHEAVES ON MANIFOLDS

Let \mathcal{C} be a presentable ∞ -category. The ∞ -categorical background given in previous talks allows to conclude the existence of a number of functors. Here we give a (somewhat) explicit formula for one.

Remark 2.1. Let \mathcal{B} be a full, dense sub-site of $\mathcal{M}fd$, recall then that, for every presentable category \mathcal{C} , the restriction functor induces an equivalence $\mathrm{Shv}(\mathcal{M}fd, \mathcal{C}) \simeq \mathrm{Shv}(\mathcal{B}, \mathcal{C})$.

Evaluation at $\{0\}$ induces an adjunction $(L, \Gamma) : \mathcal{C} \rightarrow \mathrm{Shv}(\mathcal{M}fd, \mathcal{C})$, where the functor Γ is evaluation at $\{0\}$, while the left adjoint L maps $C \in \mathcal{C}$ to the sheafification of the constant pre-sheaf with value C .

Remark 2.2. Every presentable ∞ -category \mathcal{C} is uniquely *cotensored over* \mathcal{S} , see [[Lur09](#), Remark 5.5.2.6]. More explicitly, for every space S and object C , there is an object C^S together with a natural equivalence

$$\mathrm{Hom}_{\mathcal{S}}(S, \mathrm{Hom}_{\mathcal{C}}(-, C)) \simeq \mathrm{Hom}_{\mathcal{C}}(-, C^S)$$

Definition 2.3. Denote by Sing the functor $\mathcal{M}fd \rightarrow \mathcal{S}$ mapping a manifold to its underlying space of *smooth* simplexes. Given a presentable ∞ -category \mathcal{C} , denote by \flat the composition $\mathcal{C} \rightarrow \mathrm{Fun}(\mathcal{S}^{\mathrm{op}}, \mathcal{C}) \rightarrow \mathrm{Fun}(\mathcal{M}fd^{\mathrm{op}}, \mathcal{C})$, the first functor coming from [Remark 2.2](#), the second being pre-composition with $\mathrm{Sing}^{\mathrm{op}}$.

Explicitly, given an object $C \in \mathcal{C}$, the associated pre-sheaf $\flat C$ maps a manifold M to $C^{\mathrm{Sing}(M)}$.

Remark 2.4. Given a topological space X , denote by $\text{Sing}^{\text{top}}(X)$ the corresponding space, i.e. its singular simplicial set. If $X = M$ is a smooth manifold, then $\text{Sing}(M) \subseteq \text{Sing}^{\text{top}}(M)$ and the inclusion is a homotopy equivalence by Whitney's Approximation Theorem, see [Zuo21, Theorem 1.6].

Lemma 2.5 ([BG16, Corollary 6.46]). *Sing is a cosheaf. In particular, \flat factors through $\text{Shv}(\text{Mfd}, \mathcal{C}) \subseteq \text{Fun}(\text{Mfd}^{\text{op}}, \mathcal{C})$.*

Lemma 2.5 is essentially the consequence of a generalized version of Seifert-van Kampen theorem, namely [Lur17, Proposition A.3.2], stating that, given a topological space X and a covering sieve \mathcal{O} , the space $\text{Sing}^{\text{top}}(X)$ is the colimit of $\text{Sing}^{\text{top}}(U)$, over $U \in \mathcal{J}(\mathcal{O})$.

Theorem 2.6. *$\flat : \mathcal{C} \rightarrow \text{Shv}(\text{Mfd}, \mathcal{C})$ is left adjoint to Γ .*

Proof. The composition $\mathcal{C} \xrightarrow{\flat} \text{Shv}(\text{Mfd}, \mathcal{C}) \xrightarrow{|\mathcal{E}_{\text{uc}}|} \text{Shv}(\mathcal{E}_{\text{uc}}, \mathcal{C})$ maps an object C to the sheaf $\flat C$ restricted to Euclidean spaces. Since \mathbb{R}^n is contractible, $(\flat C)(\mathbb{R}^n) = C^{\text{Sing}(\mathbb{R}^n)} \simeq C$ and so \flat restricted to \mathcal{E}_{uc} is equivalent to L , the functor taking C to the pre-sheaf with constant value C , which is left adjoint to Γ restricted to \mathcal{E}_{uc} . \square

Remark 2.7. The proof of **Theorem 2.6** shows that, given an object $C \in \mathcal{C}$ and a dense sub-site $\mathcal{B} \subseteq \text{Mfd}$ of contractible manifolds, the constant pre-sheaf with value C is equivalent to the sheaf $\flat C$ restricted to \mathcal{B} , hence it is already a sheaf.

3. CHAIN COMPLEXES AND SHEAVES

The stable Dold-Kan correspondence allows us to move freely between sheaves of $H\mathbb{Z}$ -module spectras and sheaves valued in the derived category $\mathcal{D}(\mathbb{Z})$ of abelian groups. In this section, we introduce definition and some technical lemmas regarding sheaves on manifolds valued in the derived category.

Remark 3.1. All chain complexes are \mathbb{Z} -graded. Given a ring R , we identify chain complexes with cochain complexes. More specifically, we identify a cochain complex V^* with the chain complex $V_n = V^{-n}$. Viceversa, we identify V_* with the cochain complex $V^n = V_{-n}$. In particular, operations defined on cochain complexes are extended to chain complexes via this identification.

Definition 3.2 ([BNV16, Definition 7.14]). Given $n \in \mathbb{Z}$ and a cochain complex V^* , denote by $V^{\geq n}$, resp. $V^{\leq n}$, the naive truncations

$$\cdots \rightarrow 0 \rightarrow V^n \rightarrow V^{n+1} \rightarrow \cdots, \quad \text{resp.} \quad \cdots \rightarrow V^{n-1} \rightarrow V^n \rightarrow 0 \rightarrow \cdots$$

Given $F : \text{Mfd}^{\text{op}} \rightarrow \text{Ch}(\mathbb{Z})$, denote by $F^{\geq n}$ and $F^{\leq n}$ the point-wise naive truncations of F . If F is a sheaf of chain complexes, then so are its truncations.

Lemma 3.3 ([BNV16, Lemma 7.12]). *Let $F : \text{Mfd}^{\text{op}} \rightarrow \text{Ch}(\mathbb{Z})$ a sheaf of chain complexes of C^∞ -modules, then $\text{Mfd}^{\text{op}} \xrightarrow{F} \text{Ch}(\mathbb{Z}) \rightarrow \mathcal{D}(\mathbb{Z})$ is a sheaf.*

Definition 3.4. Denote by Ω^* the sheaf $\text{Mfd}^{\text{op}} \rightarrow \text{Ch}(\mathbb{Z})$ mapping a manifold to its de Rham complex.

Lemma 3.3 ensures that the sheaf in **Definition 3.4** and the corresponding naive truncations remain sheaves after post-composition with the localization functor $\text{Ch}(\mathbb{Z}) \rightarrow \mathcal{D}(\mathbb{Z})$.

Definition 3.5. Given a sheaf $F : \text{Mfd}^{\text{op}} \rightarrow \mathcal{D}(\mathbb{Z})$, denote by HF the Eilenberg-Mac Lane sheaf of $H\mathbb{Z}$ -module spectra obtained by applying point-wise the equivalence of **Theorem 1.10**.

4. HIGHER DE RHAM THEOREM

The classical de Rham theorem gives an explicit ring isomorphism between the de Rham cohomology of a manifold M and its singular cohomology with real coefficients. Using the modern perspective we can recover de Rham theorem as a corollary of a more general \mathbb{A}_∞ -quasi-isomorphism of differential graded algebras. An \mathbb{A}_∞ -morphism of DGAs $\mu : R \rightarrow S$ consists of a sequence of graded linear maps $\{\mu_n : R^{\otimes n} \rightarrow S\}_{n \geq 1}$ satisfying a sequence of coherence conditions, see [LV12, Proposition 10.2.12]. One such condition is that μ_1 is a chain map, then a \mathbb{A}_∞ -quasi-isomorphism is an \mathbb{A}_∞ -morphism μ such that μ_1 is a quasi-isomorphism. Here we use the machinery set-up in **Section 2**.

Remark 4.1. Since $\mathcal{D}(\mathbb{Z})$ is presentable, we know that it is cotensored over \mathcal{S} . Given a space S and a chain complex M_* , the cotensor M_*^S is the chain complex of graded linear maps $C_*(S, \mathbb{Z}) \rightarrow M_*$, from the (normalized) singular chain complex of S to M_* , see [Lur17, Definition 1.3.2.1]. In particular, let $M_* = M$ be concentrated in degree 0, then M_*^S is the singular cochain complex of S with values in M .

Remark 4.2. Given a smooth manifold M , denote by $C_*^{\text{sm}}(M, \mathbb{Z})$ the *smooth* singular chain complex generated by the smooth singular simplicial set $\text{Sing}(M)$. Given an abelian group A , denote by $C_{\text{sm}}^*(M, A)$ the cochain complex valued in A associated to $C_*^{\text{sm}}(M, \mathbb{Z})$.

Definition 4.3. Consider the morphism $\Omega^*(M) \rightarrow (\mathfrak{b}\mathbb{R})(M) = C_{\text{sm}}^*(M, \mathbb{R})$ taking a form $\omega \in \Omega^n(M)$ to the linear map $\int \omega : C_n^{\text{sm}}(M, \mathbb{Z}) \rightarrow \mathbb{R}$. We call the induced transformation $\text{dR} : \Omega^* \rightarrow \mathfrak{b}\mathbb{R}$ the *de Rham morphism*.

We can now state the main theorem.

Theorem 4.4 ([Aa10, Theorem 3.25]). $\text{dR} : \Omega^* \rightarrow \mathfrak{b}\mathbb{R}$ lifts to an \mathbb{A}_∞ -quasi-isomorphism of sheaves of DGAs.

5. DELIGNE COHOMOLOGY

In this section, we give the definition of the ℓ -th Deligne sheaf as the Eilenberg-Mac Lane spectrum (see **Definition 3.5**) associated to a sheaf $F : \mathcal{M}\text{fd}^{\text{op}} \rightarrow \mathcal{D}(\mathbb{Z})$. Following that, we give explicit cochain complexes that are quasi-isomorphic to the value of F at a manifold M .

Definition 5.1. Given $\ell \in \mathbb{N}$, define $\hat{\mathbb{Z}}(\ell) : \mathcal{M}\text{fd}^{\text{op}} \rightarrow \mathcal{D}(\mathbb{Z})$ as the pullback

$$\begin{array}{ccc} \hat{\mathbb{Z}}(\ell) & \longrightarrow & \Omega^{\geq \ell} \\ \downarrow & & \downarrow \\ \mathfrak{b}\mathbb{Z} & \longrightarrow & \mathfrak{b}\mathbb{R} \end{array}$$

The vertical morphism being the composition $\Omega^{\geq \ell} \hookrightarrow \Omega^* \xrightarrow{\text{dR}} \mathfrak{b}\mathbb{R}$. We call the corresponding Eilenberg-Mac Lane spectrum $H\hat{\mathbb{Z}}(\ell)$ the ℓ -th *Deligne sheaf*.

Remark 5.2 (Model A, see [HS05, §3.2]). Let $\acute{C}(\ell)^n(M) \subseteq C_{\text{sm}}^n(M, \mathbb{Z}) \oplus C_{\text{sm}}^{n-1}(M, \mathbb{R}) \oplus \Omega^n(M)$ consist of triples (c, h, ω) for which $\omega = 0$ if $n < \ell$, with differential $\delta(c, h, \omega) = (\delta c, \text{dR}(\omega) - c - \delta h, d\omega)$. This complex $\acute{C}^*(\ell)(M)$ fits into a diagram

$$\begin{array}{ccc} \acute{C}^*(\ell)(M) & \longrightarrow & \Omega^{\geq \ell}(M) \\ \downarrow & & \downarrow \\ C_{\text{sm}}^*(M, \mathbb{Z}) & \longrightarrow & C_{\text{sm}}^*(M, \mathbb{R}) \end{array}$$

which commutes up to homotopy given by the projections $\acute{C}^n(\ell)(M) \rightarrow C_{\text{sm}}^{n-1}(M, \mathbb{R})$. The diagram above model the pullback of **Definition 5.1**, hence $\acute{C}^*(\ell)(M) \simeq \hat{\mathbb{Z}}(\ell)(M)$.

Remark 5.3 (Model B). Recall that \mathfrak{b} preserves cofiber sequences, since it is left adjoint, and that fiber sequences are the same a cofiber sequences in stable ∞ -categories. Consider the diagram

$$\begin{array}{ccc} \hat{\mathbb{Z}}(\ell) & \longrightarrow & \Omega^{\geq \ell} \\ \downarrow & & \downarrow \\ \mathfrak{b}\mathbb{Z} & \longrightarrow & \mathfrak{b}\mathbb{R} \\ \downarrow & & \downarrow \\ 0 & \longrightarrow & \mathfrak{b}(\mathbb{R}/\mathbb{Z}) \end{array}$$

Since the bottom square is an pullback, $\hat{\mathbb{Z}}(\ell)(M)$ is equivalent to the pullback of the diagram in **red**. Let $\check{C}^n(\ell)(M) \subseteq C_{\text{sm}}^{n-1}(M, \mathbb{R}/\mathbb{Z}) \oplus \Omega^n(M)$ consist of pairs (χ, ω) for which $\omega = 0$ if $n < \ell$, with differential $\delta(\chi, \omega) = (e^{2\pi i \text{dR}(\omega)} - \delta\chi, d\omega)$. Similar to **Remark 5.2**, the complex $\check{C}^*(\ell)(M)$ fits into the above diagram so that the outer square is an pullback, hence it is equivalent to $\hat{\mathbb{Z}}(\ell)(M)$.

Take an n -cocycle ($n \geq \ell$) in the model from [Remark 5.3](#), i.e. $(\chi, \omega) \in C_{\text{sm}}^{n-1}(M, \mathbb{R}/\mathbb{Z}) \oplus \Omega^n(M)$ such that $d\omega = 0$ and $\delta\chi = e^{2\pi i \text{dR}(\omega)}$. Such a cocycle determines a differential character of degree $n-1$ for M , in the sense of the following definition:

Definition 5.4 ([HS05, Definition 3.4], see also [BB14, Chapter 5]). Consider a manifold M , a *differential character* of degree n consists of a character $\chi : Z_n^{\text{sm}}(M, \mathbb{Z}) \rightarrow \mathbb{R}/\mathbb{Z}$ on the group of smooth n -cycles of M together with a n -form $\omega \in \Omega^{n+1}(M)$, such that, for every smooth $(n+1)$ -chain c ,

$$\chi(\partial c) = e^{2\pi i \int_c \omega}$$

Definition 5.5. Given a full sub-site $\mathcal{B} \subseteq \text{Mfd}$ and a manifold M , a \mathcal{B} -good open cover of M consists of an open cover \mathcal{O} such that every finite intersection of elements in \mathcal{O} is either empty or diffeomorphic to an object of \mathcal{B} . If $\mathcal{B} = \mathcal{E}\text{uc}$ the sub-site of Euclidean spaces, we call a \mathcal{B} -good open cover a *differentiably good open cover*, see [FSS11, Definition 6.3.9].

If \mathcal{B} is the sub-site of contractible manifolds, we recover the notion of a *good open cover*. In our next model, we will construct a sheaf F equivalent to $\hat{\mathbb{Z}}(\ell)$ on a dense sub-site \mathcal{B} . Given a generic manifold M , we can evaluate $\hat{\mathbb{Z}}(\ell)(M)$ using F , if we assume the existence of a \mathcal{B} -good open cover. Here we care mainly about $\mathcal{B} = \mathcal{E}\text{uc}$.

Theorem 5.6 ([FSS11, Theorem A.1]). *Every paracompact smooth manifold admits a differentiably good open cover.*

Remark 5.7 (Model C, see [ADH21, Lemma 7.3.4]). Let $\mathcal{B} \subseteq \text{Mfd}$ be a dense sub-site of contractible manifolds, consider the following diagram in the category of sheaves on \mathcal{B}

$$\begin{array}{ccccc} \hat{\mathbb{Z}}(\ell)|_{\mathcal{B}} & \longrightarrow & \Omega^{\geq \ell}|_{\mathcal{B}} & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow \\ \mathbb{Z} & \longrightarrow & \Omega^*|_{\mathcal{B}} & \longrightarrow & \Omega^{\leq \ell-1}|_{\mathcal{B}} \end{array}$$

The left square is obtained by: ① Apply the restriction to \mathcal{B} functor to the pullback diagram of [Definition 5.1](#) ② Substitute $\flat\mathbb{R}|_{\mathcal{B}}$ with $\Omega^*|_{\mathcal{B}}$ (see [Theorem 4.4](#)) and $\flat\mathbb{Z}|_{\mathcal{B}}$ with \mathbb{Z} (see [Remark 2.7](#)). Since the right square is a pullback, $\hat{\mathbb{Z}}(\ell)|_{\mathcal{B}}$ is equivalent to the pullback of the diagram in red. Let $\check{C}^*(\ell)$ be the sheaf of chain complexes $\mathbb{Z} \rightarrow \Omega^0 \rightarrow \dots \rightarrow \Omega^{\ell-1} \rightarrow 0 \rightarrow \dots$, where \mathbb{Z} is in degree 0 includes into Ω^0 as constant functions. The complex $\check{C}^*(\ell)$ fits into the above diagram, so that the outer square is an pullback, therefore $\hat{\mathbb{Z}}(\ell)|_{\mathcal{B}} \simeq \check{C}^*(\ell)$.

Remark 5.8 (Model C, Continued). Since $\hat{\mathbb{Z}}(\ell)$ is equivalent to $\check{C}^*(\ell)$ restricted to \mathcal{B} , consider a manifold M having a \mathcal{B} -good open cover \mathcal{O} and let $\mathcal{J}(\mathcal{O})$ the closure of \mathcal{O} under finite intersections, then recall

$$\hat{\mathbb{Z}}(\ell)(M) \simeq \lim_{U \in \mathcal{J}(\mathcal{O})} \check{C}^*(\ell)(U)$$

By rearranging the limit, we see it is equivalent to the limit of the cosimplicial object taking n to the product over all tuples $(U_0, \dots, U_n) \in \mathcal{O}^{n+1}$ of $\check{C}^*(\ell)(U_0 \cap \dots \cap U_n)$. We can then apply [BNV16, Lemma 7.10] to calculate the limit of this cosimplicial cochain complex as the total cochain complex of

$$\check{C}^{n,m}(\ell) := \prod_{\mathcal{O}^{n+1}} \check{C}^m(\ell)(U_0 \cap \dots \cap U_n)$$

The differential in the m -direction simply comes from $\check{C}^*(\ell)$. The differential in the n -direction is the alternating sum of the morphisms $\delta^i : \check{C}^{n,m}(\ell) \rightarrow \check{C}^{n+1,m}(\ell)$, for $0 \leq i \leq n+1$, described in components as follows: Given $\mu \in \check{C}^{n,m}(\ell)$, the (U_0, \dots, U_{n+1}) -component of $\delta^i \mu$ is the $(U_0, \dots, \widehat{U_i}, \dots, U_{n+1})$ -component of μ restricted to $U_0 \cap \dots \cap U_{n+1}$, where the hat under U_i means we're removing it from the tuple.

6. UNFOLDING THE FRACTURE SQUARE OF DELIGNE COHOMOLOGY

REFERENCES

- [Aa10] Camilo Arias Abad and Florian Schätz. The A_∞ de Rham theorem and integration of representations up to homotopy. [arXiv:1011.4693](#), 2010.

- [ADH21] Araminta Amabel, Arun Debray, and Peter J. Haine. Differential cohomology: Categories, characteristic classes, and connections. *arXiv preprint*, 2021. [arXiv:2109.12250](#).
- [BB14] Christian Bär and Christian Becker. *Differential Characters and Geometric Chains*, pages 1–90. Springer International Publishing, Cham, 2014.
- [BG16] Ulrich Bunke and David Gepner. Differential function spectra, the differential Becker-Gottlieb transfer, and applications to differential algebraic K -theory. [arXiv:1306.0247](#), 2016.
- [BNV16] Ulrich Bunke, Thomas Nikolaus, and Michael Völkl. Differential cohomology theories as sheaves of spectra. *J. Homotopy Relat. Struct.*, 11(1):1–66, 2016.
- [Dav24] Jack Davies. V4d2 - Algebraic Topology II So24 (stable and chromatic homotopy theory). *Lecture notes*, 2024. [Unpublished](#).
- [FSS11] Domenico Fiorenza, Urs Schreiber, and Jim Stasheff. Čech cocycles for differential characteristic classes - An ∞ -Lie theoretic construction. [arXiv:1011.4735](#), 2011.
- [HS05] M. J. Hopkins and I. M. Singer. Quadratic functions in geometry, topology, and M-theory. *J. Differential Geom.*, 70(3):329–452, 2005.
- [Lur09] Jacob Lurie. *Higher topos theory*, volume 170 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2009.
- [Lur17] Jacob Lurie. Higher algebra. [Available online](#), September 2017.
- [LV12] Jean-Louis Loday and Bruno Vallette. *Algebraic operads*, volume 346. Springer Science & Business Media, 2012.
- [Zuo21] Wang Zuoqin. Lecture 11 : Approximations and Deformations. *Lecture notes*, 2021. [Unpublished](#).