

DIFFERENTIAL COHOMOLOGY SEMINAR 2

TALK BY MATTHIAS FRERICHs

In this lecture we want to learn the basics of ∞ -category theory. For the ∞ -categorical background, we broadly follow [Gro10] and a little [Lur09].

1. BASICS ON $(\infty, 1)$ -CATEGORIES

$(\infty, 1)$ -categories have different models that capture its essence. The first model are *quasi-categories*.

Definition 1.1. Given a natural number n , let $\langle n \rangle$ denote the linearly ordered set $\{0, \dots, n\}$. The *simplex category* Δ is the category of finite linearly ordered sets $\langle n \rangle$, for every n , and monotone functions.

Definition 1.2. Given $0 \leq i \leq n$, the *i -face map* is the unique injective map $\delta_n^i : \langle n-1 \rangle \rightarrow \langle n \rangle$ missing i . The *i -degeneracy map* is the unique surjective map $\sigma_n^i : \langle n+1 \rangle \rightarrow \langle n \rangle$ such that i and $i+1$ have the same image.

Theorem 1.3. As a category, Δ is generated from the face and degeneracy maps subject to the simplicial identities (see [GJ99, I.4]).

Definition 1.4. A *simplicial set* is a contravariant functor $X : \Delta^{\text{op}} \rightarrow \text{Set}$. Denote by sSet the category of simplicial sets. The representable simplicial set $\text{Hom}_{\Delta}(-, \langle n \rangle)$ is denoted as Δ^n . The set of *n -simplices*, denoted by X_n , is the image of $\langle n \rangle$. By Yoneda lemma, we identify a n -simplex with the corresponding map of simplicial sets $\Delta^n \rightarrow X$. Let $d_i^n : X_n \rightarrow X_{n-1}$ be the map of sets induced by the i -face map δ_n^i , which we also call *i -face map*.

By only representing the face maps, we can depict a simplicial set as follows:

$$(1.5) \quad X_0 \xleftarrow{\quad} X_1 \xleftarrow{\quad} X_2 \xleftarrow{\quad} \dots$$

∞ -categories are then defined in terms of a lifting condition, for which we need to define horns.

Definition 1.6. Given $i \in \langle n \rangle$, the *i -horn* is the simplicial sub-set $\Lambda_i^n \subseteq \Delta^n$ defined as follows: A monotone map $f : \langle k \rangle \rightarrow \langle n \rangle$ is a *k -simplex* of Λ_i^n if there is $j \neq i$ that does not belong in the image of f .

Definition 1.7. A simplicial set X is called a *quasi-category* if every inner horn $\Lambda_i^n \rightarrow X$, i.e. $0 < i < n$, can be extended to a n -simplex $\Delta^n \rightarrow X$. If every horn $\Lambda_i^n \rightarrow X$ admits an extension, X is called a *∞ -groupoid*.

Remark 1.8. The nerve of a category \mathcal{C} is a quasi-category (one can simply check the horn filling condition directly). Moreover, the nerve is a ∞ -groupoid if and only if \mathcal{C} is a groupoid. The nerve functor $N : \text{Cat} \rightarrow \text{sSet}$ has a left adjoint, the *homotopy category* functor Ho . If X is a quasi-category, $\text{Ho}(X)$ has an explicit construction where objects are the 0-simplices of X and morphisms are homotopy classes of 1-simplices, see [Lan21, 1.2.5] and [Gro10, Proposition 1.15].

Definition 1.9. Let X be a quasi-category, a 1-simplex $f : x \rightarrow y$ is an *equivalence* if there are 2-simplices of the following form:

$$(1.10) \quad \begin{array}{ccc} & x & \\ \nearrow & \Downarrow & \searrow f \\ y & \xrightarrow{\quad} & y \end{array} \quad \begin{array}{ccc} & y & \\ \nearrow f & \Downarrow & \searrow \\ x & \xrightarrow{\quad} & x \end{array}$$

where the bottom 1-simplices are the image of y and x , respectively, under the function $X_0 \rightarrow X_1$ induced by σ_0^0 .

The following theorem shows that Kan complex have the same relation with quasi-categories, as groupoids have with categories.

Theorem 1.11. *A quasi-category X is a ∞ -groupoid if and only if every 1-simplex is an equivalence.*

Remark 1.12. [Gro10] approaches ∞ -groupoids in the other direction, namely defining a ∞ -groupoid as a quasi-category where every 1-simplex is an equivalence, then characterizing ∞ -groupoids as quasi-categories satisfying the horn filling condition for every horn.

A strict model for ∞ -categories is given by simplicially enriched categories.

Definition 1.13. Denote by \mathbf{sCat} the category of simplicially enriched categories.

Remark 1.14. Lurie constructs in [Lur09, 1.1.5.1] a functor $N_\Delta : \mathbf{sCat} \rightarrow \mathbf{sSet}$, called the *homotopy coherent nerve*. Similarly to the ordinary nerve, it admits a left adjoint, denoted \mathfrak{C} and called *rigidification*, see [DS11].

Theorem 1.15. *\mathbf{sSet} and \mathbf{sCat} underlie model structures, called Joyal and Dwyer-Kan-Bergner, respectively, such that fibrant-cofibrant objects are quasi-categories and categories enriched over ∞ -groupoids, respectively, and the adjunction*

$$(1.16) \quad \mathbf{sCat} \xrightleftharpoons[\mathfrak{C}]{N_\Delta} \mathbf{sSet}$$

lifts to a Quillen equivalence.

Definition 1.17. The category $\mathcal{K}an$ of ∞ -groupoids is a self-enriched category, so $\mathcal{S} := N_\Delta(\mathcal{K}an)$ is a quasi-category, namely the *quasi-category of ∞ -groupoids*.

Let \mathcal{Top} be the category of compactly generated topological spaces.

Theorem 1.18. *\mathbf{sSet} and \mathcal{Top} underlie model structures, both called Quillen, such that fibrant-cofibrant are ∞ -groupoids and retracts of cellular complexes, respectively, and the adjunction*

$$(1.19) \quad \mathcal{Top} \xrightleftharpoons[\text{Sing}]{\text{Sing}} \mathbf{sSet}$$

lifts to a Quillen equivalence.

2. BASIC CONSTRUCTIONS

Remark 2.1. Recall that \mathbf{sSet} is self-enriched. More specifically, given simplicial sets X, Y , the simplicial set $\text{Hom}_{\mathbf{sSet}}(X, Y)$ is characterized as follows: A n -simplex is a map of simplicial sets $\Delta^n \times X \rightarrow Y$.

Lemma 2.2. *If X is a simplicial set and \mathcal{C} a ∞ -category, then $\text{Hom}_{\mathbf{sSet}}(X, \mathcal{C})$ is a ∞ -category.*

Definition 2.3. Let \mathcal{C} and \mathcal{D} be ∞ -categories, denote by $\mathcal{F}un(\mathcal{C}, \mathcal{D})$ the *∞ -category of functors $\mathcal{C} \rightarrow \mathcal{D}$* .

3. ACCESSIBLE AND PRESENTABLE CATEGORIES

In general, a limit, resp. colimit, preserving functor need not have a left, resp. right, adjoint. Here we wish to introduce a rather large class of ∞ -categories for which the previous statement holds. We begin by recalling the definition of locally presentable 1-categories. Let κ denote a regular cardinal.

Definition 3.1. A category \mathcal{J} is *κ -filtered* if, for every \mathcal{J} with $< \kappa$ many morphisms, every diagram $\mathcal{J} \rightarrow \mathcal{J}$ has a cocone. A functor $F : \mathcal{C} \rightarrow \mathcal{D}$ is *κ -accessible* if it preserves κ -filtered colimits. Given a category \mathcal{C} , an object X is *κ -compact* if $\text{Hom}_{\mathcal{C}}(X, -) : \mathcal{C} \rightarrow \mathbf{Set}$ is κ -accessible.

Definition 3.2. A category \mathcal{C} is *κ -accessible* if there exists a set $S \subseteq \mathcal{C}_0$ of κ -compact objects that generate \mathcal{C} under κ -filtered colimits. A category is *accessible* if it is κ -accessible, for some regular cardinal κ .

Definition 3.3. A category \mathcal{C} that is accessible and cocomplete is called *locally presentable*.

Theorem 3.4. *Let \mathcal{C} be a category, then \mathcal{C} is locally presentable if and only if there exists a small category S such that the induced functor $\mathcal{C} \rightarrow \mathcal{P}(S)$ is a fully faithful, accessible right adjoint.*

Theorem 3.5. *Let \mathcal{C}, \mathcal{D} be locally presentable categories. A functor $F : \mathcal{C} \rightarrow \mathcal{D}$ is a left, resp. right, adjoint if and only if it preserves colimits, resp. it preserves limits and is accessible.*

Definition 3.6. Let X be a ∞ -category. The *∞ -category of pre-sheaves of spaces* is defined as $\mathcal{P}(X) := \text{Hom}_{\mathbf{sSet}}(X^{\text{op}}, \mathcal{S})$.

Definition of
opposite ∞ -
category, slice
 ∞ -category

Theorem 3.7 (Yoneda). *Given a ∞ -category X , there is a fully faithful functor $y : X \rightarrow \mathcal{P}(X)$, called the Yoneda embedding, such that: Given a cocomplete ∞ -category Y , pre-composition by y induces an equivalence*

$$(3.8) \quad \mathcal{F}\mathrm{un}^{\mathrm{L}}(\mathcal{P}(X), Y) \longrightarrow \mathcal{F}\mathrm{un}(X, Y)$$

where $\mathcal{F}\mathrm{un}^{\mathrm{L}}$ denotes the category of colimit preserving functors.

The definition of accessible category transfers directly to the ∞ -categorical setting.

Theorem 3.9. *A ∞ -category X is locally presentable (cocomplete and accessible) if and only if there is a small sub- ∞ -category S such that $X \rightarrow \mathcal{F}\mathrm{un}(S^{\mathrm{op}}, \mathcal{S})$ is a fully faithful, accessible right adjoint.*

Remark 3.10. In view of [Equation \(3.9\)](#), one can define a category to be *locally presentable* if it is the accessible right localization of a pre-sheaf category for some small ∞ -category S . In particular, every pre-sheaf category is locally presentable.

Theorem 3.11. *Let X, Y be presentable ∞ -categories, then a functor $f : X \rightarrow Y$ is a left, resp. right, adjoint if and only if it preserves colimits, resp. it preserves limits and is accessible.*

4. STABLE ∞ -CATEGORIES AND SPECTRA

We now use the ∞ -categorical framework to study spectra. The study of spectra originates from the study of *stable phenomena*, i.e. patterns appearing after repeated application of the suspension functor $\Sigma : \mathrm{Top}_* \rightarrow \mathrm{Top}_*$.

Example 4.1. Let $(\Sigma, \Omega) : \mathrm{Top}_{*/} \rightarrow \mathrm{Top}_{*/}$ be the suspension-loop adjunction on pointed topological spaces. *Freudenthal Suspension Theorem* states that, if X is a n -connected space, the adjunction unit $X \rightarrow \Omega \Sigma X$ is $2n$ -connected. If X is connected, $S^n \wedge X$ is n -connected, so $\Sigma^n X \rightarrow \Omega \Sigma^{n+1} X$ is $2n$ -connected. In particular, $\pi_i(\Sigma^n X) \rightarrow \pi_{i+1}(\Sigma^{n+1} X)$ is an isomorphism, for all $i < 2n$. The group $\pi_i(\Sigma^n X)$ is denoted $\pi_{n-i}^s(X)$, called the $(n-i)$ -stable homotopy group of X .

More examples?

Definition 4.2. Let \mathcal{C} be an ∞ -category, an object 0 that is both initial and terminal is called *zero object*. A category \mathcal{C} with a zero object is called a *pointed category*.

Example 4.3. Let $1 \in \mathcal{C}$ be a terminal object, then the identity of 1 is the zero object in the slice category $\mathcal{C}_{1/}$ of objects under 1 . In particular, the category $\mathcal{S}_* = \mathcal{S}_{*/}$ of pointed spaces is pointed.

Proposition 4.4. *Let \mathcal{D} be a pointed ∞ -category. Evaluation at the 0-sphere S^0 induces an equivalence*

$$(4.5) \quad \mathcal{F}\mathrm{un}^{\mathrm{L}}(\mathcal{S}_*, \mathcal{C}) \longrightarrow \mathcal{C}$$

We now introduce the notion of a triangle.

Definition 4.6. Let \mathcal{C} be a pointed ∞ -category. A *triangle* in \mathcal{C} is a commutative diagram of the form

$$(4.7) \quad \begin{array}{ccc} X & \longrightarrow & Y \\ \downarrow & & \downarrow \\ 0 & \longrightarrow & Z \end{array}$$

A triangle is *exact*, resp. *coexact*, if it is a pullback, resp. pushout, square.

Definition 4.8. Let \mathcal{C} be a pointed ∞ -category. Denote by \mathcal{C}^{Σ} , resp. \mathcal{C}^{Ω} , the full sub-category of $\mathcal{F}\mathrm{un}(\Delta^1 \times \Delta^1, \mathcal{C})$ of coexact, resp. exact, triangles of the form

$$(4.9) \quad \begin{array}{ccc} X & \longrightarrow & 0 \\ \downarrow & & \downarrow \\ 0 & \longrightarrow & Y \end{array},$$

If \mathcal{C} is finitely cocomplete, resp. complete, for every object X , resp. Y , there is a contractible space of coexact, resp. exact, triangles as [Equation \(4.9\)](#). In particular, $\mathcal{C} \simeq \mathcal{C}^{\Sigma}$ and $\mathcal{C} \simeq \mathcal{C}^{\Omega}$.

Proposition 4.10. *If \mathcal{C} is a finitely complete and cocomplete pointed ∞ -category, define the following functors. Then the functors*

$$(4.11) \quad \Sigma : \mathcal{C} \longrightarrow \mathcal{C}^\Sigma \xrightarrow{\text{ev}_{(1,1)}} \mathcal{C} \quad \Omega : \mathcal{C} \longrightarrow \mathcal{C}^\Omega \xrightarrow{\text{ev}_{(0,0)}} \mathcal{C}$$

are adjoint (Σ is left adjoint to Ω).

Theorem 4.12. *Let \mathcal{C} be a finitely bicomplete pointed ∞ -category. The following are equivalent:*

- (1) *A triangle is exact if and only if it is coexact.*
- (2) *(Σ, Ω) is an adjoint equivalence.*
- (3) *A commutative square is a pullback if and only if it is a pushout.*

Definition 4.13. A finite bicomplete pointed ∞ -category \mathcal{C} satisfying any of the equivalent conditions in [Equation \(4.12\)](#) is called *stable*.

If \mathbb{A} denotes a nice abelian category, there is a *derived ∞ -category* of \mathbb{A} , denoted $\mathcal{D}(\mathbb{A})$ such that $\text{Ho}\mathcal{D}(\mathbb{A}) = D(\mathbb{A})$ is the ordinary derived category of \mathbb{A} , i.e. the localization of chain complex at quasi-isomorphisms. From homological algebra, it is known that $D(\mathbb{A})$ underlies the structure of a triangulated category, which turns out to be the 1-categorical reflection of $\mathcal{D}(\mathbb{A})$ being stable.

Proposition 4.14 ([Lur17, 3.11]). *If \mathcal{C} is a stable ∞ -category, then $\text{Ho}\mathcal{C}$ has a canonical structure of a triangulated category.*

To construct the stabilization of a pointed ∞ -category, there are several approaches, such as reduced excisive functors on S_*^{fin} , the category of pointed, finite spaces, see [Lur17, 1.4.2.8]. Here we consider the more explicit approach using spectrum objects.

Definition 4.15. Let \mathcal{C} be a pointed ∞ -category. A *pre-spectrum object* in \mathcal{C} consists of a functor $E : \mathbb{Z} \times \mathbb{Z} \rightarrow \mathcal{C}$ such that $E(n, m) \simeq 0$, for all $n \neq m$. Denote by $\mathcal{P}\text{Sp}(\mathcal{C})$ the category of pre-spectrum objects. The functor $\Omega^{\infty-n} : \mathcal{P}\text{Sp}(\mathcal{C}) \rightarrow \mathcal{C}$ is defined as evaluation at (n, n) .

For every n , the diagram

$$(4.16) \quad \begin{array}{ccc} E(n, n) & \longrightarrow & 0 \\ \downarrow & & \downarrow \\ 0 & \longrightarrow & E(n+1, n+1) \end{array}$$

determines a pair of adjoint morphisms

$$\alpha_n : \Sigma E(n, n) \rightarrow E(n+1, n+1), \quad \beta_n : E(n, n) \rightarrow \Omega E(n+1, n+1)$$

Definition 4.17. Let \mathcal{C} be a pointed ∞ -category. A *spectrum object* in \mathcal{C} consists of a pre-spectrum object E such that β_n is an equivalence, for all n . Denote by $\text{Sp}(\mathcal{C}) \subseteq \mathcal{P}\text{Sp}(\mathcal{C})$ the full sub-category of spectrum objects.

Theorem 4.18. *Let \mathcal{C} be a presentable pointed ∞ -category, then $\Omega^{\infty-n} : \text{Sp}(\mathcal{C}) \rightarrow \mathcal{C}$ admits a left adjoint $\Sigma^{\infty-n} : \mathcal{C} \rightarrow \text{Sp}(\mathcal{C})$, for every n .*

In particular, $\Sigma^\infty : \mathcal{C} \rightarrow \text{Sp}(\mathcal{C})$ has the following universal property.

Theorem 4.19. *Let \mathcal{C} be a presentable pointed ∞ -category. Given a stable ∞ -category \mathcal{D} , pre-composition by Σ^∞ induces an equivalence*

$$(4.20) \quad \mathcal{F}\text{un}^L(\text{Sp}(\mathcal{C}), \mathcal{D}) \longrightarrow \mathcal{F}\text{un}^L(\mathcal{C}, \mathcal{D})$$

In particular, for $\mathcal{C} = S_$, evaluation at the sphere spectrum $\mathbb{S} = \Sigma^\infty S^0$ induces an equivalence*

$$(4.21) \quad \mathcal{F}\text{un}^L(\text{Sp}(S_*), \mathcal{D}) \longrightarrow \mathcal{D}$$

Definition 4.22. The *∞ -category of spectra* is the category of spectrum objects in pointed spaces.

5. GENERALIZED COHOMOLOGY THEORIES

We shall now use the language of ∞ -categories to reformulate the concept of generalized cohomology theory à-la Eilenberg-Steenrod. In this new context, we recall a representability theorem for cohomology theories by spectrum object.

Remark 5.1. Denote by $\mathrm{Set}^{\mathbb{Z}}$ the category of \mathbb{Z} -indexes families of sets. Given an object S and $n \in \mathbb{Z}$, denote by $\Sigma^n S$ the shifted family $(\Sigma^n S)_i = S_{i-n}$.

Definition 5.2 ([Lur17, 1.4.1.6]). Let \mathcal{C} be a finitely cocomplete pointed ∞ -category, $\Sigma_{\mathcal{C}} : \mathcal{C} \rightarrow \mathcal{C}$ the induced suspension functor. A *generalized cohomology theory* is a functor $H : \mathrm{Ho}\mathcal{C}^{\mathrm{op}} \rightarrow \mathrm{Set}^{\mathbb{Z}}$ together with a natural isomorphism $\partial : \Sigma H \rightarrow H\Sigma_{\mathcal{C}}$ such that:

- H preserves arbitrary products. In particular, $H^n(0)$ is the one-point set. Given an object X , the unique morphism $X \rightarrow 0$ induces an element $* \simeq H^n(0) \rightarrow H^n(X)$, which we denote by 0 .
- Given a coexact triangle $X' \rightarrow X \rightarrow X''$, if $\eta \in H^n(X)$ has image $0 \in H^n(X'')$, then it lies in the image of $H^n(X') \rightarrow H^n(X)$.

Theorem 5.3 ([Lur17, 1.4.1.10]). *Let \mathcal{C} be a nice ∞ -category and (H, ∂) a generalized cohomology theory, then, for every n , the functor H^n is representable by an object $E(n)$.*

The natural isomorphism ∂ translates into an equivalence $E(n) \simeq \Omega E(n+1)$, which is then used to construct a spectrum object representing the cohomology theory H^n , see [Lur17, 1.4.1.11].

Remark 5.4. For $\mathcal{C} = \mathcal{S}_*$, the above definition of cohomology theory reduces to the classical Eilenberg-Steenrod definition. Since \mathcal{S}_* is nice, we thus recover the classical *Brown representability theorem*.

REFERENCES

- [DS11] Daniel Dugger and David I. Spivak. Rigidification of quasi-categories. *Algebr. Geom. Topol.*, 11(1):225–261, 2011.
- [GJ99] Paul G. Goerss and John F. Jardine. *Simplicial homotopy theory*, volume 174 of *Progress in Mathematics*. Birkhäuser Verlag, Basel, 1999.
- [Gro10] Moritz Groth. A short course on ∞ -categories. *arXiv preprint*, 2010. [arXiv:1007.2925](https://arxiv.org/abs/1007.2925).
- [Lan21] Markus Land. *Introduction to ∞ -categories*. Birkhäuser Cham, 2021.
- [Lur09] Jacob Lurie. *Higher topos theory*, volume 170 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2009.
- [Lur17] Jacob Lurie. Higher algebra. [Available online](https://www.math.berkeley.edu/~lurie/), September 2017.