

DIFFERENTIAL COHOMOLOGY SEMINAR 3

TALK BY HANNES BERKENHAGEN

In this lecture we cover the basics of sheaf theory and learn about differential cohomology theories as sheaves of spectra.

1. SHEAVES IN CATEGORY THEORY

Let us review sheaves in classical category theory. A good reference remains [MLM94]. We start with the case of sheaves on a topological space.

Definition 1.1. Let \mathcal{C} be a category. A *pre-sheaf* on \mathcal{C} is a functor $F : \mathcal{C}^{\text{op}} \rightarrow \mathbf{Set}$.

Definition 1.2. Let X be a topological space, we denote by $\mathcal{O}pen_X$ the partial order of open subsets of X , viewed as a category. We call a pre-sheaf on $\mathcal{O}pen_X$ a *pre-sheaf on X* . Given $F : \mathcal{O}pen_X^{\text{op}} \rightarrow \mathbf{Set}$ and $u \in F(U)$, for every $V \subseteq U$, we write $u|_V$ for the image of u under the map $F(U) \rightarrow F(V)$.

Definition 1.3. Let X be a topological space. A pre-sheaf on X is a *sheaf* if, for every open $U \subseteq X$ and open cover $\{U_i \subseteq U\}_i$, the following is an equalizer diagram:

$$F(U) \longrightarrow \prod_i F(U_i) \rightrightarrows \prod_{i,j} F(U_i \cap U_j)$$

Now, we wish to extend the notion of sheaf to pre-sheaves on a generic category \mathcal{C} . To do so, we first generalize the concept of cover, for examples by specifying for each object C a class of families of maps $\{f_i : C_i \rightarrow C\}$ that cover C . Finally, in the sheaf condition, we replace intersections $U_i \cap U_j$ with fiber products (assuming \mathcal{C} has the necessary limits). Then a pre-sheaf $F : \mathcal{C}^{\text{op}} \rightarrow \mathbf{Set}$ is a sheaf if, for every object C and any covering family $\{f_i : C_i \rightarrow C\}_i$, the following is an equalizer diagram:

$$F(C) \longrightarrow \prod_i F(C_i) \rightrightarrows \prod_{i,j} F(C_i \times_C C_j)$$

The above idea leads to the notion of *coverage*, here we will use another language, namely that of *sieves*.

2. SHEAVES VIA GROTHENDIECK TOPOLOGIES

We now generalize from sheaves on a topological space to sheaves on a category via the additional data of a Grothendieck topology. We begin by recalling a few definitions. Denote by y the *Yoneda embedding* $\mathcal{C} \rightarrow \mathcal{P}(\mathcal{C})$, mapping C to the pre-sheaf $y(C) := \text{Hom}_{\mathcal{C}}(-, C) : \mathcal{C}^{\text{op}} \rightarrow \mathbf{Set}$.

Definition 2.1. A *sub-functor* to a pre-sheaf $F : \mathcal{C}^{\text{op}} \rightarrow \mathbf{Set}$ consists of a pre-sheaf $S : \mathcal{C}^{\text{op}} \rightarrow \mathbf{Set}$ such that $S(C) \subseteq F(C)$, for all objects C , and the subset inclusions define a natural transformation $S \rightarrow F$. The last condition is equivalent to the following: Given $f : D \rightarrow C$ and $u \in S(C)$, then $F(f)(u) \in S(D)$.

Definition 2.2. Given a pre-sheaf $F : \mathcal{C}^{\text{op}} \rightarrow \mathbf{Set}$, denote by $\mathcal{C}_{/F}$ the category of pairs (C, c) , consisting of an object C and a morphism of pre-sheaves $y(C) \xrightarrow{c} F$.

Theorem 2.3 (Density Theorem). *F is canonically isomorphic to the colimit of the diagram*

$$(2.4) \quad \mathcal{C}_{/F} \longrightarrow \mathcal{C} \xrightarrow{y} \mathcal{P}(\mathcal{C})$$

where the first functor is the canonical projection.

Definition 2.5. Let \mathcal{C} be a category. A *sieve* on an object C is a sub-functor of $y(C)$, with $y(C)$ being the *maximal sieve*. Given a morphism $f : D \rightarrow C$ and a sieve S on C , denote by $f^*(S)$ the *pullback sieve*, mapping X to the set of arrows $g : X \rightarrow D$ such that $fg \in S(C)$.

Definition 2.6. Let \mathcal{F} be a set of morphisms with codomain $C \in \mathcal{C}$. Define the *sieve generated by \mathcal{F}* as the sieve $S_{\mathcal{F}} \subseteq \text{Hom}_{\mathcal{C}}(-, C)$ mapping D to the set of morphisms $D \rightarrow C$ that factor through some element of \mathcal{F} .

Definition 2.7. Let \mathcal{C} be a category. A *Grothendieck topology* on \mathcal{C} is a collection $J(\mathcal{C})$ of sieves, called *covering sieves*, on every object C in \mathcal{C} , satisfying the following axioms:

- (1) The maximal sieve is a covering sieve.
- (2) Given a covering sieve S and a morphism $f: D \rightarrow C$, the pullback sieve $f^*(S)$ is a covering sieve on D .
- (3) A sieve S on C is a covering sieve if, for some covering sieve S' on C and every map $f \in S'$, the pullback sieve $f^*(S)$ is also a covering sieve.

A *Grothendieck site* is a pair (\mathcal{C}, J) where \mathcal{C} is a category and J is a Grothendieck topology on \mathcal{C} .

Example 2.8. If every sieve is a covering sieve, we obtain the *discrete Grothendieck topology*. If the only covering sieves are the maximal ones, we obtain the *indiscrete Grothendieck topology*.

Example 2.9. Let X be a topological space. A sieve on an object U of $\mathcal{O}pen_X$ corresponds to a collection of open subsets of U . Define a sieve to be a covering sieve on U if the corresponding family of subsets covers U in the classical sense.

Example 2.10. Let $\mathcal{T}op$ be the category of topological spaces and continuous maps. Define a sieve S to be a covering sieve if generated, in the sense of Equation (2.6), by a family \mathcal{O} of jointly surjective, open embeddings.

Example 2.11. Let $\mathcal{M}fd$ be the category of smooth manifolds and smooth maps. We can define a topology in the same way as Equation (2.10), using families of jointly surjective, smooth open embeddings.

Definition 2.12. Let (\mathcal{C}, J) be a site and $\mathcal{D} \subseteq \mathcal{C}$ a full sub-category. The *sub-category Grothendieck topology* on \mathcal{D} assigns to each object $C \in \mathcal{D}$ the collection of sieves $S \subseteq \text{Hom}_{\mathcal{D}}(-, C)$ that are the restriction to \mathcal{D} of covering sieves of C in \mathcal{C} .

Definition 2.13. Let (\mathcal{C}, J) be a site and $C \in \mathcal{C}$. Given $f: D \rightarrow C$, there is a canonical identification $\alpha_f: (\mathcal{C}/C)_{/f} \simeq \mathcal{C}/D$. The *slice Grothendieck topology* on \mathcal{C}/C assigns to each object $f: D \rightarrow C$ the collection of sieves $\mathcal{D} \subseteq (\mathcal{C}/C)_{/f}$ that correspond to covering sieves of D under α_f .

We are now ready to define sheaves on a Grothendieck site. Given a sieve S on C , notice that the category \mathcal{C}/S can be identified with the full sub-category of \mathcal{C}/C spanned by morphisms $f \in S$.

Remark 2.14. A sieve on C is equivalent to the datum of a sub-category $\mathcal{D} \subseteq \mathcal{C}/C$ closed un pre-composition, i.e. if $f \in \mathcal{D}$, then $fg \in \mathcal{D}$, for every g composable with f .

Definition 2.15. Given a pre-sheaf $F: \mathcal{C}^{op} \rightarrow \text{Set}$, denote by $\text{Hom}(\mathcal{C}/S, F)$ the limit of the following diagram:

$$(2.16) \quad (\mathcal{C}/S)^{op} \longrightarrow \mathcal{C}^{op} \xrightarrow{F} \text{Set}$$

The inclusion $\mathcal{C}/S \subseteq \mathcal{C}/C$ induces a natural comparison map $\text{Hom}(\mathcal{C}/C, F) \rightarrow \text{Hom}(\mathcal{C}/S, F)$.

Notice that $\text{Hom}(\mathcal{C}/C, F)$ is the limit of a diagram indexed by the category $(\mathcal{C}/C)^{op}$, which has an initial object, the identity of C , therefore the limit is canonically isomorphic to $F(C)$, the value of the diagram at the initial object.

Definition 2.17. Let (\mathcal{C}, J) be a Grothendieck site. A pre-sheaf $F: \mathcal{C}^{op} \rightarrow \text{Set}$ is a *sheaf* if, for every object C and covering sieve S of C , the canonical morphism

$$F(C) \cong \text{Hom}(\mathcal{C}/C, F) \rightarrow \text{Hom}(\mathcal{C}/S, F)$$

is an isomorphism. Explicitly, an object in $\text{Hom}(\mathcal{C}/S, F)$ consists of an element $x_f \in F(D)$, for every map $f \in S$ (where D is the domain of f), such that $x_{fg} = F(g)(x_f)$, for every f and g composable. The canonical morphism takes $x \in F(C)$ to the tuple with $x_f := F(f)(x)$.

Using [Equation \(2.3\)](#) and Yoneda's lemma, one can construct a canonical isomorphism $\mathrm{Hom}(\mathcal{C}_{/S}, F) \simeq \mathrm{Hom}_{\mathcal{P}(\mathcal{C})}(S, F)$, such that the following diagram commutes

$$(2.18) \quad \begin{array}{ccc} F(C) & \longrightarrow & \mathrm{Hom}(\mathcal{C}_{/S}, F) \\ \downarrow \cong & & \downarrow \cong \\ \mathrm{Hom}_{\mathcal{P}(\mathcal{C})}(y(C), F) & \longrightarrow & \mathrm{Hom}_{\mathcal{P}(\mathcal{C})}(S, F) \end{array}$$

where the bottom map is the restriction map induced by the inclusion $S \subseteq y(C)$. In particular, we can rephrase the sheaf condition as follows:

Corollary 2.19. *A pre-sheaf F on a Grothendieck site (\mathcal{C}, J) is a sheaf if and only if for every object C and covering sieve S of C , the canonical morphism*

$$F(C) \cong \mathrm{Hom}_{\mathcal{P}(\mathcal{C})}(y(C), F) \rightarrow \mathrm{Hom}_{\mathcal{P}(\mathcal{C})}(S, F)$$

is an isomorphism.

3. SHEAVES ON ∞ -CATEGORIES

Now that we know how to define sheaves on 1-categories, we generalize the definition of sheaves to the case of ∞ -categories. Here we largely follow [\[Lur09\]](#). Recall that a sieve on C determines a full sub-category of $\mathcal{C}_{/C}$. Conversely, a full sub-category \mathcal{D} of $\mathcal{C}_{/C}$ determines a sieve if and only if $f \in \mathcal{D}$ implies $fg \in \mathcal{D}$, for every g composable with f .

Definition 3.1. Let \mathcal{C} be an ∞ -category and C an object of \mathcal{C} . A *sieve* on C is a full sub- ∞ -category \mathcal{D} of $\mathcal{C}_{/C}$ such that $f \in \mathcal{D}$ implies $fg \in \mathcal{D}$, for every g composable with f .

A sieve $\mathcal{D} \subseteq \mathcal{C}_{/C}$ induces a sieve $\mathcal{D}' \subseteq \mathrm{Ho}(\mathcal{C})_{/C}$ by taking homotopy classes of 1-morphisms in \mathcal{D} .

Definition 3.2. Let \mathcal{C} be an ∞ -category. A *Grothendieck topology* on \mathcal{C} is a collection $J(C)$ of sieves for every object C , such that it induces a Grothendieck topology on $\mathrm{Ho}(\mathcal{C})$.

Notice, we have the following compatibility observation.

Lemma 3.3. *Let \mathcal{C} be a 1-category. A Grothendieck topology on \mathcal{C} is precisely a Grothendieck topology on \mathcal{C} seen as an ∞ -category.*

Definition 3.4. Let \mathcal{C} be an ∞ -category and C an object of \mathcal{C} . Given a sieve $\mathcal{D} \subseteq \mathcal{C}_{/C}$ and a pre-sheaf $F : \mathcal{C}^{\mathrm{op}} \rightarrow \mathcal{S}$, denote by $\mathrm{Hom}(\mathcal{D}, F)$ the limit of the diagram

$$(3.5) \quad \mathcal{D}^{\mathrm{op}} \longrightarrow \mathcal{C}^{\mathrm{op}} \xrightarrow{F} \mathcal{S}$$

where the first functor is the opposite of the canonical projection $\mathcal{C}_{/C} \rightarrow \mathcal{C}$ restricted to \mathcal{D} . Using the same argument as in the 1-categorical case, there is a comparison map $\mathrm{Hom}(\mathcal{C}_{/C}, F) \rightarrow \mathrm{Hom}(\mathcal{D}, F)$ and a canonical equivalence $\mathrm{Hom}(\mathcal{C}_{/C}, F) \simeq F(C)$.

Definition 3.6. Let \mathcal{C} be an ∞ -category. A pre-sheaf $F : \mathcal{C}^{\mathrm{op}} \rightarrow \mathcal{S}$ is a *sheaf* if for every object C and covering sieve \mathcal{D} of C , the canonical morphism

$$F(C) \simeq \mathrm{Hom}(\mathcal{C}_{/C}, F) \rightarrow \mathrm{Hom}(\mathcal{D}, F)$$

is an equivalence. Denote by $\mathrm{Shv}(\mathcal{C}, J)$ the full sub- ∞ -category of $\mathcal{F}\mathrm{un}(\mathcal{C}^{\mathrm{op}}, \mathcal{S})$ spanned by the sheaves on (\mathcal{C}, J) .

Note we defined sheaves as local objects in a locally presentable ∞ -category. Hence, using the formalism of locally presentable ∞ -categories, we immediately have the following result.

Proposition 3.7. *The ∞ -category of sheaves on $\mathrm{Shv}(\mathcal{C}, J)$ is locally locally presentable.*

4. SHEAVES WITH ARBITRARY VALUES

We started with sheaves valued in sets. Then generalized to ∞ -categorical sheaves valued in spaces. However, we want ∞ -categorical sheaves valued in spectra. Hence the next step is to generalize the values of our ∞ -categorical sheaves. Abstractly we obtain such sheaves via the *tensor product* of locally presentable ∞ -categories, which we review now.

Theorem 4.1 ([Lur17, Proposition 4.8.1.15]). *Let \mathcal{C}, \mathcal{D} be locally presentable ∞ -categories. Then there exists a locally presentable ∞ -category $\mathcal{C} \otimes \mathcal{D}$ together with a functor $F : \mathcal{C} \times \mathcal{D} \rightarrow \mathcal{C} \otimes \mathcal{D}$ such that*

- (1) *The functor F preserves colimits component-wise.*
- (2) *For any locally presentable ∞ -category \mathcal{E} , pre-composition with F induces an equivalence between the sub- ∞ -category $\mathcal{F}\mathrm{un}^L(\mathcal{C} \otimes \mathcal{D}, \mathcal{E}) \subseteq \mathcal{F}\mathrm{un}(\mathcal{C} \otimes \mathcal{D}, \mathcal{E})$ of colimit preserving functors, and the sub- ∞ -category $\mathcal{F}\mathrm{un}^{L,L}(\mathcal{C} \times \mathcal{D}, \mathcal{E}) \subseteq \mathcal{F}\mathrm{un}(\mathcal{C} \times \mathcal{D}, \mathcal{E})$ of functors preserving colimits in each component.*

Theorem 4.2 ([Lur17, Proposition 4.8.1.17]). *For \mathcal{C} and \mathcal{D} locally presentable ∞ -categories, there is a canonical equivalence between $\mathcal{C} \otimes \mathcal{D}$ and $\mathcal{F}\mathrm{un}^R(\mathcal{C}^{\mathrm{op}}, \mathcal{D})$, of limit preserving functors $\mathcal{C}^{\mathrm{op}} \rightarrow \mathcal{D}$.*

Definition 4.3. Let (\mathcal{C}, J) be \mathcal{C} be a Grothendieck site, and \mathcal{D} be a locally presentable ∞ -category. A *sheaf* on (\mathcal{C}, J) with values in \mathcal{D} is a limit preserving functor $F : \mathrm{Shv}(\mathcal{C}, J)^{\mathrm{op}} \rightarrow \mathcal{D}$, which corresponds to an object in $\mathrm{Shv}(\mathcal{C}, J) \otimes \mathcal{D}$.

Of course, this description is very abstract and ideally we want a more explicit description that we can use in the future. That is the aim of the next sections.

5. AN EXPLICIT DESCRIPTION OF SHEAVES VALUED IN SPECTRA

We have given a formal definition of sheaves valued in spectra, via the tensor product of locally presentable ∞ -categories. We now want a more explicit description thereof. For this we make use of the following result.

Proposition 5.1 ([Lur17], see also [GGN15]). *The inclusion $\mathcal{P}\mathrm{r}_{\mathrm{st}}^L \rightarrow \mathcal{P}\mathrm{r}^L$ of stable locally presentable ∞ -categories into the category of locally presentable ∞ -categories admits a left adjoint, which is explicitly given by tensoring with Sp .*

We can recall from our previous talk that the stabilization of a locally presentable ∞ -category \mathcal{C} is given by the ∞ -category $\mathrm{Sp}(\mathcal{C})$ of spectrum objects in \mathcal{C} , therefore $\mathcal{C} \otimes \mathrm{Sp} \simeq \mathrm{Sp}(\mathcal{C})$, for every locally presentable ∞ -category \mathcal{C} . In particular, the ∞ -category $\mathrm{Shv}(\mathcal{C}, J) \otimes \mathrm{Sp}$ is equivalent to the category of spectrum objects in $\mathrm{Shv}(\mathcal{C}, J)$. Finally, since the inclusion $\mathrm{Shv}(\mathcal{C}, J) \rightarrow \mathcal{F}\mathrm{un}(\mathcal{C}^{\mathrm{op}}, \mathcal{S})$ preserves and reflects limits, a commutative square in $\mathrm{Shv}(\mathcal{C}, J)$ is a pullback square if and only if it is a pullback square in $\mathcal{F}\mathrm{un}(\mathcal{C}^{\mathrm{op}}, \mathcal{S})$.

Proposition 5.2. *A spectrum object in sheaves on the Grothendieck site (\mathcal{C}, J) is given by a spectrum object in pre-sheaves on (\mathcal{C}, J) such that each component is a sheaf.*

This gives us the following explicit description of sheaves valued in spectra. Recall that $\Omega^{\infty-n} : \mathrm{Sp} \rightarrow \mathcal{S}$ is the functor mapping a spectrum object into its n -th component.

Theorem 5.3. *There is a canonical equivalence between $\mathrm{Shv}(\mathcal{C}, J) \otimes \mathrm{Sp}$ and the category of functors $F : \mathcal{C}^{\mathrm{op}} \rightarrow \mathrm{Sp}$ such that $\Omega^{\infty-n} F : \mathcal{C}^{\mathrm{op}} \rightarrow \mathcal{S}$ is a sheaf of spaces, for every n .*

6. AN EXPLICIT DESCRIPTION OF SHEAVES ON THE SITE OF MANIFOLDS

We continue our analysis of sheaves with the aim of providing explicit descriptions. Now we focus on the case of sheaves on the site of manifolds. Recall that a sieve S on a manifold M is a covering sieve if an open cover \mathcal{O} of M exists, such that every morphism in S factors through the inclusion $U \subseteq M$ of some element $U \in \mathcal{O}$. From the definition of the topology on $\mathcal{M}\mathrm{fd}$ we have the following: Given a covering sieve S generated by an open cover \mathcal{O} . Denote by $\mathcal{I}(\mathcal{O})$ the poset (viewed as a category) consisting of finite intersections of elements of \mathcal{O} .

Lemma 6.1. *The functor $i : \mathcal{I}(\mathcal{O}) \rightarrow \mathcal{M}\mathrm{fd}_{/S}$ is cofinal.*

In particular, since $\mathrm{Hom}(\mathcal{M}\mathrm{fd}_{/S}, F)$ is defined as the limit $(\mathcal{M}\mathrm{fd}_{/S})^{\mathrm{op}} \rightarrow \mathcal{M}\mathrm{fd}^{\mathrm{op}} \xrightarrow{F} \mathcal{S}$ and i^{op} is final, there is a canonical equivalence

$$\mathrm{Hom}(\mathcal{M}\mathrm{fd}_{/S}, F) \simeq \lim(\mathcal{I}(\mathcal{O})^{\mathrm{op}} \rightarrow \mathcal{M}\mathrm{fd}^{\mathrm{op}} \xrightarrow{F} \mathcal{S}) = \lim_{U \in \mathcal{I}(\mathcal{O})} F(U)$$

Proposition 6.2. *A pre-sheaf $F: \mathcal{Mfd}^{\text{op}} \rightarrow \mathcal{S}\mathcal{P}$ is a sheaf if and only if for every manifold M and every open cover \mathcal{O} of M , the canonical morphism $F(M) \rightarrow \lim_{U \in \mathcal{I}(\mathcal{O})} F(U)$ is an equivalence.*

In fact we can check the sheaf condition for sheaves on the site of manifolds with a very specific type of covering families. For a proof of the following result, we refer to [ADH21, Proposition 3.6.6].

Proposition 6.3. *A pre-sheaf $F: \mathcal{Mfd}^{\text{op}} \rightarrow \mathcal{S}\mathcal{P}$ is a sheaf if and only if*

- (1) $F(\emptyset)$ is terminal.
- (2) For every $M = U \cup V$, the canonical square

$$\begin{array}{ccc} F(M) & \longrightarrow & F(U) \\ \downarrow & & \downarrow \\ F(V) & \longrightarrow & F(U \cap V) \end{array}$$

is a pullback square.

- (3) For a sequential family of opens $U_1 \subseteq U_2 \subseteq \dots$ covering M , the canonical map $F(M) \rightarrow \lim_n F(U_n)$ is an equivalence.

7. ENOUGH POINTS AND EQUIVALENCES OF SHEAVES

We now have established a very solid understanding of sheaves on the site of manifolds with arbitrary values as pre-sheaves with suitable limit conditions. We now want to use these explicit descriptions to better understand equivalences of sheaves on the site of manifolds.

Definition 7.1 ([Lur09, Remark 6.5.4.8]). Given a ∞ -topos \mathcal{X} , a *point* of \mathcal{X} is a functor $p: \mathcal{S} \rightarrow \mathcal{X}$ admitting a left exact left adjoint p^* . We say \mathcal{X} has *enough points* if the following holds: Given a morphism f , if $p^*(f)$ is an equivalence for every point p , then f is an equivalence.

Here we are mainly interested in ∞ -topoi of sheaves, i.e. $\mathcal{X} = \mathcal{S}\mathcal{h}\mathcal{v}(\mathcal{C})$, for some site \mathcal{C} .

Remark 7.2. A functor $p: \mathcal{S} \rightarrow \mathcal{X}$ admitting a left exact left adjoint is called a *geometric morphism*, see [Lur09, Definition 6.3.1.1].

Definition 7.3. Let X be a topological space and \mathcal{C} a locally presentable category. Given a point $x \in X$, denote by x^* the functor $\mathcal{S}\mathcal{h}\mathcal{v}(X, \mathcal{C}) \rightarrow \mathcal{C}$ given by $x^*(F) = \text{colim}_{x \in U} F(U)$. The functor x^* is called *stalk at x* and is left adjoint to the *skyscraper at x* functor, denoted as x_* and given by $x_*(S)(U) = S$, if $x \in U$, and $= 1$, otherwise (where $1 \in \mathcal{C}$ is the terminal object).

Remark 7.4. Using that filtered colimits commute with finite limits (see [Lur09, Example 7.3.4.7]), one can prove that x^* is left exact and x_* is a point of $\mathcal{S}\mathcal{h}\mathcal{v}(X)$ according to Equation (7.1). In particular, every point $x \in X$ induces a point of $\mathcal{S}\mathcal{h}\mathcal{v}(X)$.

Theorem 7.5. *Let X be a smooth manifold, then $\mathcal{S}\mathcal{h}\mathcal{v}(X)$ has enough points.*

Proof. [Lur09, Corollary 7.2.1.17] and [Lur09, Theorem 7.2.3.6] reduce the claim to X having finite Lebesgue's covering dimension (see [Lur09, Definition 7.2.3.1]). The Fundamental Urysohn's Identity states that, for separable metric spaces (such as smooth paracompact manifolds), Lebesgue's covering dimension coincides with the *small inductive dimension*, or SID, for short. The SID of a space X , denoted by $\text{ind}X$, is defined inductively as follows:

- (1) $\text{ind}\emptyset = -1$.
- (2) $\text{Ind}X = n$ is the smallest natural number such that every point has a neighborhood basis of open sets U such that $\text{ind}\partial U \leq n - 1$. If no such natural number n exists, we set $\text{ind}X = \infty$.

One can check by induction that $\text{ind}X = n$, for every n -dimensional topological manifold X . □

Remark 7.6. The proof of [Lur09, Corollary 7.2.1.17] actually shows that the collection of points $x_*: \mathcal{S} \rightarrow \mathcal{S}\mathcal{h}\mathcal{v}(X)$, for all $x \in X$, is enough to detect equivalences, namely f is an equivalence if $x^*(f)$ is an equivalence, for every $x \in X$.

Theorem 7.7. *The ∞ -category $\mathcal{S}\mathcal{h}\mathcal{v}(\mathcal{Mfd})$ has enough points.*

Proof. Let M be a smooth manifold, there is a faithful functor $\mathcal{O}pen_M \rightarrow \mathcal{M}fd$ that is a morphism of sites (see [Pst22, Definition A.10]) and satisfies the *covering lifting property* (see [Pst22, Definition A.12]), therefore the restriction functor $-|_M : \mathcal{S}hv(\mathcal{M}fd) \rightarrow \mathcal{S}hv(M)$ is a left adjoint (see [Pst22, Proposition A.13]) and a right adjoint (hence left exact). In particular, a point $p : \mathcal{S} \rightarrow \mathcal{S}hv(M)$, for any manifold M , induces a point of $\mathcal{S}hv(\mathcal{M}fd)$ as the composition of $p : \mathcal{S} \rightarrow \mathcal{S}hv(M)$ with the right adjoint $\mathcal{S}hv(M) \rightarrow \mathcal{S}hv(\mathcal{M}fd)$. The left adjoint is $\mathcal{S}hv(\mathcal{M}fd) \xrightarrow{-|_M} \mathcal{S}hv(M) \xrightarrow{p^*} \mathcal{S}$.

Let f be a morphism such that $p^*(f)$ is an equivalence for every point p of $\mathcal{S}hv(\mathcal{M}fd)$. In particular, $p^*(f|_M)$ is an equivalence for every point of $\mathcal{S}hv(M)$, hence $f|_M$ is an equivalence in $\mathcal{S}hv(M)$, by [Equation \(7.5\)](#). Finally, if $f|_M$ is an equivalence, for every M , then f is an equivalence. \square

The proof of [Equation \(7.7\)](#) together with [Equation \(7.6\)](#) implies the following:

Corollary 7.8. *A morphism f in $\mathcal{S}hv(\mathcal{M}fd)$ is an equivalence if and only if for every manifold M and every point $x \in M$, the map on stalks $x^*(f)$ is an equivalence.*

Note, for every n -manifold M and $x \in M$, there is an open embedding $x' : \mathbb{R}^n \rightarrow M$ such that $x'(0) = x$. The induced map $\mathcal{O}pen_{\mathbb{R}^n, 0}^{\text{op}} \rightarrow \mathcal{O}pen_{M, x}^{\text{op}}$ is final (since $\{x'(U) | 0 \in U \subseteq \mathbb{R}^n\}$ forms a neighborhood basis for x), therefore it induces a natural equivalence $0^* \simeq x^*$. Denote by 0_n the origin of \mathbb{R}^n , the previous argument together with [Equation \(7.8\)](#) implies the following:

Corollary 7.9. *A morphism f of sheaves on $\mathcal{M}fd$ is an equivalence if and only if for all $n \geq 0$, the induced map on stalks $(0_n)^*(f)$ is an equivalence.*

8. SHEAVES OF MANIFOLDS VIA THE EUCLIDEAN SITE

According to [Equation \(2.12\)](#), every full sub-category $\mathcal{B} \subseteq \mathcal{C}$ of a site \mathcal{C} inherits a Grothendieck topology. If \mathcal{B} is *dense* (see [Joh02, Definition 2.2.1]) then the restriction functor $\mathcal{S}hv(\mathcal{C}, \text{Set}) \rightarrow \mathcal{S}hv(\mathcal{B}, \text{Set})$ is an equivalence, this is the *comparison lemma* ([Joh02, Theorem 2.2.3]). However, this is no longer true with sheaves valued in ∞ -categories, i.e. a dense sub-site $\mathcal{B} \subseteq \mathcal{C}$ does not induce an equivalence between ∞ -categories of sheaves (see [Lur18, Counterexample 20.4.0.1]). But, the comparison lemma from ∞ -sheaves *does* hold for $\mathcal{C} = \mathcal{M}fd$ because of *hypercompleteness*, as we'll show. This comparison lemma allows us to relate statements about sheaves on dense sub-categories of $\mathcal{M}fd$ to statements about sheaves on all manifolds.

Let \mathcal{C} be a 1-site and $\mathcal{P}_\Delta(\mathcal{C}) = \mathcal{F}un(\mathcal{C}^{\text{op}}, \text{sSet})$ the ordinary category of simplicial pre-sheaves on \mathcal{C} . Given a pre-sheaf $F : \mathcal{C}^{\text{op}} \rightarrow \text{Set}$, denote by F_δ the simplicial pre-sheaf mapping C to the constant simplicial set with value $F(C)$.

Definition 8.1. Given a morphism of pre-sheaves of sets $f : F \rightarrow \text{Hom}_{\mathcal{C}}(-, C)$, the *image* of f is the sieve S_f mapping D to the image of the map of sets $F(D) \rightarrow \text{Hom}_{\mathcal{C}}(D, C)$. If $C \in \mathcal{C}$ and \mathcal{C} is a site, we call f a *local epimorphism* if S_f is a covering sieve of C . A morphism of pre-sheaves $F \rightarrow G$ is a local epimorphism if, for every $y(C) \rightarrow G$, the projection $y(C) \times_G F \rightarrow y(C)$ is a local epimorphism.

Definition 8.2 ([DHI03, Lemma 4.9]). Given $C \in \mathcal{C}$, an *hypercover* of C consists of a simplicial pre-sheaf $U : \mathcal{C}^{\text{op}} \rightarrow \text{sSet}$ together with a morphism $U \rightarrow y(C)_\delta$, called *augmentation*, such that the maps $U_0 \rightarrow y(C)$ and $U_n \simeq \text{Hom}_{\text{sSet}}(\Delta^n, U) \rightarrow \text{Hom}_{\text{sSet}}(\partial\Delta^n, U)$, for every n , are local epimorphisms, see [Equation \(8.1\)](#). An hypercover has *height* k if $U_n \rightarrow \text{Hom}_{\text{sSet}}(\partial\Delta^n, U)$ is an isomorphism, for all $n \geq k$.

Remark 8.3. An hypercover has height 0 if and only if it is the Čech nerve of the morphism $U_0 \rightarrow y(C)$.

Definition 8.4. Denote by $\widehat{\mathcal{S}hv}(\mathcal{C}) \subseteq \mathcal{S}hv(\mathcal{C})$ the full sub- ∞ -category of sheaves satisfying *hyperdescent*, meaning that, for every object C and hypercover U of C , the augmentation induced an equivalence

$$F(C) \simeq \text{Hom}_{\mathcal{P}(\mathcal{C})}(y(C), F) \rightarrow \lim_n \text{Hom}_{\mathcal{P}(\mathcal{C})}(U_n, F)$$

$\mathcal{S}hv(\mathcal{C})$ is called *hypercomplete* if every sheaf satisfies hyperdescent.

Definition 8.5 ([BGH20, Definition 3.12.2]). Let \mathcal{C} be a site. A full sub-category $\mathcal{B} \subseteq \mathcal{C}$ is *dense* if for every object $C \in \mathcal{C}$ there is a family \mathcal{F} of morphisms with codomain C and domain in \mathcal{B} , such that $S_{\mathcal{F}}$ (see [Equation \(2.6\)](#)) is a covering sieve.

Theorem 8.6. *$\mathcal{S}hv(\mathcal{M}fd)$ is hypercomplete. If $i : \mathcal{D} \subseteq \mathcal{M}fd$ is a full, dense sub-site, then $\mathcal{S}hv(\mathcal{D})$ is also hypercomplete. Moreover, the adjunction $(i_*, i^*) : \mathcal{S}hv(\mathcal{D}) \rightarrow \mathcal{S}hv(\mathcal{M}fd)$ is an equivalence.*

Proof. Hypercompleteness of $\mathrm{Shv}(\mathcal{M}\mathrm{fd})$ follows from [Equation \(7.7\)](#), since having enough points implies hypercompleteness, see [\[BGH20, Example 3.11.10\]](#). The second and third claims are a direct consequence of [\[BGH20, Corollary 3.12.13\]](#) and the first claim. By [\[BGH20, Proposition 3.12.11\]](#), pre-sheaves restriction i^* preserves sheaves satisfying hyperdescent. \square

Recall that, given a locally presentable category \mathcal{E} , there is a canonical equivalence $\mathrm{Shv}(\mathcal{C}, \mathcal{E}) \simeq \mathrm{Shv}(\mathcal{C}) \otimes \mathcal{E}$. In particular, the following is a simple corollary of [Equation \(8.6\)](#):

Corollary 8.7. *For every locally presentable category \mathcal{E} and full, dense sub-site $\mathcal{D} \subseteq \mathcal{M}\mathrm{fd}$, the restriction functor $\mathrm{Shv}(\mathcal{M}\mathrm{fd}, \mathcal{E}) \rightarrow \mathrm{Shv}(\mathcal{D}, \mathcal{E})$ is an equivalence.*

We will mainly focus on the following sub-site of $\mathcal{M}\mathrm{fd}$.

Definition 8.8. Denote by $\mathcal{E}\mathrm{uc}$ the full sub-category of $\mathcal{M}\mathrm{fd}$ generated by Euclidean manifolds, i.e. \mathbb{R}^n , for every n . $\mathcal{E}\mathrm{uc} \subseteq \mathcal{M}\mathrm{fd}$ is dense in the sense of [Equation \(8.5\)](#).

Related to the notion of hypercompleteness is the notion of *Postnikov completeness*. A space S is *n-truncated*, $n \geq 0$, if $\pi_i(S) \simeq 0$, for all $i > n$. We say S is *(-1)-truncated* if S is either empty or contractible. Denote by $\tau_{\leq n} \mathcal{S} \subseteq \mathcal{S}$ the full sub-category of *n-truncated* spaces. The inclusion functor admits a left adjoint $\tau_{\leq n}$, and so every space has a natural map $S \rightarrow \tau_{\leq n} S$. A classical result of algebraic geometry is that $S \simeq \lim_n \tau_{\leq n} S$. The notion of truncated object makes sense in a general ∞ -category.

Definition 8.9 ([\[Lur09, Definition 5.5.6.1\]](#)). Let $n \geq -1$, an object $C \in \mathcal{C}$ is *n-truncated* if $\mathrm{Hom}_{\mathcal{C}}(D, C)$ is a *n-truncated* space, for every $D \in \mathcal{C}$. Denote by $\tau_{\leq n} \mathcal{C}$ the full sub-category of *n-truncated* objects. A morphism $f : C \rightarrow D$ *exhibit D as the n-truncation of C* if D is *n-truncated* and the induced morphism $\tau_{\leq n} C \rightarrow D$ is an equivalence.

If \mathcal{C} is a locally presentable ∞ -category, $\tau_{\leq n} \mathcal{C}$ can be characterized as a class of local objects (see [\[Lur09, Proposition 5.5.6.18\]](#)). In particular, the inclusion functor $\tau_{\leq n} \mathcal{C} \subseteq \mathcal{C}$ admits a left adjoint $\tau_{\leq n} : \mathcal{C} \rightarrow \tau_{\leq n} \mathcal{C}$. Given an object $C \in \mathcal{C}$, its truncations form a sequence

$$(8.10) \quad \cdots \rightarrow \tau_{\leq 2} C \rightarrow \tau_{\leq 1} C \rightarrow \tau_{\leq 0} C$$

where $\tau_{\leq n+1} C \rightarrow \tau_{\leq n} C$ is the $(n+1)$ -truncation of the map $C \rightarrow \tau_{\leq n} C$ together with the fact that $\tau_{\leq n+1} E \simeq E$ naturally, for every *n-truncated* E . Let $\mathbb{Z}_{\geq 0}$ be the partial order of natural numbers.

Definition 8.11 ([\[Lur18, Definition A.7.2.1\]](#) and [\[Lur09, Definition 5.5.6.23\]](#)). A *Postnikov pre-tower* is a functor $X : \mathbb{Z}_{\geq 0}^{\mathrm{op}} \rightarrow \mathcal{C}$ such that $X_{n+1} \rightarrow X_n$ exhibits X_n as the *n-truncation* of X_{n+1} , for every $n \geq 0$. Denote by $\mathrm{Pos}(\mathcal{C})$ the category of Postnikov pre-towers, then there is a comparison functor $\tau : \mathcal{C} \rightarrow \mathrm{Pos}(\mathcal{C})$. We say \mathcal{C} is *Postnikov complete* if τ is an equivalence.

One can show that τ is left left adjoint to the limit functor $\lim : \mathcal{F}\mathrm{un}(\mathbb{Z}_{\geq 0}^{\mathrm{op}}, \mathcal{C}) \rightarrow \mathcal{C}$ restricted to Postnikov pre-towers. In particular, if \mathcal{C} is Postnikov complete, then $C \simeq \lim_n \tau_{\leq n} C$. Let $\mathcal{X} = \mathrm{Shv}(\mathcal{C})$, for some site \mathcal{C} , and $\widehat{\mathcal{X}} = \widehat{\mathrm{Shv}(\mathcal{C})}$, then $\tau_{\leq n} \mathcal{X} \subseteq \widehat{\mathcal{X}}$, for every n , see [\[Lur09, Lemma 6.5.2.9\]](#). In particular, if \mathcal{X} is Postnikov complete, then every object C is equivalent to the limit of its truncations, which implies C is hypercomplete (Since $\widehat{\mathcal{X}}$ is a category of local objects in \mathcal{X} , it is closed under limits in \mathcal{X}).

Theorem 8.12. *If $\mathrm{Shv}(\mathcal{C})$ is Postnikov complete, then it is hypercomplete.*

REFERENCES

- [ADH21] Araminta Amabel, Arun Debray, and Peter J. Haine. Differential cohomology: Categories, characteristic classes, and connections. *arXiv preprint*, 2021. [arXiv:2109.12250](#).
- [BGH20] Clark Barwick, Saul Glasman, and Peter Haine. Exodromy. [arXiv:1807.03281](#), 2020.
- [DHI03] Daniel Dugger, Sharon Hollander, and Daniel C. Isaksen. Hypercovers and simplicial presheaves. [arXiv:math/0205027](#), 2003.
- [GGN15] David Gepner, Moritz Groth, and Thomas Nikolaus. Universality of multiplicative infinite loop space machines. *Algebr. Geom. Topol.*, 15(6):3107–3153, 2015.
- [Joh02] Peter T. Johnstone. *Sketches of an elephant: a topos theory compendium. Vol. 1*, volume 43 of *Oxford Logic Guides*. The Clarendon Press, Oxford University Press, New York, 2002.
- [Lur09] Jacob Lurie. *Higher topos theory*, volume 170 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2009.
- [Lur17] Jacob Lurie. Higher algebra. [Available online](#), September 2017.

- [Lur18] Jacob Lurie. Spectral algebraic geometry. <https://www.math.ias.edu/~lurie/papers/SAG-rootfile.pdf>, February 2018.
- [MLM94] Saunders Mac Lane and Ieke Moerdijk. *Sheaves in geometry and logic*. Universitext. Springer-Verlag, New York, 1994.
A first introduction to topos theory, Corrected reprint of the 1992 edition.
- [Pst22] Piotr Pstragowski. Synthetic spectra and the cellular motivic category. [arXiv:1803.01804](https://arxiv.org/abs/1803.01804), 2022.