

Astronomía Avanzada I (Semester 1 2025)

## **Stellar Atmospheres (3)**

Radiative Transfer Effects

**Nina Hernitschek**

Centro de Astronomía CITEVA

Universidad de Antofagasta

May 30, 2025

# Recap: Flux and Brightness

The **(specific) intensity**  $I_\lambda$  is a measure of brightness:

$$I_\lambda = \frac{E_\lambda}{\cos \theta \, d\lambda \, d\sigma \, d\omega \, dt}$$

The **flux**  $F_\lambda$  is the projection of the specific intensity in the radial direction, integrated over all solid angles:

$$F_\lambda = \oint I_\lambda \cos \theta \, d\omega$$

$I_\lambda$  is independent of the distance from the source.  $F_\lambda$  obeys the inverse square law.

# Recap: Flux and Brightness

Stellar  
Atmospheres  
(3)

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

Flux is related to the **intensity**:

The flux  $F_\lambda$  is a measure of the net energy flow across an area  $d\sigma$ , over a time  $dt$ , in a  $d\lambda$ . The only directional significance is whether the energy crosses  $d\sigma$  from the top or from the bottom.

Then we can write:

$$\begin{aligned} F_\lambda &= \frac{\oint dE_\lambda}{d\lambda d\sigma dt} \\ &= \oint I_\lambda \cos \theta d\omega \end{aligned}$$

The solid angle  $d\omega$  appears for  $I_\lambda$  but not for  $F_\lambda$ .

# Mean Intensity and Energy Density

Stellar  
Atmospheres  
(3)

## The mean intensity

$$J_{\lambda} = \frac{1}{4\pi} \oint I_{\lambda} d\Omega$$

is related to the energy density  $u_{\lambda}$ :

Energy radiated through area element  $d\sigma$  during time  $dt$ :


$$dE_{\lambda} = I_{\lambda} d\lambda d\sigma d\Omega dt$$

with  $l = c dt$ :  $dV = l d\sigma = c dt d\sigma$

Hence, the energy contained in the volume element  $dV$  per wavelength interval, the **energy density**, is

$$u_{\lambda} dV d\lambda = \oint I_{\lambda} d\Omega d\lambda d\sigma dt = 4\pi J_{\lambda} \frac{dV}{c} d\lambda$$

$$u_{\lambda} = \frac{4\pi}{c} J_{\lambda} \left[ \frac{\text{erg}}{\text{cm}^3 \text{ \AA}} \right]$$



Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

# Radiation Pressure

Stellar  
Atmospheres  
(3)

Because a photon possesses an energy  $E$ , it also carries a **momentum**  $p_\lambda = E_\lambda/c$  and can exert a **radiation pressure**:

Consider photons transferring momentum to a solid wall. We get the force:


$$F = \frac{dp_{\lambda,\perp}}{dt} = \frac{1}{c} \frac{dE_\lambda}{dt} \cos \theta$$

The pressure is:

$$dP_\lambda = \frac{F}{d\sigma} = \frac{1}{c} \frac{dE_\lambda}{dt} \frac{\cos \theta}{d\sigma} = \frac{1}{c} I_\lambda \cos^2 \theta d\Omega d\lambda$$

with

$$I_\lambda = \frac{dE_\lambda}{\cos \theta d\lambda d\sigma d\Omega dt}$$



Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limbo  
Darkening

Eddington  
Approximation

Summary

# Radiation Pressure

The change in the  $z$  component of the momentum of photons with wavelengths between  $\lambda$  and  $\lambda + d\lambda$  that are reflected from the area  $dA$  in a time interval  $dt$  is

$$\begin{aligned}\Delta p_{\lambda} d\lambda &= [p_{\lambda,z}^{\text{final}} - p_{\lambda,z}^{\text{initial}}] d\lambda \\ &= \left[ \frac{E_{\lambda} \cos \theta}{c} - \left( -\frac{E_{\lambda} \cos \theta}{c} \right) \right] d\lambda \\ &= \frac{2E_{\lambda} \cos \theta}{c} d\lambda \\ &= \frac{2}{c} I_{\lambda} d\lambda dt dA \cos^2 \theta d\Omega\end{aligned}$$

Stellar  
Atmospheres  
(3)

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

# Radiation Pressure

From Newton's 2nd and 3rd laws, the radiation pressure exerted by these photons with wavelengths between  $\lambda$  and  $\lambda + d\lambda$  in the case of reflection:

$$P_{\lambda}^{\text{rad}} = \frac{2}{c} \int_{\text{hemisphere}} I_{\lambda} d\lambda \cos^2 \theta d\Omega = \frac{2}{c} \int_{\phi=0}^{2\pi} \int_{\theta=0}^{\pi/2} I_{\lambda} d\lambda \cos^2 \theta \sin \theta d\theta d\phi$$

Stellar  
Atmospheres  
(3)

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

# Radiation Pressure

Stellar  
Atmospheres  
(3)

From Newton's 2nd and 3rd laws, the radiation pressure exerted by these photons with wavelengths between  $\lambda$  and  $\lambda + d\lambda$  in the case of reflection:

$$P_{\lambda}^{\text{rad}} = \frac{2}{c} \int_{\text{hemisphere}} I_{\lambda} d\lambda \cos^2 \theta d\Omega = \frac{2}{c} \int_{\phi=0}^{2\pi} \int_{\theta=0}^{\pi/2} I_{\lambda} d\lambda \cos^2 \theta \sin \theta d\theta d\phi$$

Recap  
Radiative  
Transfer (II)  
Grey  
Atmosphere

Imagine removing the reflecting surface  $dA$  and replacing it with a mathematical surface. Instead of being reflected, photons will be streaming through  $dA$  from the other side. We have:

Limb  
Darkening  
Eddington  
Approximation

$$\begin{aligned} P_{\lambda}^{\text{rad}} d\lambda &= \frac{1}{c} \int I_{\lambda} d\lambda \cos^2 \theta d\Omega \quad (\text{transmission}) \\ &= \frac{1}{c} \int_{\pi=0}^{2\pi} \int_{\theta=0}^{\pi} I_{\lambda} d\lambda \cos^2 \theta \sin \theta d\theta d\phi \\ &= \frac{4\pi}{3c} I_{\lambda} d\lambda \quad (\text{isotropic radiation field}) \end{aligned}$$

Summary



# Radiation Pressure

The total radiation pressure produced by photons with all wavelengths is

$$P_{\text{rad}} = \int_0^{\infty} P_{\lambda}^{\text{rad}} d\lambda.$$

Stellar  
Atmospheres  
(3)

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

# Radiation Pressure

The total radiation pressure produced by photons with all wavelengths is

$$P_{\text{rad}} = \int_0^{\infty} P_{\lambda}^{\text{rad}} d\lambda.$$

For blackbody radiation, the total radiation pressure can be expressed as

$$\begin{aligned} P_{\text{rad}} &= \frac{4\pi}{3c} \int_0^{\infty} B_{\lambda}(T) d\lambda \\ &= \frac{4\sigma_{\text{SB}} T^4}{3c} = \frac{1}{3} u \end{aligned}$$

Stellar  
Atmospheres  
(3)

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

# Radiation Pressure Gradient

Despite photon's *random walk* journey to the stellar surface, the energy from the deep interior of the star manages to escape into space.

The temperature in the stellar atmosphere decreases outward, the radiation pressure is smaller at larger radii. This gradient in the radiation pressure causes a **net flow of radiation energy moving outward** from the center of the star and eventually emerging at the stellar surface. This process can be described by

$$\frac{dP_{\text{rad}}}{dr} = -\frac{\rho \bar{\kappa}}{c} F_{\text{rad}}$$

Stellar  
Atmospheres  
(3)

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

# Radiation Pressure Gradient

Despite photon's *random walk* journey to the stellar surface, the energy from the deep interior of the star manages to escape into space.

The temperature in the stellar atmosphere decreases outward, the radiation pressure is smaller at larger radii. This gradient in the radiation pressure causes a **net flow of radiation energy moving outward** from the center of the star and eventually emerging at the stellar surface. This process can be described by

$$\frac{dP_{\text{rad}}}{dr} = -\frac{\rho \bar{\kappa}}{c} F_{\text{rad}}$$

The transfer of radiation energy is a subtle process involving the slow outward diffusion of randomly walking photons, drifting towards the surface in response to small differences in the radiation pressure.

Stellar  
Atmospheres  
(3)

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

# Radiation Pressure Gradient

Stellar  
Atmospheres  
(3)

Assume a **grey atmosphere** (= opacity is independent of wavelength and is described by the Rosseland mean opacity  $\bar{\kappa}$ ), and ignore all the remaining wavelength dependencies by integrating over all wavelengths

$$I = \int_0^\infty I_\lambda \, d\lambda$$

and

$$S = \int_0^\infty S_\lambda \, d\lambda$$

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

# Radiation Pressure Gradient

Stellar  
Atmospheres  
(3)

The transfer equation for a plane-parallel gray atmosphere is

$$\mu \frac{dI}{d\tau_{\perp}} = I - S$$

with  $\mu = \cos \theta$ .

Integrating over all solid angles and recalling that  $S$  is isotropic, we have

$$\frac{d}{d\tau_{\perp}} \int I \mu d\Omega = \int I d\Omega - \int S d\Omega = \Omega$$

$$\frac{dF_{\text{rad}}}{d\tau_{\perp}} = 4\pi(\langle I \rangle - S)$$

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

# Radiation Pressure Gradient

Furthermore, multiplying the transfer equation by  $\mu = \cos \theta$  and integrating over all solid angles, we find

$$\frac{d}{d\tau_{\perp}} \underbrace{\int I \mu^2 d\Omega}_{=cP_{\text{rad}}} = \underbrace{\int I \mu d\Omega}_{=F_{\text{rad}}} - S \underbrace{\int \mu d\Omega}_{=0}$$

Recall that

$$F_{\lambda} d\lambda = \int I_{\lambda} d\lambda \mu d\Omega$$

and

$$P_{\lambda, \text{rad}} d\lambda = \frac{1}{c} \int I_{\lambda} d\lambda \mu^2 d\Omega$$

Therefore, we have

$$\frac{dP_{\text{rad}}}{d\tau_{\perp}} = \frac{1}{c} F_{\text{rad}}.$$



# Radiation Pressure

Stellar  
Atmospheres  
(3)

A photon has **momentum**  $p_\lambda = E_\lambda/c$ .

We consider photons hitting a solid wall, transferring momentum. We have the **force**

$$F = \frac{dp_{\lambda\perp}}{dt} = \frac{1}{c} \frac{dE_\lambda}{dt} \cos \theta$$



This generates **radiation pressure**:

$$dP_\lambda = \frac{F}{d\sigma} = \frac{1}{c} \frac{dE_\lambda}{dt} \frac{\cos \theta}{d\sigma} = \frac{1}{c} I_\lambda \cos^2 \theta d\omega d\lambda$$
$$\Rightarrow P(\lambda) = \frac{1}{c} \oint_{4\pi} I_\lambda \cos^2 \theta d\omega = \frac{4\pi}{c} K_\lambda$$

Where  $K_\lambda$  is the **K-integral**

$$K_\lambda = \frac{1}{4\pi} \oint I_\lambda \cos^2 \theta d\omega.$$

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary



# Spherical vs. Plane-Parallel Geometry

Stellar  
Atmospheres  
(3)


When calculating the transfer equations in stars, in principal, we would have to consider spherical geometry.

Fortunately, the **geometrical thickness** of most photospheres is small compared to their radii, thus permitting the plane-parallel approximation  $r \rightarrow \infty$ :

$$\frac{dl_\nu}{ds} = -\cos\theta \frac{\partial l_\nu}{\partial t}$$

which we can derive from

$$\begin{aligned} \frac{dl_\nu}{ds} &= \frac{\partial l_\nu}{\partial r} \frac{dr}{ds} + \frac{\partial l_\nu}{\partial \theta} \frac{d\theta}{ds} \\ \frac{dr}{ds} &= \cos\theta, \quad \frac{d\theta}{ds} = -\frac{\sin\theta}{r} \end{aligned}$$



Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

# Plane-Parallel Atmosphere Approximation

Stellar  
Atmospheres  
(3)

Consider the atmosphere as a plane-parallel medium, define a vertical optical depth,  $\tau_{\lambda,\perp}(z)$ , as


$$\tau_{\lambda,\perp}(z) \equiv \int_z^0 \rho \kappa_{\lambda}(z) dz$$

Since  $dz = \cos \theta ds$ , the optical depth of a ray traveling upward at an angle  $\theta$  from the same initial position  $z$  can be related with  $\tau_{\lambda,\perp}(z)$  by

$$\tau_{\lambda} = \frac{\tau_{\lambda,\perp}}{\cos \theta} = \frac{\tau_{\lambda,\perp}}{\mu}$$

Hence:

$$\mu \frac{dI_{\lambda}}{d\tau_{\lambda,\perp}} = I_{\lambda} - S_{\lambda}$$



Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

# Plane-Parallel Atmosphere Approximation

Stellar  
Atmospheres  
(3)

In most cases, stellar atmospheres are in an equilibrium state, no net energy is subtracted from or added to the radiation field. For a plane-parallel atmosphere, this means that the radiative flux must have the same value at any level, including its surface.

Hence,

$$F_{\text{rad}} = F_{\text{surf}} = \sigma_{\text{SB}} T_e^4 = \text{const}$$

Also, since the flux is a constant, we have

$$\frac{dF_{\text{rad}}}{d\tau_{\perp}} = 0 = 4\pi(\langle I \rangle - S) \Rightarrow \langle I \rangle = S$$

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

# Plane-Parallel Atmosphere Approximation

Stellar  
Atmospheres  
(3)

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

In most cases, stellar atmospheres are in an equilibrium state, no net energy is subtracted from or added to the radiation field. For a plane-parallel atmosphere, this means that the radiative flux must have the same value at any level, including its surface.

Hence,

$$F_{\text{rad}} = F_{\text{surf}} = \sigma_{\text{SB}} T_e^4 = \text{const}$$

Also, since the flux is a constant, we have

$$\frac{dF_{\text{rad}}}{d\tau_{\perp}} = 0 = 4\pi(\langle I \rangle - S) \Rightarrow \langle I \rangle = S$$

The radiation pressure can be found by integrating

$$\frac{dF_{\text{rad}}}{dr} = -\frac{\rho \bar{\kappa}}{c} F_{\text{rad}}$$

$$P_{\text{rad}} = \frac{1}{c} F_{\text{rad}} \tau_{\perp} + C$$

where  $C$  is the constant of integration. A **boundary condition** is needed to solve  $C$ .

# The Plane-Parallel Transfer Equation

The plane-parallel transfer equation for stars with thin photospheres


$$\cos \theta \frac{dI_{\lambda}(\theta)}{d\tau_{\lambda}} = I_{\lambda}(\theta) - S_{\lambda}$$

is identical to the parallel-ray transfer equation (for ISM studies)

$$\frac{dI_{\theta}}{d\tau_{\theta}} = -I_{\lambda} + S_{\lambda}$$

except for:

- the  $\cos \theta$  term, because the optical depth is measured along the radial direction  $x$  and not along the line of sight, e.g.  
 $d\tau_{\lambda} = -\kappa_{\lambda} \rho dx$
- sign change, since we are now looking from the outside in, along the direction  $x$ .



# The Plane-Parallel Transfer Equation

The plane-parallel transfer equation for stars with thin photospheres


$$\cos \theta \frac{dI_{\lambda}(\theta)}{d\tau_{\lambda}} = I_{\lambda}(\theta) - S_{\lambda}$$

is identical to the parallel-ray transfer equation (for ISM studies)

$$\frac{dI_{\theta}}{d\tau_{\theta}} = -I_{\lambda} + S_{\lambda}$$

except for:

- the  $\cos \theta$  term, because the optical depth is measured along the radial direction  $x$  and not along the line of sight, e.g.  
 $d\tau_{\lambda} = -\kappa_{\lambda} \rho dx$
- sign change, since we are now looking from the outside in, along the direction  $x$ .



The full spherical geometry transfer equation is necessary for **supergiants**.  
There is no flux going into the star, i.e.

# Surface Intensity

To derive the intensity at the surface, we multiply the plane-parallel transfer equation by an integrating factor  $e^{-\tau/\cos\theta} = e^{-u}$ ,

$$\frac{dI_{\lambda}(\theta)}{du} e^{-u} - I_{\lambda}(\theta) e^{-u} = -S_{\lambda} e^{-u}$$

Stellar  
Atmospheres  
(3)

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

# Surface Intensity

To derive the intensity at the surface, we multiply the plane-parallel transfer equation by an integrating factor  $e^{-\tau/\cos\theta} = e^{-u}$ ,

$$\frac{dI_{\lambda}(\theta)}{du} e^{-u} - I_{\lambda}(\theta) e^{-u} = -S_{\lambda} e^{-u}$$

This can be written as

$$\frac{d(I_{\lambda}(\theta) e^{-u})}{du} = -S_{\lambda} e^{-u}$$

Stellar  
Atmospheres  
(3)

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary



# Surface Intensity

To derive the intensity at the surface, we multiply the plane-parallel transfer equation by an integrating factor  $e^{-\tau/\cos\theta} = e^{-u}$ ,

$$\frac{dI_{\lambda}(\theta)}{du} e^{-u} - I_{\lambda}(\theta) e^{-u} = -S_{\lambda} e^{-u}$$

This can be written as

$$\frac{d(I_{\lambda}(\theta) e^{-u})}{du} = -S_{\lambda} e^{-u}$$

Integrating  $du$  from 0 to  $\infty$ :

$$[I_{\lambda}(\theta) e^{-u}]_0^{\infty} = - \int_0^{\infty} S_{\lambda}(\tau_{\lambda}) e^{-u} du$$

$$I_{\lambda}(0, \theta) = \int_0^{\infty} S_{\lambda}(\tau_{\lambda}) e^{-u} du$$

# Grey Atmosphere

Stellar  
Atmospheres  
(3)

In the previous section, we assumed that the opacity can be independent of  $\lambda$ , i.e.  $\kappa_\lambda = \kappa$ . We call such a hypothetical atmosphere a **grey atmosphere**.

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

In the theory of stellar atmospheres, much of the technical effort goes into iteration schemes using equations of radiative equilibrium (such as discussed today) to find the source function  $S_\lambda$ .

Often, a starting point for such iterations is the grey atmosphere.

# Thermal (Radiative) Equilibrium

Stellar  
Atmospheres  
(3)

In stellar atmospheres, radiation dominates transfer of energy, so we can discuss three conditions of radiative equilibrium which can be used to derive the **temperature structure in the photosphere**.

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

The radiation we see from the Sun comes from a layer of geometrical height of a few hundred km.

In a column of 100 km height and  $1 \text{ cm}^2$  cross-section there are  $10^{24}$  particles (since  $\rho \sim 10^{17} / \text{cm}^3$  in the Sun), each of which has a thermal energy of this column is therefore  $10^{12} \text{ erg/cm}^2$ . The observed radiative energy loss (per  $\text{cm}^2$ ) of the Solar surface is  $F_{\odot} = 6.3 \times 10^{10} \text{ erg/cm}^2/\text{s}$ .

# Thermal (Radiative) Equilibrium

Stellar  
Atmospheres  
(3)

In stellar atmospheres, radiation dominates transfer of energy, so we can discuss three conditions of radiative equilibrium which can be used to derive the **temperature structure in the photosphere**.

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

The radiation we see from the Sun comes from a layer of geometrical height of a few hundred km.

In a column of 100 km height and  $1 \text{ cm}^2$  cross-section there are  $10^{24}$  particles (since  $\rho \sim 10^{17} / \text{cm}^3$  in the Sun), each of which has a thermal energy of this column is therefore  $10^{12} \text{ erg/cm}^2$ . The observed radiative energy loss (per  $\text{cm}^2$ ) of the Solar surface is  $F_{\odot} = 6.3 \times 10^{10} \text{ erg/cm}^2/\text{s}$ .

If the Sun shines at a constant rate, the **energy content** of the Solar photosphere can last only for 15 seconds without being resupplied from below. Exact that amount of energy has to be supplied or else the photosphere would quickly change temperature.

# First Equation of Radiative Equilibrium

Stellar  
Atmospheres  
(3)

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

As the photosphere is not changing temperature,  $dF/dt = 0$  or  $dF/dx = 0$  or  $dF/d\tau = 0$ , i.e. the total flux must be constant at all depth of the photosphere because of conservation of energy. This is the **first equation of radiative equilibrium**:

$$F(x) = F(0) = \text{const} = \sigma_{\text{SB}} T_{\text{eff}}^4$$

When all the energy is carried by radiation, we have

$$F(x) = \int_0^\infty F_\lambda(\tau_\lambda) d\lambda = F(0)$$

Although the shape of  $F_\lambda$  can be expected to change very significantly with depth, its integral remains invariant.

If **other sources of energy transport** must be taken into account, then a more general expression of flux constancy must be applied:

$$\Phi(x) + \int_0^\infty F_\lambda(\tau_\lambda) d\lambda = F(0)$$

Here,  $\Phi(x)$  is for example the convective flux.

# Second Equation of Radiative Equilibrium

Stellar  
Atmospheres  
(3)

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

We integrate the plane-parallel transfer equation over the solid angle  $\omega$ :

$$\int \cos \theta \frac{dI_{\lambda}(\tau_{\lambda}, \theta)}{d\tau_{\lambda}} d\Omega = \int I_{\lambda}(\tau_{\lambda}, \theta) d\Omega - \int S_{\lambda}(\tau_{\lambda}) d\Omega$$

$$\frac{d}{d\tau_{\lambda}}(F_{\lambda}(\tau_{\lambda})) = 4\pi(J_{\lambda}(\tau_{\lambda})) - \int S_{\lambda}(\tau_{\lambda}) d\Omega$$

Based on the definition of the mean intensity:

$$J_{\lambda} = \frac{1}{4\pi} \oint I_{\lambda} d\Omega$$

Finally, assuming  $S_{\lambda}$  to be isotropic, we obtain

$$\frac{1}{4\pi} \frac{d}{d\tau_{\lambda}}(F_{\lambda}(\tau_{\lambda})) = J_{\lambda}(\tau_{\lambda}) - S_{\lambda}(\tau_{\lambda})$$

In the case of a grey atmosphere (opacity  $\kappa$  independent of wavelength)

$$\frac{1}{4\pi} \frac{d}{d\tau} F(\tau) = -S(\tau) + J(\tau) = 0$$

Since  $dF/d\tau = 0$ , the source function must equal the mean intensity  $J$ .

# Second Equation of Radiative Equilibrium

Stellar  
Atmospheres  
(3)

If the atmosphere is not grey, which is the situation for most stars, let's incorporate the opacity  $\kappa$  into the RHS, and integrating over wavelength

$$\frac{1}{4\pi} \frac{d}{dt} \left( \int_0^\infty F(\tau_\lambda) d\lambda \right) = \int_0^\infty (\kappa_\lambda S_\lambda - \kappa_\lambda J_\lambda) d\lambda = 0$$

Since  $dF/dt = 0$ , we can get the radiative balance equation (energy conservation)

$$\int_0^\infty \kappa_\lambda J_\lambda d\lambda = \int_0^\infty \kappa_\lambda S_\lambda d\lambda$$

This is the **second equation of radiative equilibrium** and can be understood as the total energy absorbed (LHS) must equal the total energy re-emitted (RHS) if no heating or cooling is taking place.

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

# Third Equation of Radiative Equilibrium

Stellar  
Atmospheres  
(3)

The third equation is obtained by multiplying the transfer equation by  $\cos \theta$  and integrating over the solid angle and the over the wavelength:

$$\oint \cos^2 \theta \frac{dI_\lambda(\tau_\lambda, \theta)}{d\tau_\lambda} d\Omega = \oint \cos \theta dI_\lambda(\tau_\lambda, \theta) \Omega - \int \cos \theta S_\lambda(\tau_\lambda, \theta) d\Omega$$

$$K_\lambda(\tau_\lambda) = \frac{1}{4\pi} \oint I_\lambda \cos^2 \theta d\Omega \quad F_\lambda = \oint I_\lambda \cos \theta d\Omega \quad 0 \text{ (} S_\lambda \text{ is isotropic)}$$

$$4\pi \int \frac{dK_\lambda}{d\tau_\lambda} d\lambda = \int F_\lambda d\lambda = F(\tau)$$

We get the **third radiative equilibrium condition**:

$$\int_0^\infty \frac{dK_\lambda}{d\tau_\lambda} d\lambda = \frac{F(\tau)}{4\pi}$$

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary



# Equations of Radiative Equilibrium

All three radiative equilibrium conditions are not independent.

A  $S_\lambda$  that is a solution of one will be the solution of all three.

The **flux constant**  $F(0)$  is often expressed in terms of an effective temperature  $F(0) = \sigma_{\text{SB}} T_{\text{eff}}^4$ .

When model photospheres are constructed using flux constancy as a condition to be fulfilled by the model, the effective temperature becomes one of the fundamental parameters characterizing the model.

Stellar  
Atmospheres  
(3)

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

# Equations of Radiative Equilibrium

All three radiative equilibrium conditions are not independent.

A  $S_\lambda$  that is a solution of one will be the solution of all three.

The **flux constant**  $F(0)$  is often expressed in terms of an effective temperature  $F(0) = \sigma_{\text{SB}} T_{\text{eff}}^4$ .

When model photospheres are constructed using flux constancy as a condition to be fulfilled by the model, the effective temperature becomes one of the fundamental parameters characterizing the model.

In **real stars**, energy is created or lost from the radiation field through e.g. convection, magnetic fields, plus in supernovae atmospheres energy conservation is not valid (radioactive decay of Ni to Fe), so the energy constraints are more complicated in reality.

Stellar  
Atmospheres  
(3)

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

# The Depth Dependence of the Source Function

In a **grey atmosphere**, with  $K(\tau) = \int_0^\infty K_\lambda d\lambda$ , the third equation implies:

$$\frac{dK(\tau)}{d\tau} = \frac{F(\tau)}{4\pi}$$

We can differentiate this, and insert our earlier result:

$$\frac{d^2 K(\tau)}{d\tau^2} = \frac{1}{4\pi} \frac{dF(\tau)}{d\tau} = J(\tau) - S(\tau) = 0$$

Integration of the equation w.r.t.  $\tau$  gives  $K(\tau) = c_1\tau + c_2$  where  $dK/d\tau = c_1 = F/(4\pi)$ .

# The Depth Dependence of the Source Function

Stellar  
Atmospheres  
(3)

In a **grey atmosphere**, with  $K(\tau) = \int_0^\infty K_\lambda d\lambda$ , the third equation implies:

$$\frac{dK(\tau)}{d\tau} = \frac{F(\tau)}{4\pi}$$

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

We can differentiate this, and insert our earlier result:

$$\frac{d^2 K(\tau)}{d\tau^2} = \frac{1}{4\pi} \frac{dF(\tau)}{d\tau} = J(\tau) - S(\tau) = 0$$

Integration of the equation w.r.t.  $\tau$  gives  $K(\tau) = c_1\tau + c_2$  where  $dK/d\tau = c_1 = F/(4\pi)$ .

For a given  $F$ , we now have two equations to determine the three unknowns  $J$ ,  $S$ , and  $K$  (or  $c_2$ ). We need an additional relation between two of these variables in order to determine all three.

# Limb Darkening

Stellar  
Atmospheres  
(3)

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary





FIG. 1.—A typical drift scan plotted against scan angle  $\phi/\phi_{\odot}$ . The intensities plotted here have been corrected for zero point and normalized. Representative values of  $\mu = \cos \theta$  are marked along the following half of the scan.

left: Sun, observed at Venus transit

right: angular dependence of Solar intensity, credit: Petro et al. (1984)

# Limb Darkening

Stellar  
Atmospheres  
(3)

Limb darkening is the gradual decrease in brightness of the disk of a star as observed from its centre to its edge, or limb. It can be easily seen in photographs of the Sun.

Its understanding offered astronomers an opportunity to construct models with such gradients. This encouraged the development of the **theory of radiative transfer**.

Recap


Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary



# Limb Darkening

Stellar  
Atmospheres  
(3)

Let us assume a linear source function,  $S_\lambda(\tau_\lambda) = a_\lambda + b_\lambda \tau_\lambda$ .  
We derive

$$\begin{aligned} I_\lambda(0, \theta) &= \int_0^\infty a_\lambda e^{-u} du + \int_0^\infty b_\lambda \tau_\lambda e^{-u} du \\ &= a_\lambda \int_0^\infty e^{-u} du + b_\lambda \cos \theta \int_0^\infty u e^{-u} du \end{aligned}$$

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

# Limb Darkening

Stellar  
Atmospheres  
(3)

Let us assume a linear source function,  $S_\lambda(\tau_\lambda) = a_\lambda + b_\lambda \tau_\lambda$ .  
We derive

$$\begin{aligned} I_\lambda(0, \theta) &= \int_0^\infty a_\lambda e^{-u} du + \int_0^\infty b_\lambda \tau_\lambda e^{-u} du \\ &= a_\lambda \int_0^\infty e^{-u} du + b_\lambda \cos \theta \int_0^\infty u e^{-u} du \end{aligned}$$

Recall  $u = \tau / \cos(\theta)$ , so  $\tau = u \cos(\theta)$  and we can use the standard integral

$$\int_0^\infty u^n e^{-u} du = n!$$

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary



# Limb Darkening

Stellar  
Atmospheres  
(3)

Let us assume a linear source function,  $S_\lambda(\tau_\lambda) = a_\lambda + b_\lambda \tau_\lambda$ .  
We derive

$$\begin{aligned} I_\lambda(0, \theta) &= \int_0^\infty a_\lambda e^{-u} du + \int_0^\infty b_\lambda \tau_\lambda e^{-u} du \\ &= a_\lambda \int_0^\infty e^{-u} du + b_\lambda \cos \theta \int_0^\infty u e^{-u} du \end{aligned}$$

Recall  $u = \tau / \cos(\theta)$ , so  $\tau = u \cos(\theta)$  and we can use the standard integral

$$\int_0^\infty u^n e^{-u} du = n!$$

We obtain

$$I_\lambda(0, \theta) = a_\lambda + b_\lambda \cos \theta = S_\lambda(\tau_\lambda = \cos \theta)$$

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

# Limb Darkening

Let us assume a linear source function,  $S_\lambda(\tau_\lambda) = a_\lambda + b_\lambda \tau_\lambda$ .  
We derive

$$\begin{aligned} I_\lambda(0, \theta) &= \int_0^\infty a_\lambda e^{-u} du + \int_0^\infty b_\lambda \tau_\lambda e^{-u} du \\ &= a_\lambda \int_0^\infty e^{-u} du + b_\lambda \cos \theta \int_0^\infty u e^{-u} du \end{aligned}$$

Recall  $u = \tau / \cos(\theta)$ , so  $\tau = u \cos(\theta)$  and we can use the standard integral

$$\int_0^\infty u^n e^{-u} du = n!$$

We obtain

$$I_\lambda(0, \theta) = a_\lambda + b_\lambda \cos \theta = S_\lambda(\tau_\lambda = \cos \theta)$$

In the linear approximation for the source function, the optical depth lies between 0 and 1. From the center of the star we see radiation leaving the star perpendicular to the surface:  $I_\lambda(0, 0^\circ) = a_\lambda + b_\lambda$ , whilst at the limb the light leaves the surface at an angle  $I_\lambda(0, 90^\circ) = a_\lambda$ .

# Limb Darkening

Let us assume a linear source function,  $S_\lambda(\tau_\lambda) = a_\lambda + b_\lambda \tau_\lambda$ .  
We derive

$$\begin{aligned} I_\lambda(0, \theta) &= \int_0^\infty a_\lambda e^{-u} du + \int_0^\infty b_\lambda \tau_\lambda e^{-u} du \\ &= a_\lambda \int_0^\infty e^{-u} du + b_\lambda \cos \theta \int_0^\infty u e^{-u} du \end{aligned}$$

Recall  $u = \tau / \cos(\theta)$ , so  $\tau = u \cos(\theta)$  and we can use the standard integral

$$\int_0^\infty u^n e^{-u} du = n!$$

We obtain

$$I_\lambda(0, \theta) = a_\lambda + b_\lambda \cos \theta = S_\lambda(\tau_\lambda = \cos \theta)$$

In the linear approximation for the source function, the optical depth lies between 0 and 1. From the center of the star we see radiation leaving the star perpendicular to the surface:  $I_\lambda(0, 0^\circ) = a_\lambda + b_\lambda$ , whilst at the limb the light leaves the surface at an angle  $I_\lambda(0, 90^\circ) = a_\lambda$ .

**Limb darkening** occurs: less light from the limb vs. from the center if  $b_\lambda > 0$ .

# Limb Darkening

Stellar  
Atmospheres  
(3)

We convert the solution of the transfer equation by using the vertical optical depth,  $\tau_{\perp}$ :

$$I_{\lambda}(0) = I_{\lambda,0} e^{-\tau_{\lambda,\perp}/\mu} - \int_{\tau_{\lambda,\perp}/\mu}^0 \frac{S_{\lambda}}{\mu} e^{-\tau_{\lambda,\perp}/\mu} d\tau_{\lambda,\perp}$$

Taking the initial position of the rays to be at the center, where  $\tau_{\lambda,\perp} = \infty$ , we have

$$I_{\lambda}(0) = \int_0^{\infty} \frac{S_{\lambda}}{\mu} e^{-\tau_{\lambda,\perp}/\mu} d\tau_{\lambda,\perp}$$

The source function is often assumed to have the form

$$S_{\lambda} = a_{\lambda} + b_{\lambda} \tau_{\lambda,\perp}$$

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

# Limb Darkening

Applying our model of plane-parallel gray atmosphere in LTE with the Eddington approximation, we can obtain

$$S = \langle I \rangle = \frac{3\sigma_{\text{SB}}}{4\pi} T_e^4 \left( \tau_{\perp} + \frac{2}{3} \right) = a + b\tau_{\perp}$$

Therefore,

$$a = \frac{\sigma_{\text{SB}}}{2\pi} T_e^4, \quad b = \frac{3\sigma_{\text{SB}}}{4\pi} T_e^4$$

Stellar  
Atmospheres  
(3)

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

# Limb Darkening

Stellar  
Atmospheres  
(3)

Applying our model of plane-parallel gray atmosphere in LTE with the Eddington approximation, we can obtain


$$S = \langle I \rangle = \frac{3\sigma_{\text{SB}}}{4\pi} T_e^4 \left( \tau_{\perp} + \frac{2}{3} \right) = a + b\tau_{\perp}$$

Therefore,

$$a = \frac{\sigma_{\text{SB}}}{2\pi} T_e^4, \quad b = \frac{3\sigma_{\text{SB}}}{4\pi} T_e^4$$

The emergent intensity will have the form  $I(0) = a + b\mu$ , and the ratio of the emergent intensity is

$$\begin{aligned} \frac{I(\theta)}{I(\theta=0)} &= \frac{a + b \cos \theta}{a + b} \\ &= \frac{2}{5} + \frac{3}{5} \cos \theta. \end{aligned}$$



Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

# The Quadratic Source Function

Until now: We assumed a **linear source function**.

More generally: If

$$S_{\lambda}(\tau_{\lambda}) = \sum_n a_{\lambda n} \tau_{\lambda}^n$$

then

$$I_{\lambda}(0, \theta) = \sum_n A_n \cos^n \theta, \quad A_n = a_{\lambda n} \int_0^{\infty} u^n e^{-u} du = a_{\lambda n} n$$

# The Quadratic Source Function

Until now: We assumed a **linear source function**.

More generally: If

$$S_{\lambda}(\tau_{\lambda}) = \sum_n a_{\lambda n} \tau_{\lambda}^n$$

then

$$I_{\lambda}(0, \theta) = \sum_n A_n \cos^n \theta, \quad A_n = a_{\lambda n} \int_0^{\infty} u^n e^{-u} du = a_{\lambda n} n$$

We still get  $S_{\lambda}(0)$  at the limb, but a more complicated result at the center.  
For example, a quadratic term requires the solution of

$$S(\tau_{\lambda}) = a_{0\lambda} + a_{1\lambda} \tau_{\lambda} + a_{2\lambda} \tau_{\lambda}^2, \quad I_{\lambda}(0, \theta) = a_{0\lambda} + a_{1\lambda} \cos \theta + 2a_{2\lambda} \cos^2 \theta$$



# The Quadratic Source Function

Until now: We assumed a **linear source function**.

More generally: If

$$S_{\lambda}(\tau_{\lambda}) = \sum_n a_{\lambda n} \tau_{\lambda}^n$$

then

$$I_{\lambda}(0, \theta) = \sum_n A_n \cos^n \theta, \quad A_n = a_{\lambda n} \int_0^{\infty} u^n e^{-u} du = a_{\lambda n} n$$

We still get  $S_{\lambda}(0)$  at the limb, but a more complicated result at the center. For example, a quadratic term requires the solution of


$$S(\tau_{\lambda}) = a_{0\lambda} + a_{1\lambda} \tau_{\lambda} + a_{2\lambda} \tau_{\lambda}^2, \quad I_{\lambda}(0, \theta) = a_{0\lambda} + a_{1\lambda} \cos \theta + 2a_{2\lambda} \cos^2 \theta$$

At  $\theta = 90^\circ$ ,  $\tau_{\lambda} = 0$ , whilst at  $\theta = 0^\circ$ ,  $\tau_{\lambda} \sim 1 + 2a_{1\lambda}/a_{2\lambda}$  providing  $a_{2\lambda} \ll a_{1\lambda}$ . This gives a **ratio of the limb-to-center intensity** of

$$I_{\lambda}(0, 90^\circ)/I_{\lambda}(0, 0^\circ) = a_{0\lambda}/(a_{0\lambda} + a_{1\lambda} + 2a_{2\lambda})$$

# Wavelength Dependency of Limb Darkening

Limb darkening is observed to be greatest at shorter wavelengths. The temperature distribution of the upper atmosphere of the Sun can be obtained from limb darkening measurements, carried out via e.g. multi-filter images of the Solar continuum.



Limb darkening in various wavelengths. The variable  $x$  decreases with the distance from the center. Credit: Koepke et al. (2001)

# Limb Darkening for other Stars

Stellar  
Atmospheres  
(3)

Until recently, the Sun was the only star for which limb darkening was observed, as one needs to spatially resolve the disk to measure it. Other methods are now possible.

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary


Limb darkening for other stars than our Sun can be measured by using the following methods:

1. Direct interferometry, using a high spatial resolution imaging - e.g. COAST array, providing the star is very large and nearby (a cool supergiant).
2. The light curve of an eclipsing binary system during secondary eclipse allows us to study limb darkening of the primary, although this is not trivial. A similar approach followed by extrasolar planets occulting their star (e.g. HD209458).
3. Analyzing the light curve from gravitational microlensing of a background star by a foreground source (e.g. PLANET team).

# Limb Darkening and Transit Light Curves

Limb darkening must be taken into account when working with transit curves from such as binary systems or exoplanets.

Limb darkening is a second-order, but key effect for determining the exact radius of the planet, because limb darkening not only modifies the shape of the transit light curve, but it also significantly affects the true transit depth.




credit: [juliaastro.org](http://juliaastro.org)

# Limb Darkening from Microlensing

Stellar  
Atmospheres  
(3)

Galactic gravitational microlensing occurs when a foreground object (lens) passes in front of a background star (source). The gravitational deflection of light by the lens causes the flux from the source to be amplified.

One such event, MACHO 97-BLG-28, was studied to reveal limb darkening information for the background K giant (Albrow et al. 1999).



Thick lines show how much fainter the K giant becomes at its limb in the red I (left) and blue-green V filter (right). If the star would emit a uniform amount of light across its whole stellar disk, the profile would look like the "Uniformly Bright" profile. credit: Albrow et al. (1999)

# Eddington Approximation

Stellar  
Atmospheres  
(3)

A plane-parallel gray atmosphere needs a description of the angular distribution of the intensity.

The Eddington approximation assumes an isotropic radiation field,

$$\langle I \rangle = \frac{1}{2}(I_{\text{out}} + I_{\text{in}})$$

$$F_{\text{rad}} = \pi(I_{\text{out}} - I_{\text{in}})$$

$$F_{\text{rad}} = \frac{2\pi}{3c}(I_{\text{out}} - I_{\text{in}}) = \frac{4\pi}{3c}\langle I \rangle$$

Applying  $P_{\text{rad}}$  to the plane-parallel gray atmosphere, we find

$$\frac{4\pi}{3c}\langle I \rangle = \frac{1}{c}F_{\text{rad}}\tau_{\perp} + C$$

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

# Eddington Approximation

The constant  $C$  can be determined by evaluating the first two equations of the Eddington approximation at the stellar surface, where  $\tau_{\perp} = 0$  and  $I_{\text{in}} = 0$ . Given  $\langle I(\tau_{\perp} = 0) \rangle = \frac{F_{\text{rad}}}{2\pi}$ , we get

$$C = \frac{2}{3c} F_{\text{rad}}.$$

Therefore,

$$\frac{4\pi}{3} \langle I \rangle = F_{\text{rad}} \left( \tau_{\perp} + \frac{2}{3} \right).$$

Substituting  $F_{\text{rad}} = \sigma_{\text{SB}} T_e^4$ , one obtains

$$\langle I \rangle = \frac{3\sigma_{\text{SB}}}{4\pi} T_e^4 \left( \tau_{\perp} + \frac{2}{3} \right)$$

# Eddington Approximation

Given

$$\langle I \rangle = \frac{3\sigma_{SB}}{4\pi} T_e^4 \left( \tau_{\perp} + \frac{2}{3} \right).$$

If the atmosphere is assumed to be in LTE, the source function is equal to the Planck function,  $S_{\lambda} = B_{\lambda}$ . Integrating over all wavelengths, we have

$$\langle I \rangle = S = B = \frac{\sigma_{SB} T^4}{\pi}$$

Finally, we obtain a description for the variation of the temperature in a plane-parallel gray atmosphere in LTE with the Eddington approximation

$$T^4 = \frac{3}{4} T_e^4 \left( \tau_{\perp} + \frac{2}{3} \right)$$



# Eddington Approximation

Given

$$\langle I \rangle = \frac{3\sigma_{SB}}{4\pi} T_e^4 \left( \tau_{\perp} + \frac{2}{3} \right).$$

If the atmosphere is assumed to be in LTE, the source function is equal to the Planck function,  $S_{\lambda} = B_{\lambda}$ . Integrating over all wavelengths, we have

$$\langle I \rangle = S = B = \frac{\sigma_{SB} T^4}{\pi}$$

Finally, we obtain a description for the variation of the temperature in a plane-parallel gray atmosphere in LTE with the Eddington approximation

$$T^4 = \frac{3}{4} T_e^4 \left( \tau_{\perp} + \frac{2}{3} \right)$$

Note that  $T = T_e$  at  $\tau_{\perp} = \frac{2}{3}$ .

By definition, the surface of the star is where  $T = T_e$ , i.e.  $\tau_{\perp} = 2/3$  since  $L = 4\pi R_*^2 \sigma_{SB} T_e^4$ .

Stellar  
Atmospheres  
(3)

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

# Eddington Approximation

Stellar  
Atmospheres  
(3)

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

Previously we had seen that for the determination of the flux the anisotropy in the radiation field is very important: in the flux integral the inward-going intensities are subtracted from the outward-going ones, due to the factor  $\cos \theta$ .

But for  $K$ , a small anisotropy is unimportant: the intensities are multiplied by  $\cos^2 \theta$ , which does not change sign for inward and outward radiation.

In order to evaluate  $K$  or  $c_2$  we can approximate the radiation field by an isotropic radiation field of the mean intensity  $J$ :  $I = J$  (by definition).

From the definition of  $K_\lambda$  we obtain

$$4\pi K_\lambda = \oint I_\lambda(\tau_\lambda, \theta) \cos^2 \theta \, d\omega = J_\lambda(\tau_\lambda) \oint \cos^2 \theta \, d\omega = \frac{4\pi}{3} J_\lambda(\tau_\lambda)$$

or after division by  $4\pi$ :

$$K_\lambda(\tau_\lambda) = \frac{1}{3} J_\lambda(\tau_\lambda)$$

This approximation for the  $K$  function is known as the **Eddington approximation**.

# Eddington Approximation

Stellar  
Atmospheres  
(3)

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

Inserting the Eddington approximation into the above equation

$$\frac{dK(\tau)}{d\tau} = \frac{F(\tau)}{4\pi}$$

we find

$$\frac{dK(\tau)}{d\tau} = \frac{1}{3} \frac{dJ(\tau)}{d\tau} = \frac{F(\tau)}{4\pi} = c_1, \quad \frac{dJ(\tau)}{d\tau} = \frac{3}{4\pi} F(\tau)$$

Since the mean intensity  $J$  equals the source function  $S$  in a grey atmosphere, when integrating the latter result we obtain

$$S(\tau) = \frac{3}{4\pi} \tau F(0) + C = J(\tau)$$

From the conditions of radiative equilibrium, we finally obtained the law for the depth dependence of the source function (for a grey atmosphere assuming the Eddington approximation).

We evaluate  $C$  using boundary conditions for the known emerging flux (there is no flux going into the star), i.e.  $I(0, \theta) = I^- = 0$  for  $\pi/2 < \theta < \pi$ , plus assuming the outward intensity is independent of  $\theta$ , i.e.  $I(0, \theta) = I^+$  for  $0 < \theta < \pi/2$ .

# Eddington Approximation

Stellar  
Atmospheres  
(3)

**Boundary condition:** There is no flux going into the star, i.e.

$$I(0, \theta) = I^- = 0 \text{ for } \pi/2 < \theta < \pi.$$

We also assume that the outward intensity does not depend upon  $\theta$ , i.e.

$$I(0, \theta) = I^+ = \text{const for } 0 < \theta < \pi/2.$$

From this, we get

$$J(0) = \frac{1}{2\pi} I^+ = \frac{1}{2\pi} F(0)$$

Hence  $C = J(0) = F(0)/2\pi$ , so

$$S(\tau) = \frac{1}{\pi} \left( \frac{3}{4} \tau + \frac{1}{2} \right) F(0)$$

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

# Eddington Approximation

Stellar  
Atmospheres  
(3)

**Boundary condition:** There is no flux going into the star, i.e.

$$I(0, \theta) = I^- = 0 \text{ for } \pi/2 < \theta < \pi.$$

We also assume that the outward intensity does not depend upon  $\theta$ , i.e.

$$I(0, \theta) = I^+ = \text{const for } 0 < \theta < \pi/2.$$

From this, we get

$$J(0) = \frac{1}{2\pi} I^+ = \frac{1}{2\pi} F(0)$$

Hence  $C = J(0) = F(0)/2\pi$ , so

$$S(\tau) = \frac{1}{\pi} \left( \frac{3}{4} \tau + \frac{1}{2} \right) F(0)$$

To find the depth dependence of  $T$ , we also need to assume LTE.

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

# Temperature Structure of the Grey Atmosphere

Stellar  
Atmospheres  
(3)

In LTE, the source function is the Planck function,  
 $S(\tau) = B(\tau) = \sigma_{\text{SB}} T^4 / \pi$ .

$$B(\tau) = \frac{\sigma_{\text{SB}}}{\pi} T^4(\tau) = \frac{3}{4\pi} \left( \tau + \frac{2}{3} \right) F(0)$$

Recall that  $F(0) = \sigma_{\text{SB}} T_{\text{eff}}^4$ , by definition, so

$$\frac{1}{\pi} \sigma_{\text{SB}} T^4(\tau) = \frac{3}{4\pi} \left( \tau + \frac{2}{3} \right) \sigma_{\text{SB}} T_{\text{eff}}^4 \quad \text{or} \quad T^4(\tau) = \frac{3}{4} \left( \tau + \frac{2}{3} \right) T_{\text{eff}}^4$$

With this, we derived the **temperature dependence on optical depth**.  
Note  $T(\tau = 2/3) = T_{\text{eff}}$  as we obtained earlier, and  $T^4(\tau = 0) = T_{\text{eff}}^4/2$

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary


# Temperature Structure of the Grey Atmosphere

Stellar  
Atmospheres  
(3)

A **complete solution** of the grey case, using accurate boundary conditions, without Eddington approximation, leads to a solution only slightly different from this, usually expressed as

$$T^4(\tau) = \frac{3}{4} (\tau + q(\tau)) T_{\text{eff}}^4$$

Here,  $q(\tau)$  is a slowly varying function, called the Hopf function, with  $q = 1/\sqrt{3} = 0.577$  at  $\tau = 0$  to  $0.710$  at  $\tau = \infty$ .



Comparison between  $T(\tau)$  in the Solar atmosphere using the simplified Eddington approximation (solid) versus the exact grey atmosphere (dashed) using the Hopf function,  $q(\tau) \sim 0.0710 - 0.133 \times e^{-2\tau}$ .

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation


Summary

# Grey Temperature Structure

How good is the approximation of a grey atmosphere?

We must next look at the frequency dependence of the sources of opacity. The grey temperature distribution is shown here versus the observed Solar temperature distribution as a function of optical depth  $\tau$  at  $5000\text{\AA}$  (D. Gray, Table 9.2).

The poor match is because the opacity is wavelength dependent.





# Summary

Three equations of radiative equilibrium can be derived:

- a constant flux with depth
- energy absorbed equals energy emitted
- the  $K$ -integral is linear in  $\tau$ .

Stellar  
Atmospheres  
(3)

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

# Summary

Stellar  
Atmospheres  
(3)

Three equations of radiative equilibrium can be derived:

- a constant flux with depth
- energy absorbed equals energy emitted
- the  $K$ -integral is linear in  $\tau$ .

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

From these, the grey temperature distribution  $T(\tau)$  may be derived, assuming

- the Eddington approximation and
- LTE, in reasonable agreement with the exact science case.

# Summary

Stellar  
Atmospheres  
(3)

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

Three equations of radiative equilibrium can be derived:

- a constant flux with depth
- energy absorbed equals energy emitted
- the  $K$ -integral is linear in  $\tau$ .

From these, the grey temperature distribution  $T(\tau)$  may be derived, assuming

- the Eddington approximation and
- LTE, in reasonable agreement with the exact science case.

Solution to plane-parallel transfer equation at surface explains limb darkening in Sun.

# Summary

Stellar  
Atmospheres  
(3)

Recap

Radiative  
Transfer (II)

Grey  
Atmosphere

Limb  
Darkening

Eddington  
Approximation

Summary

Three equations of radiative equilibrium can be derived:

- a constant flux with depth
- energy absorbed equals energy emitted
- the  $K$ -integral is linear in  $\tau$ .

From these, the grey temperature distribution  $T(\tau)$  may be derived, assuming

- the Eddington approximation and
- LTE, in reasonable agreement with the exact science case.

Solution to plane-parallel transfer equation at surface explains limb darkening in Sun.

Limb darkening in other stars can be estimated from interferometry, eclipsing binaries, microlensing.