

ORIGINAL ARTICLE

Effect of Inhaled Glucocorticoids in Childhood on Adult Height

H. William Kelly, Pharm.D., Alice L. Sternberg, Sc.M., Rachel Lescher, M.D., Anne L. Fuhlbrigge, M.D., Paul Williams, M.D., Robert S. Zeiger, M.D., Ph.D., Hengameh H. Raissy, Pharm.D., Mark L. Van Natta, M.H.S., James Tonascia, Ph.D., and Robert C. Strunk, M.D.,
for the CAMP Research Group*

ABSTRACT

BACKGROUND

The use of inhaled glucocorticoids for persistent asthma causes a temporary reduction in growth velocity in prepubertal children. The resulting decrease in attained height 1 to 4 years after the initiation of inhaled glucocorticoids is thought not to decrease attained adult height.

METHODS

We measured adult height in 943 of 1041 participants (90.6%) in the Childhood Asthma Management Program; adult height was determined at a mean (\pm SD) age of 24.9 ± 2.7 years. Starting at the age of 5 to 13 years, the participants had been randomly assigned to receive 400 μ g of budesonide, 16 mg of nedocromil, or placebo daily for 4 to 6 years. We calculated differences in adult height for each active treatment group, as compared with placebo, using multiple linear regression with adjustment for demographic characteristics, asthma features, and height at trial entry.

RESULTS

Mean adult height was 1.2 cm lower (95% confidence interval [CI], -1.9 to -0.5) in the budesonide group than in the placebo group ($P=0.001$) and was 0.2 cm lower (95% CI, -0.9 to 0.5) in the nedocromil group than in the placebo group ($P=0.61$). A larger daily dose of inhaled glucocorticoid in the first 2 years was associated with a lower adult height (-0.1 cm for each microgram per kilogram of body weight) ($P=0.007$). The reduction in adult height in the budesonide group as compared with the placebo group was similar to that seen after 2 years of treatment (-1.3 cm; 95% CI, -1.7 to -0.9). During the first 2 years, decreased growth velocity in the budesonide group occurred primarily in prepubertal participants.

CONCLUSIONS

The initial decrease in attained height associated with the use of inhaled glucocorticoids in prepubertal children persisted as a reduction in adult height, although the decrease was not progressive or cumulative. (Funded by the National Heart, Lung, and Blood Institute and the National Center for Research Resources; CAMP ClinicalTrials.gov number, NCT00000575.)

From the University of New Mexico, Albuquerque (H.W.K., H.H.R.); Johns Hopkins University, Baltimore (A.L.S., M.L.V.N., J.T.); Washington University, St. Louis (R.L., R.C.S.); Alaska Native Medical Center, Anchorage (R.L.); Brigham and Women's Hospital, Boston (A.L.F.); University of Washington, Seattle (P.W.); and the University of California, San Diego, and Kaiser Permanente Southern California Region, San Diego (R.S.Z.). Address reprint requests to Dr. Kelly at 9828 Guadalupe Trail NW, Albuquerque, NM 87114, or at hwkelly@salud.unm.edu.

*Additional members of the Childhood Asthma Management Program (CAMP) Research Group are listed in the Supplementary Appendix, available at NEJM.org.

This article was published on September 3, 2012, at NEJM.org.

N Engl J Med 2012;367:904-12.
DOI: 10.1056/NEJMoa1203229
Copyright © 2012 Massachusetts Medical Society.

INHALED GLUCOCORTICOIDS ARE THE RECOMMENDED therapy for persistent asthma in children.¹ In prepubertal children, however, the use of inhaled glucocorticoids has been shown to reduce growth velocity, resulting in a linear growth reduction of 0.5 to 3.0 cm (approximately 1 cm on average) during the first few years of therapy.²⁻⁴ This reduction has been reported for low-to-medium doses, but the degree of reduction is dependent on the type of inhaled glucocorticoid and the delivery method.⁵⁻⁹

Although growth velocity returns to normal within a few years after the initiation of inhaled glucocorticoid therapy (resulting in a deficit in height that is not progressive), the long-term effect of the initial decrease in growth velocity on attainment of adult height is still unclear.¹⁰⁻¹² Retrospective, cross-sectional studies of the effect of inhaled glucocorticoid therapy in childhood on the attainment of adult height have had conflicting results.^{10,13-15} In one small longitudinal study,¹¹ there was no significant difference between actual adult height and predicted adult height in 142 participants who had received budesonide in variable daily doses (mean, 427 μ g) during childhood for a mean of 9.2 years.

At the end of the Childhood Asthma Management Program (CAMP) clinical trial, we predicted that the children receiving budesonide (Pulmicort Turbuhaler, AstraZeneca) for a mean follow-up of 4.3 years would attain the same adult height as the children receiving nedocromil (Tilade, Rhone-Poulenc Rorer) or placebo.² It has been hypothesized that children receiving inhaled glucocorticoids grow for a longer period of time, eventually catching up and having no long-term effects from the decreased growth velocity seen in the first few years of therapy.¹⁶ Thus, the effect of inhaled glucocorticoids on height has often been characterized as growth retardation rather than growth suppression. We later reported that the heights of the participants in the budesonide group had not caught up with the heights of participants in the placebo group after an additional 4.8 years of follow-up (total, 9.1 years), until the ages of 12 to 23 years.¹² The purpose of the current study was to continue the height follow-up of the CAMP participants to assess the effects of budesonide and nedocromil on adult height.

METHODS

PARTICIPANTS AND STUDY DESIGN

From December 1993 through September 1995, we randomly assigned 1041 children between the ages of 5 and 13 years with mild-to-moderate asthma to one of three study groups in the double-blind, placebo-controlled CAMP trial. In this study, we compared the efficacy and safety of 200 μ g of budesonide administered by means of a dry-powder inhaler twice daily (400 μ g per day), 8 mg of nedocromil administered by means of a metered-dose inhaler twice daily (16 mg per day), and placebo. Albuterol was used for asthma symptoms in all three groups. The design, methods, and results of the trial have been described previously.^{2,17} At the end of the trial (mean follow-up, 4.3 years), the children were recruited into an observational cohort, as described previously.¹² During follow-up, the children were treated for asthma by their primary care physicians under advisement from the CAMP physicians on the basis of the guidelines of the National Asthma Education and Prevention Program.¹ Height and weight were measured every 6 months during the initial 4.5 years of observational follow-up and 1 to 2 times a year during the next 8 years. Adult height was determined at a mean (\pm SD) age of 24.9 \pm 2.7 years. Tanner staging was performed annually until the participants were 18 years of age or attained sexual maturity. (For details regarding the trial design and phases of the observational follow-up, see Fig. S1 in the Supplementary Appendix, available with the full text of this article at NEJM.org.) All study phases were approved by the institutional review board at each study center. Participants who were 18 years of age or older provided written informed consent; those under the age of 18 years provided assent, with written informed consent provided by their parents or guardians.

At each clinic visit, a trained technician used a stadiometer (Harpender 603, Holtain) to measure height, as described in the protocol (available at NEJM.org), and investigators obtained an interim history of participants' health and medication use. During the first 8 years of follow-up, this information was supplemented with information obtained during telephone contacts that were evenly spaced around the clinic visits.

On the basis of reported recommendations,¹⁸⁻²¹ adult height was defined as either the mean of all measurements performed at the age of 18 years or older for women and at the age of 20 years or older for men or, if height was not measured at these ages, the most recently obtained height that was less than 1 cm greater than a height obtained at least 1 year previously.

STATISTICAL ANALYSIS


We used a multiple linear regression model to compare mean height in the budesonide and nedocromil groups with that in the placebo group, with adjustment for eight covariates at trial entry: age, race or ethnic group, sex, clinic, height, duration of asthma (<3 years, ≥3 to <7 years, or ≥7 years), severity of asthma (moderate or mild), and presence or absence of skin-test reactivity. Data for each participant were analyzed in the study group to which the participant had been randomly assigned, regardless of subsequent asthma treatment. The adjusted mean height in each study group was computed from the regression model at the mean values for the covariates.²² To assess the statistical significance of potential variations in height differences between the budesonide or nedocromil group and the placebo group across subgroups defined by race or ethnic group, age at trial entry, and duration of asthma at entry, we added covariate interaction terms to the regression model. Although these post hoc interaction tests may have a low statistical power for detecting variations in height effects across subgroups, we wanted to examine the qualitative consistency of the budesonide and nedocromil effects across subgroups.

We used chi-square tests for categorical variables and t-tests for continuous variables to assess differences in baseline characteristics between participants for whom data regarding adult height were missing and those for whom such data were available. We used logistic-regression analysis to check for systematic differences between participants with adult-height data and those without such data with respect to study-group distribution in subgroups, as defined by the baseline covariates. Effects of missing data on adult height were further addressed in sensitivity analyses with the use of two alternative methods to impute missing adult height. In the first method, we used a single imputation of adult height on the basis of bone age obtained at

the end of the trial and the equations of Tanner et al.^{2,19} In the second method, we used multivariate multiple imputation with 13 imputations from simulations of a Bayesian posterior predictive distribution of the missing data, using an iterative Markov chain Monte Carlo method based on the multivariate normal distribution provided by Stata software.²³ All eight covariates in the model for difference in adult height were used together with serial height measurements obtained 2 to 102 months after trial entry. The missing adult height data were assumed to be missing completely at random. Additional sensitivity analyses were performed with the use of seven alternative definitions of adult height.

We used a linear regression model of height in relation to age, using all available age–height pairs for each participant in each 1-year age span, to determine age-specific growth velocity for each participant during the first 2 years of the trial. We assessed study-group differences in growth-velocity trends for participants between the ages of 5 and 15 years separately for each sex by means of a repeated-measures multiple linear regression model,²⁴ including an indicator variable for treatment, splines for each year of age, and interaction terms for treatment according to age. This analysis was also adjusted for the covariates used in the models of adult height. The significance of study-group differences in velocity trends was determined on the basis of the P value from an F-test of the simultaneous null effects of the study-group indicator and all age splines according to treatment interaction terms. We performed a secondary analysis of adult height in relation to the mean weight-adjusted dose of daily inhaled glucocorticoids (in micrograms per kilogram of body weight) during the first 2 years of the trial, with adjustment for demographic characteristics, status with respect to exposure to cigarette smoke in utero, and total prednisone dose throughout follow-up until adult height was attained, as well as asthma features, physical development, and vitamin D sufficiency or insufficiency at trial entry. The dose of inhaled glucocorticoids and weight were assumed to be constant over each reporting interval.

A two-sided P value of less than 0.05 was considered to indicate statistical significance, except in interaction analyses, for which a P value of less than 0.01 was considered to indicate significance. Statistical analyses were performed with



the use of Stata software, version 12 (StataCorp) or SAS software, version 9.2 (SAS Institute).

RESULTS

ADULT HEIGHT

We obtained measurements of adult height for 943 of the original 1041 CAMP participants (90.6%). Of these measurements, 96.8% were obtained from women who were at least 18 years of age or men who were at least 20 years of age; 3.2% of measurements were the most recently obtained heights that were less than 1 cm greater than a height obtained at least 1 year previously (Fig. 1). Adult height was not known for 98 participants (9.4%): 35 female participants with a median age of 12 years (range, 8 to 17) and 63 male participants with a median age of 14 years (range, 7 to 19) at the last height measurement.

The adjusted mean adult height was 1.2 cm lower in the budesonide group than in the placebo group (171.1 cm vs. 172.3 cm, $P=0.001$);

the mean adult height in the nedocromil group (172.1 cm) was similar to that in the placebo group ($P=0.61$) (Table 1 and Fig. 2).²⁵ The deficit in adult height in the budesonide group, as compared with the placebo group, was greater for women (−1.8 cm, $P=0.001$) than for men (−0.8, $P=0.10$), but the effect of budesonide on adult height did not vary significantly according to sex ($P=0.10$ for interaction) (Table 1). Although the deficit was greater for participants who were younger at trial entry than for those who were older and greater for whites than for those of other races or ethnic groups, the effect of budesonide on adult height did not vary significantly according to the age at trial entry ($P=0.12$ for interaction), race or ethnic group ($P=0.50$ for interaction), or duration of asthma at trial entry ($P=0.35$ for interaction) (Table 1, and Table S1 in the Supplementary Appendix). The deficit in the adjusted mean height in the budesonide group, as compared with the placebo group, was 1.3 cm (95% confidence interval [CI], −1.7 to −0.9) after

Table 1. Adjusted Mean Adult Height among 943 Study Participants.*

Variable	Mean Adult Height			Difference in Height			
	Budesonide (N=281)	Nedocromil (N=285)	Placebo (N=377)	Budesonide vs. Placebo (95% CI)	P Value	Nedocromil vs. Placebo (95% CI)	P Value
		cm		cm		cm	
All participants	171.1	172.1	172.3	-1.2 (-1.9 to -0.5)	0.001	-0.2 (-0.9 to 0.5)	0.61
Sex							
Female	162.8	163.9	164.6	-1.8 (-2.9 to -0.7)	0.001	-0.7 (-1.8 to 0.5)	0.26
Male	176.8	177.6	177.6	-0.8 (-1.8 to 0.2)	0.10	-0.0 (-0.9 to 0.9)	0.98
P value for interaction					0.10		0.49
Age at entry							
5–8 yr	170.7	171.8	172.6	-1.9 (-3.2 to -0.6)	0.004	-0.8 (-2.1 to 0.5)	0.22
9–13 yr	171.4	172.4	171.9	-0.5 (-1.7 to 0.6)	0.37	0.5 (-0.8 to 1.6)	0.48
P value for interaction					0.12		0.15

* Mean values for adult height have been adjusted for age, race or ethnic group, sex, clinic, duration of asthma, asthma severity, presence or absence of skin-test reactivity, and height at trial entry. In the analysis of mean adult height according to sex, 385 participants were female and 558 were male. In the analysis of age at study entry, 489 participants were 5 to 8 years of age and 454 were 9 to 13 years of age.

2 years of treatment and 1.2 cm (95% CI, -2.0 to -0.4) at the end of the CAMP trial and persisted into adulthood without progressing further (-1.2 cm; 95% CI, -1.9 to -0.5) (Fig. 2B).

GROWTH VELOCITY

Overall, age trends with respect to growth velocity in the budesonide and placebo groups differed during the first 2 years of the trial for women ($P=0.007$) and men ($P<0.001$) (Fig. S2 in the Supplementary Appendix). For both sexes, the difference in velocity reduction that was seen in the first 2 years of assigned treatment in the budesonide group, as compared with the placebo group, was primarily among prepubertal participants (girls 5 to 10 years of age, $P=0.001$; girls 11 to 15 years of age, $P=0.54$; boys 5 to 11 years of age, $P<0.001$; and boys 12 to 15 years of age, $P=0.57$).

SENSITIVITY ANALYSES


In sensitivity analyses performed to assess the potential effect of missing adult heights, the reduction in adult height in the budesonide group as compared with the placebo group was also calculated with imputation of missing adult heights based on bone age at the end of the trial¹⁹ (-1.1 cm, $P=0.004$) and multivariate multiple imputation (-1.2 cm, $P=0.002$). The between-group difference in adjusted mean adult height remained numerically similar to that in the primary analysis and significant for all seven alternative definitions of adult height (Table 2).

At trial entry, data for participants for whom measurements of adult height were not available were similar to data for those with available measurements, except for the variable of the clinic site ($P=0.002$) (Table S2 in the Supplementary Appendix). There was no interaction between the study group and any of the baseline covariates, suggesting that the study-group comparison among the 943 participants with known adult height may reasonably be generalized to the entire CAMP population.

EFFECTS OF GLUCOCORTICOID DOSE

During the trial period of 4 to 6 years, the mean adjusted total doses of inhaled glucocorticoids were 636.1 mg in the budesonide group and 88.5 mg in the nedocromil group versus 109.4 mg in the placebo group ($P<0.001$ and $P=0.14$, respectively). During follow-up after the trial ended, the mean adjusted total doses were 381.0 mg in the budesonide group and 347.9 mg in the nedocromil group versus 355.0 mg in the placebo group ($P=0.55$ and $P=0.87$, respectively).

The as-treated secondary analysis of the daily weight-adjusted dose of inhaled glucocorticoids during the first 2 years of the CAMP trial showed that a larger daily dose was associated with a lower adult height (-0.1 cm for each microgram per kilogram, $P=0.007$). In addition, lower adult height was associated with Hispanic ethnic group ($P<0.001$) and female sex ($P<0.001$), as well as a higher Tanner stage ($P<0.001$), lower height



($P < 0.001$), greater body-mass index ($P < 0.001$), longer duration of asthma ($P < 0.001$), skin-test reactivity ($P < 0.001$), and vitamin D insufficiency (≤ 30 ng

per milliliter, $P = 0.004$) at baseline. The cumulative prednisone dose from trial entry until attainment of adult height did not affect adult height ($P = 0.76$).

Table 2. Adjusted Mean Adult Height, According to the Imputation Strategy Used for Missing Data and Alternative Height Definitions.*

Variable	Total No. of Participants	Mean Adult Height			Difference in Height	
		Budesonide	Nedocromil	Placebo	Budesonide vs. Placebo (95% CI) <i>cm</i>	P Value Nedocromil vs. Placebo (95% CI) <i>cm</i>
Primary analysis	943	171.1	172.1	172.3	-1.2 (-1.9 to -0.5)	0.001
Imputation strategy						
Adult height or projected adult height†	991	171.2	172.2	172.3	-1.1 (-1.7 to -0.3)	0.004
Adult height or multiple-imputation height‡	1041	171.1	172.1	172.3	-1.2 (-1.9 to -0.4)	0.002
Alternative height definition						
Mean height at age ≥18 yr for women and ≥20 yr for men	913	171.0	172.1	172.3	-1.3 (-2.0 to -0.5)	0.001
Minimum height at age ≥18 yr for women and ≥20 yr for men	913	170.6	171.7	171.8	-1.2 (-2.0 to -0.5)	0.001
Maximum height at age ≥18 yr for women and ≥20 yr for men	913	171.4	172.6	172.7	-1.3 (-2.0 to -0.5)	0.001
Most recent height at age ≥18 yr for women and ≥20 yr for men	913	171.1	172.2	172.3	-1.2 (-2.0 to -0.5)	0.001
Mean height at age ≥21 yr for women and ≥23 yr for men	780	170.5	171.4	171.6	-1.1 (-1.8 to -0.3)	0.01
Height at time of growth of <1 cm/yr§	928	171.1	172.1	172.3	-1.2 (-1.9 to -0.5)	0.001
Mean height at age ≥21 yr for women and ≥23 yr for men or at growth of <1 cm/yr	937	171.1	172.1	172.3	-1.2 (-1.9 to -0.4)	0.002

* Mean values for adult height have been adjusted for age, race or ethnic group, sex, clinic, duration of asthma, asthma severity, presence or absence of skin-test reactivity, and height at trial entry.

† Projected adult height was calculated from the prediction equations of Tanner et al.,¹⁹ which use height, chronologic age, bone age, and (for girls) age at first menses.

‡ Adult height was imputed for the 98 participants for whom data regarding adult height were missing. The imputation analysis was completed with the use of multivariate normal regression of the available height data for the CAMP group (adult height for 943 participants and up to 21 serial height measurements performed 2 to 102 months after randomization for 1041 participants) on age, race or ethnic group, sex, clinic, duration of asthma, asthma severity, presence or absence of skin-test reactivity, and height at trial entry, stratified according to study group.

§ In this category, adult height was defined as the chronologically later height in the most recent pair of height measurements that were performed at least 1 year apart and differed by less than 1 cm.

DISCUSSION

In an intention-to-treat analysis of the growth-suppressive effect of long-term inhaled glucocorticoid therapy for asthma initiated in children between the ages of 5 and 13 years, we found that the height deficit observed at 1 to 2 years after treatment initiation persisted into adulthood, although the deficit was neither progressive nor cumulative. Our conclusion is based on a randomized comparison of adult height, with height data available for 91% of the CAMP cohort, with the use of recognized definitions of adult height,¹⁸⁻²¹ and with consistent findings in sensitivity analyses using imputation strategies for missing data and alternative definitions of adult height (Table 2). We found little evidence that the 98 participants for whom data regarding adult height were missing differed at trial entry from the 943 participants with available data (Table S2 in the Supplementary Appendix).

In contrast, the only other prospective longitudinal cohort study that followed patients into adulthood¹¹ was an open-label study, and by the time the patients reached adulthood, only 15 of the original controls were available, so the investigators recruited 51 healthy siblings of the patients with asthma to be controls. The investigators in that study based their conclusion of a lack of long-term effect on height on the finding that both controls and participants receiving budesonide attained predicted adult height rather than on a randomized comparison of the adult heights reached by the two groups.

The growth-velocity deficit that we observed in the budesonide group, as compared with the placebo group, during the first 2 years of treatment was primarily among prepubertal children (Fig. S2 in the Supplementary Appendix). Our ability to further disentangle the effects of duration of treatment, age at treatment, and puberty status during treatment on growth velocity was limited because of confounding. Nevertheless, the effect on adult height of budesonide as compared with placebo was clearly demonstrated in the CAMP population.

Two previous 1-year studies of beclomethasone dipropionate also showed a greater reduction in growth in prepubertal children than in pubertal children.^{26,27} Like other studies in which prepubertal children received different doses of the same inhaled glucocorticoids that have shown a

dose-response effect on growth,^{28,29} our study showed a weight-based, dose-dependent effect in the CAMP participants (Table S3 in the Supplementary Appendix).

We found that a longer time since asthma diagnosis at trial entry and atopy (any positive skin test) were independent risk factors for shorter adult height (Table S3 in the Supplementary Appendix). Other investigators have reported an increased incidence of short stature in children with atopy and asthma.³⁰⁻³² One of these studies³¹ showed that short stature was associated with an early onset of asthma (before the age of 3 years), a finding that is consistent with our data. However, atopy-induced growth retardation has been associated with a delay in bone maturation and thus was thought to be unlikely to affect adult height.^{32,33} Our results suggest that when asthma and atopy impair growth, the deficit may persist into adulthood.

We selected a daily dose of 400 μ g of budesonide for the CAMP trial to ensure a therapeutic effect in both children with mild asthma and those with moderate asthma.² Since then it has been shown that daily administration of 200 μ g of budesonide by means of a dry-powder inhaler effectively controls asthma symptoms and reduces exacerbations in children 5 to 11 years of age.³ Even at this lower dose, there was a reported mean reduction of 1.0 cm in height during the first 2 years of therapy.³ Although the systemic effects of inhaled glucocorticoids are dose-dependent, they are also dependent on the therapeutic index of the specific inhaled glucocorticoid and the delivery device used.⁴⁻⁹ Thus, it seems prudent to select inhaled glucocorticoids and devices with higher therapeutic indexes and to use them in the lowest effective doses in children with persistent asthma.^{1,9}

In conclusion, the reduction in growth seen in the first few years of administration of inhaled glucocorticoids in prepubertal children persists as lowered adult height. However, in the information about inhaled glucocorticoids and their side effects that is provided to parents, the potential effect on adult height must be balanced against the large and well-established benefit of these drugs in controlling persistent asthma. It is appropriate to use the lowest effective dose for symptom control in order to minimize concern about the effects of inhaled glucocorticoids on adult height.

Supported by contracts with the National Heart, Lung, and Blood Institute (NO1-HR-16044, 16045, 16046, 16047, 16048, 16049, 16050, 16051, and 16052) and General Clinical Research Center grants from the National Center for Research Resources (M01RR00051, M01RR0099718-24, M01RR02719-14, and RR00036). Phases 2 and 3 of the CAMP Continuation Study were supported by grants from the National Heart, Lung, and Blood Institute (U01HL075232, U01HL075407, U01HL075408, U01HL075409, U01HL075415, U01HL075416, U01HL075417, U01HL075419, U01HL075420, and U01HL075408).

Dr. Kelly reports serving on steering committees for and receiving consulting fees from AstraZeneca, GlaxoSmithKline, Merck, and Novartis; Dr. Fuhlbrigge, receiving consulting fees from

Merck, GlaxoSmithKline, Dmogi, Lovelace Respiratory Research Institute, and Sunovion and serving as a member of an adjudication committee for studies sponsored by AstraZeneca, GlaxoSmithKline, Merck, and Novartis; Dr. Williams, receiving lecture fees from GlaxoSmithKline; and Dr. Zeiger, serving on a steering committee for a study sponsored by GlaxoSmithKline and receiving consulting fees from Aerocrine, AstraZeneca, Genentech, GlaxoSmithKline, MedImmune, Novartis, Schering-Plough, and Sunovion and grant support from Aerocrine, Genentech, GlaxoSmithKline, MedImmune, Merck, and Thermofisher. No other potential conflict of interest relevant to this article was reported.

Disclosure forms provided by the authors are available with the full text of this article at NEJM.org.

REFERENCES

1. Expert Panel Report 3 (EPR3): guidelines for the diagnosis and management of asthma. Bethesda, MD: National Institutes of Health, National Heart, Lung, and Blood Institute, 2007 (<http://www.nhlbi.nih.gov/guidelines/asthma>).
2. The Childhood Asthma Management Program Research Group. Long-term effects of budesonide or nedocromil in children with asthma. *N Engl J Med* 2000;343:1054-63.
3. Pauwels RA, Pedersen S, Busse WW, et al. Early intervention with budesonide in mild persistent asthma: a randomized, double-blind trial. *Lancet* 2003;361:1071-6.
4. Sharek PJ, Bergman DA. The effect of inhaled steroids on linear growth of children with asthma: a meta-analysis. *Pediatrics* 2000;106(1):e8.
5. Ferguson AC, Van Bever HP, Teper AM, Lasysya O, Goldfrad CH, Whitehead PJ. A comparison of the relative growth velocities with budesonide and fluticasone propionate in children with asthma. *Respir Med* 2007;101:118-29.
6. de Benedictis FM, Teper A, Green RJ, et al. Effects of 2 inhaled corticosteroids on growth: results of a randomized controlled trial. *Arch Pediatr Adolesc Med* 2001;155:1248-54.
7. Acun C, Tomac N, Ermis B, et al. Effects of inhaled corticosteroids on growth in asthmatic children: a comparison of fluticasone propionate with budesonide. *Allergy Asthma Proc* 2005;26:204-6.
8. Guilbert TW, Morgan WJ, Zeiger RS, et al. Long-term inhaled corticosteroids in preschool children at high risk for asthma. *N Engl J Med* 2006;354:1985-97.
9. Kelly HW. Comparison of inhaled corticosteroids: an update. *Ann Pharmacother* 2009;43:519-27.
10. Van Bever HP, Desager KN, Lijssens N, Weyler JJ, Du Caju MVL. Does treatment of asthmatic children with inhaled corticosteroids affect their adult height? *Pediatr Pulmonol* 1999;27:369-75.
11. Agertoft L, Pedersen S. Effect of long-term treatment with budesonide on adult height in children with asthma. *N Engl J Med* 2000;343:1064-9.
12. Strunk RC, Sternberg AL, Szefer SJ, et al. Long-term budesonide or nedocromil treatment, once discontinued, does not alter the course of mild to moderate asthma in children and adolescents. *J Pediatr* 2009;154:682-7.
13. Silverstein MD, Yunginger JW, Reed CE, et al. Attained adult height after childhood asthma: effect of glucocorticoid therapy. *J Allergy Clin Immunol* 1997;99:466-74.
14. Inoue T, Doi S, Takamatsu I, Murayama N, Kameda M, Toyoshima K. Effect of long-term treatment with inhaled beclomethasone dipropionate on growth of asthmatic children. *J Asthma* 1999;36:159-64.
15. Larsson L, Gerhardsson de Verdier M, Lindmark B, Norjavarra E. Budesonide-treated asthmatic adolescents attain target height: a population-based follow-up study from Sweden. *Pharmacoepidemiol Drug Saf* 2002;11:715-20.
16. Pedersen S. Clinical safety of inhaled corticosteroids for asthma in children: an update of long-term trials. *Drug Saf* 2006;29:599-612.
17. Childhood Asthma Management Program Research Group. The Childhood Asthma Management Program (CAMP): design, rationale, and methods. *Control Clin Trials* 1999;20:91-120.
18. Tanner JM, Whitehouse RH, Marshall WA, Carter BS. Prediction of adult height from height, bone age, and occurrence of menarche at ages 4 to 16 with allowance for midparent height. *Arch Dis Child* 1975;50:14-26.
19. Tanner JM, Landt KW, Cameron N, Carter BS, Patel J. Prediction of adult height from height and bone age in childhood: a new system of equations (TW Mark II) based on a sample including very tall and very short children. *Arch Dis Child* 1983;58:767-76.
20. Lee JJ, Escher JC, Shuman MJ, et al. Final adult height of children with inflammatory bowel disease is predicted by parental height and patient minimum height Z-score. *Inflamm Bowel Dis* 2010;16:1669-77.
21. Fine RN, Sullivan EK, Tejani A. The impact of recombinant human growth hormone treatment on final adult height. *Pediatr Nephrol* 2000;14:679-81.
22. Searle SR, Speed FM, Milliken GA. Population marginal means in the linear model: an alternative to least squares means. *Am Stat* 1980;34:216-21.
23. Stata multiple-imputation reference manual, release 12. College Station, TX: StataCorp, 2011.
24. Diggle PJ, Heagerty P, Liang KY, Zeger SL. Analysis of longitudinal data. 2nd ed. New York: Oxford University Press, 2002.
25. Tukey JW. Exploratory data analysis. Reading, MA: Addison-Wesley, 1977.
26. Tinkelman DG, Reed CE, Nelson HS, Offord KP. Aerosol beclomethasone dipropionate compared with theophylline as primary treatment of chronic, mild to moderately severe asthma in children. *Pediatrics* 1993;92:64-77.
27. Verberne AA, Frost C, Roorda RJ, van der Laag H, Kerrebijn KF. One year treatment with salmeterol compared with beclomethasone in children with asthma. *Am J Respir Crit Care Med* 1997;156:688-95.
28. Allen DB, Bronsky EA, LaForce CF, et al. Growth in asthmatic children treated with fluticasone propionate. *J Pediatr* 1998;132:472-7.
29. Skoner DP, Meltzer EO, Milgrom H, Stryczak P, Teper A, Staudinger H. Effects of inhaled mometasone furoate on growth velocity and adrenal function: a placebo-controlled trial in children 4-9 years old with mild persistent asthma. *J Asthma* 2011;48:848-59.
30. Snyder RD, Collipp PJ, Greene JS. Growth and ultimate height of children with asthma. *Clin Pediatr* 1967;6:389-92.
31. Murray AB, Fraser BM, Hardwick DF, Pirie GE. Chronic asthma and growth failure in children. *Lancet* 1976;2:197-8.
32. Ferguson AC, Murray AB, Tze W-J. Short stature and delayed skeletal maturation in children with allergic disease. *J Allergy Clin Immunol* 1982;69:461-6.
33. Patel L, Clayton PE, Addison GM, Price DA, David TJ. Linear growth in prepubertal children with atopic dermatitis. *Arch Dis Child* 1998;79:169-72.

Copyright © 2012 Massachusetts Medical Society.