RESEARCH SUMMARY

LUCAS R. KERBS

1. Introduction

I am a geometric topologist studying dynamical structures in 3-manifolds and geometric group theory. Thurston pioneered groundbreaking techniques that bridged low dimensional topology, hyperbolic geometry, and dynamical systems. A common motif is simplifying the dynamics by introducing lower-dimensional structures carrying those same dynamics—what started with a careful study of manifolds that fiber over the circle has grown into a field ripe for exploration. My work involves using the large-scale geometry of these dynamical systems to understand the manifolds and groups supporting them.

The prototypical example of this simplification technique is that of a ***universal circle***, a circle S^1_u together with an action of the fundamental group $\pi_1 M \curvearrowright S^1_u$. These circles are associated to many different dynamical structures and distinct structures on the same manifold will produce different universal circles. Universal circles can usually (and conjecturally always) be studied via $\pi_1 M$-equivariant ***sphere filling curves*** $S^1_u \to S^2$ which contain additional dynamical information. In the group theoretic setting, these same techniques unlock new descriptions of hyperbolic groups and their splittings. The bulk of my work has been focused on two fronts:

(1) In work with my advisor Steven Frankel, we show that any sphere filling curve $S^1 \to S^2$ approximated by embeddings has the same rich structure as those coming from 3-manifold constructions. [FK26].
(2) In work with Maxwell Plummer, we construct and classify the universal Cantor sets of a free-by-cyclic group in a way that parallels how the Thurston norm organizes universal circles. [KP26]

In the future, I plan to build on this work and:

(3) Use (1) to construct 3-manifolds from sphere-filling curves.
(4) Use (1) to construct new sphere-filling curves associated to dynamical structures which are not known to admit them.
(5) Compare the maps produced in (2) to one another to better understand the various splittings of a free-by-cyclic group.
(6) Generalize the results of (2) to topological structures in these groups that do *not* come from group-theoretic splittings.

2. Motivation: Fibered 3-manifolds and Cannon-Thurston Maps

Suppose that $f : \Sigma \to \Sigma$ is a homeomorphism of a closed surface with genus at least two. Using f, we may create a 3-manifold as follows: First, observe that $\Sigma \times [0,1]$ is a compact 3-manifold with boundary given by two copies of Σ. We then create M_f, the ***mapping torus*** of f, by gluing one boundary component to the other along f:

$$M_f = \Sigma \times [0,1]/(p,0) \sim (f(p),1)$$

Such an M_f is a closed 3-manifold with a fibration $M_f \to S^1$ whose point-preimages are each a copy of Σ. Further, the dynamics of f determines the geometric structure of M_f. A celebrated theorem of Thurston is that M_f is hyperbolic (a geometric property) if and only if no power of f fixes the isotopy class of any simple closed curve on Σ (a topological property) [Thu88]. Homeomorphisms with this property are called ***pseudo-Anosov***.

M_f comes with a ***suspension flow*** φ given by flowing in the 'vertical' direction which is transverse to the ***foliation by fibers***, the image of $\{\Sigma \times \{x\} \mid x \in [0,1]\}$ under the quotient map. We construct the ***orbit space*** of φ by lifting φ to a flow $\tilde{\varphi}$ on the universal cover $\tilde{M}$ and quotienting each flow line of $\tilde{\varphi}$ to a point: $\mathcal{O}_\varphi = \tilde{M}/\tilde{\varphi}$. The orbit space $\mathcal{O}_\varphi$ is the prototypical example of a ***bi-foliated plane***—a plane P supporting a pair of foliations (decompositions of P into lines and pronged singularities) which meet transversely at every point.

Giving Σ and M_f their natural hyperbolic structures, we may identify their universal covers with hyperbolic 2- and 3-space respectively: as such, the inclusion $\Sigma \hookrightarrow M_f$ lifts to a map $\mathbb{H}^2 \to \mathbb{H}^3$. A celebrated theorem of Cannon and Thurston [CT07] is that this map extends continuously to their natural compactifications $\mathbb{H}^2 \sqcup S^1_\infty \to \mathbb{H}^3 \sqcup S^2_\infty$. Restricting our attention to the boundaries, the ***Cannon-Thurston map*** $CT : S^1_\infty \to S^2_\infty$ is $\pi_1 M$ equivariant. As the action $\pi_1 M \curvearrowright S^2_\infty$ is minimal, the image is dense—thus CT is surjective.

These results have two main directions for generalization. Staying in the world of 3-manifolds we can build additional topological or dynamical structures and construct their universal circles. On the other hand, Cannon and Thurston's construction has a natural group theoretic analogue—building the geometric group theory of certain hyperbolic groups fibering over the circle.

2.1. **Overview.** My research program can be broadly divided into two subfields and I devote a section below to each of them. "Cannon-Thurston Maps onto S^2" covers the work done in [FK26] in the setting of hyperbolic 3-manifolds. "Cannon-Thurston Maps for Free-by-Cyclic Groups" covers the work done in [KP26] generalizing well-known results in 3-manifolds to a new setting.

3. Cannon-Thurston Maps onto S^2

A flow ψ on a closed hyperbolic 3-manifold is a ***pseudo-Anosov flow*** if it is locally modeled on the suspension flow of a pseudo-Anosov map. The orbit space $\mathcal{O}_\psi$ is topologically a bifoliated plane whose foliations are determined by the dynamics of the flow [FM01]. In [Fen12], Fenley shows that $\mathcal{O}_\psi$ can be compactified into a closed disk $\mathcal{D}_\psi = \mathcal{O}_\psi \sqcup S^1_\psi$ and that there is a Cannon-Thurston map $S^1_\psi \to S^2_\infty$. On the other hand, a flow φ on a closed hyperbolic 3-manifold is a ***quasigeodesic flow*** if, after lifting φ to a flow $\tilde{\varphi}$ on $\tilde{M}$, each flowline of $\tilde{\varphi}$ stays a bounded distance from a geodesic; as such every flow line determines two points in S^2_∞. By work of Calegari, the orbit space $\mathcal{O}_\varphi$ of a quasigeodesic flow is a plane which can be naturally compactified to a closed disk, $\mathcal{D}_\varphi = \mathcal{O}_\varphi \sqcup S^1_\varphi$ [Cal06], Frankel then shows the existence of a Cannon-Thurston Map $S^1_\varphi \to S^2_\infty$ for φ [Fra15].

Recent exciting work of Frankel and Landry unifies these two pictures: If φ is a quasigeodesic flow on M, then there is a closed, flow-invariant subset $M_L \subset M$ and a surjection $p : M_L \to M$ such that p semiconjugates the quasigeodesic flow on M_L to a pseudo-Anosov flow on M [FL25]. In particular, this is done *without* changing the Cannon-Thurston map. The key insight is that one can modify the flow space by considering the corresponding ***circle laminations***—a closed set of unordered pairs of points in S^1; the laminations provide data about how the Cannon-Thurston map identifies points on S^1_φ.

In its simplest form, a ***universal circle*** for a 3-manifold is a circle S^1_u and an action $\pi_1 M \curvearrowright S^1_u$ which contains information about a geometric or dynamical structure in M. The prototypical examples are the circle at the boundary of the orbit space S^1_φ of a quasigeodesic pseudo-Anosov flow

and the ***leftmost/rightmost circles*** $\mathcal{C}^{\ell/r}$ of an ***Anosov foliation***. Recent work by Buckminster [Buc26] produces a Cannon-Thurston map for $\mathcal{C}^{\ell/r}$, inspiring the following question:

Question 1. *Suppose that S^1_u is a universal circle for M. Does there exist a Cannon-Thurston map $\varphi : S^1_u \to S^2_\infty$?*

While universal circles arising from orbit spaces come with the additional data in the form of a pair of circle laminations and corresponding bi-foliated plane, this is not always the case—$\mathcal{C}^{\ell/r}$ lack this structure. Conjecturally, every universal circle is the boundary of the orbit space of some pseudo-Anosov flow [LMT24].

Question 2. *Suppose that S^1_u is a universal circle for M admitting a Cannon-Thurston map $\varphi : S^1_u \to S^2_\infty$. Does φ admit a pair of laminations extending to a bifoliated plane?*

My work with Frankel opens a new angle of attack for this question:

Theorem 3.1 (Frankel-K. [FK26])**.** *Suppose that $f_\infty \colon S^1 \to S^2$ is a sphere filling curve and that there is a continuous family $f_t : S^1 \to S^2$, $t \in [0, \infty]$, such that f_t is an embedding when $t < \infty$. Then f_∞ supports a pair of laminations $\Lambda^\pm$ extending to bifoliated plane.*

Thus, we have reduced Question 2 to showing that the Cannon-Thurston maps are approximated by embeddings. Further, the construction in Theorem 3.1 is natural in the sense that if $f_\infty : S^1 \to S^2$ is equivariant under the action of some group G, then the action of G in S^1 preserves the laminations $\Lambda^\pm$ and G acts on the corresponding bifoliated plane.

This work is a part of a larger program by Calegari as work towards ***Cannon's Conjecture***. Let M be a compact, metrizable space and $\Theta(M)$ the space of distinct triples of point in M. A group action $G \curvearrowright M$ is a ***uniform convergence group action*** if the induced action $G \curvearrowright \Theta(M)$ is properly discontinuous and co-compact. Combining results of Tukia, Gabai, Casson-Jungreis and Freden, [Tuk86; Gab92; CJ94; Fre95] we have the following characterization of uniform convergence group action on circles.

Theorem 3.2. *Suppose that G acts as a uniform convergence group action on S^1. Then there is a discrete group $\Gamma \subset \mathrm{Isom}(\mathbb{H}^2)$ such that $\Gamma \curvearrowright \mathbb{H}^2 \sqcup \partial_\infty \mathbb{H}^2$ is properly discontinuous and cocompact and the actions $G \curvearrowright S^1$ and $\Gamma \curvearrowright \partial_\infty \mathbb{H}^2$ are conjugate.*

A natural follow up question is to ask if there are similar statements for $G \curvearrowright S^n$ and $\Gamma \subset \mathrm{Isom}(\mathbb{H}^{n+1})$. If $n > 2$, the statement is false; Cannon's Conjecture is the case where $n = 2$:

Conjecture 3.3 (Cannon's Conjecture)**.** *Suppose that G acts as a uniform convergence group on S^2. Then G is conjugate to a discrete group $\Gamma \subset \mathrm{Isom}(\mathbb{H}^3)$.*

If the group G is torsion free, then Cannon's Conjecture implies that G is $\pi_1 M$ for some closed, hyperbolic 3-manifold M. In their September release, OpenAI claims to have resolved Cannon's Conjecture positively, [Ope26], but their proof does not interface with any internal dynamical structures (flows, foliations, etc.) of the corresponding 3-manifold, an important part of Calegari's program.

Calegari and Loukidou introduce ***CaTherine Wheels*** and ***Zippers*** [CL24; CL26] as the start of a dictionary between structures on 3-manifolds and structures on S^2. CaTherine wheels are a class of sphere filling curves $f \colon S^1 \to S^2$ such that for every closed interval $I \subset S^1$, $f(I)$ is a closed disk and $f(\partial I) \subset \partial f(I)$. All such maps are approximated by embeddings and Theorem 3.1 is a key step in developing their dictionary. Importantly however, there are many topological and dynamical structures whose Cannon-Thurston maps are *not* CaTherine Wheels—but they ***are*** approximated by embeddings. The proper generalizations—P-Zippers and P-CaTherine Wheels—are yet to be formalized and the ideas are rich for exploration. As such, a major future direction for my research is to build 3-manifolds from their Cannon-Thurston Maps:

Conjecture 3.4. *Suppose G acts as a uniform convergence group on S^2 and that $f\colon S^1 \to S^2$ is a G-invariant sphere filling curve approximated by embeddings. Then there is a closed, hyperbolic 3-manifold M and a quasigeodesic flow φ on M such that $G = \pi_1 M$ and f is the Cannon-Thurston map for φ.*

4. Cannon-Thurston Maps for Free-by-Cyclic Groups

A celebrated result of Mj says that whenever

$$1 \to H \to G \to Q \to 1$$

is a short exact sequence of infinite word hyperbolic groups, there is a surjective Cannon-Thurston map $\partial_\infty H \to \partial_\infty G$ between the Gromov boundaries of H and G [Mit98]. Cannon and Thurston's result is then the case where $H = \pi_1\Sigma$, $G = \pi_1 M$, and $Q = \mathbb{Z}$; notice that we can write G as the semidirect product $\pi_1 M \simeq \pi_1\Sigma \rtimes_{f_*} \mathbb{Z}$ where $f_*\colon \pi_1\Sigma \to \pi_1\Sigma$ is induced by gluing map $f : \Sigma \to \Sigma$.

Let F_N denote the free group on N letters; given an outer automorphism $\varphi \in \mathrm{Out}(F_N)$, we may form the ***free-by-cyclic*** $G \simeq F_N \rtimes_\varphi \mathbb{Z}$. As long φ is fully irreducible and atoroidal (technical conditions on φ to ensure it is sufficiently mixing), G is a hyperbolic group that shares many properties with a fibered hyperbolic 3-manifold. By the aforementioned result of Mj, there is a Cannon-Thurston map $\partial_\infty F_N \to \partial_\infty G$. Unlike the fibered setting, these spaces at infinity are less well-behaved: if $N \geq 3$, then $\partial_\infty F_N$ is a Cantor set and $\partial_\infty G$ is the Menger Sponge [KK00].

While a hyperbolic free-by-cyclic group $G \simeq F_N \rtimes_\varphi \mathbb{Z}$ is a purely group theoretic construction, there are many ways to reintroduce topology. An ***expanding train track map*** for φ is a graph Γ with smoothing data around each vertex and a homotopy equivalence $f\colon \Gamma \to \Gamma$ inducing φ on $\pi_1\Gamma$. Dowall-Kapovich-Leininger use the data of an expanding train track map to construct the ***folding mapping torus***, a $K(G,1)$ with a fibration $X_f \to S^1$ whose point preimages are graphs homotopy equivalent to Γ and monodromy f [DKL15]. The folded mapping torus comes equipped with the ***suspension semi-flow*** ψ_t (an action $\mathbb{R}_+ \curvearrowright X_f$) defined analogously to the suspension flow on a mapping torus; as f is not invertible, we can only flow forwards, hence the restriction to a semi-flow.

The first natural question about a hyperbolic free-by-cyclic group is if the splitting is unique. In general, the answer is a resounding no—most free-by-cyclic groups split infinitely many ways with infinitely many different monodromies. The splittings of G are governed by the ***symmetric BNS invariant***, a finite union of open cones $\Sigma_s(G) \subset H^1(G;\mathbb{R})$ [BNS87]. A surjective homomorphism $u : G \to \mathbb{Z}$ satisfies $[u] \in \Sigma_s(G)$ if and only if $\ker u$ is finitely generated; in this case, $\ker u$ is free and we can write $G \simeq \ker u \rtimes \mathbb{Z}$ [GMSW01]. The folded mapping torus is an invariant of an entire component of the BNS invariant. If $[u]$ and $[v]$ belong to the same component of the BNS invariant with monodromies φ, ψ and expanding train track maps $f : \Gamma \to \Gamma$ and $g : \Theta \to \Theta$, then there is a homeomorphism $X_f \to X_g$ between their folded mapping tori sending semiflow lines to semiflow lines and inducing an isomorphism between the splittings $F_N \rtimes_\phi \mathbb{Z} \simeq G \simeq F_{N'} \rtimes_\psi \mathbb{Z}$. The utility of the BNS invariant motivates:

Question 3. *Given a splitting of a free-by-cyclic group $G \simeq F_N \rtimes_\varphi \mathbb{Z}$, what properties of the splitting depend only on the group G? Which properties depend only on the component of the BNS Invariant?*

Question 3 has many partial answers. For example, Mutanguha proved that (full) irreducibility of φ is an invariant of the group [Mut21], building on work of Dowall–Kapovich–Leininger, who showed it was an invariant of a component of the BNS invariant [DKL17]. Maxwell Plummer and I answer Question 3 for the Cannon-Thurston Map.

Theorem 4.1 (K.-Plummer [KP26])**.** *Let $G \simeq F_N \rtimes_\varphi \mathbb{Z}$ be a hyperbolic free-by-cyclic group with fully irreducible monodromy. The Cannon-Thurston map $\partial_\infty F_N \to \partial_\infty G$ depends only on the component of the BNS Invariant.*

We construct a ***Universal Cantor*** set $\mathfrak{C}$ and a map $\mathcal{R} : \mathfrak{C} \to \partial_\infty G$ such that for any $u : G \to \mathbb{Z}$ which belongs to the same component as φ there is an identification $h_u \colon \partial_\infty \ker u \to \mathfrak{C}$ such that the Cannon-Thurston map $CT_u : \ker u \to \partial_\infty G$ factors as $CT_u = \mathcal{R} \circ h_u$.

Building on this result, I hope to compare the universal Cantor sets across different components of the BNS Invariant. In the world of 3-manifolds, universal circles from different components of the BNS Invariant for $\pi_1 M$ are distinct in both their $\pi_1 M$ actions and their Cannon-Thurston maps. Thus, I hope to prove

Conjecture 4.2. *Suppose that G is a hyperbolic free-by-cyclic group with fully irreducible monodromy and let $\mathfrak{C}$ and $\mathfrak{C}'$ denote the universal Cantor sets associated to distinct component of $\Sigma_s(G)$. Then the actions $G \curvearrowright \mathfrak{C}$ and $G \curvearrowright \mathfrak{C}'$ are not conjugate.*

References

[BNS87] Robert Bieri, Walter D. Neumann, and Ralph Strebel. "A Geometric Invariant of Discrete Groups". In: ***Inventiones mathematicae*** 90.3 (Oct. 1987), pp. 451–477. DOI: 10.1007/BF01389175.

[Buc26] Ellis Buckminster. ***Cannon–Thurston Maps for Anosov Foliations***. Apr. 2026. DOI: 10.48550/arXiv.2604.21201. arXiv: 2604.21201 [math.GT].

[Cal06] Danny Calegari. "Universal Circles for Quasigeodesic Flows". In: ***Geometry & Topology*** 10.4 (Nov. 2006), pp. 2271–2298. DOI: 10.2140/gt.2006.10.2271. arXiv: math/0406040.

[CJ94] Andrew Casson and Douglas Jungreis. "Convergence Groups and Seifert Fibered 3-Manifolds". In: ***Inventiones mathematicae*** 118.1 (Dec. 1994), pp. 441–456. DOI: 10.1007/BF01231540.

[CL24] Danny Calegari and Ino Loukidou. ***Zippers***. Nov. 2024. DOI: 10.48550/arXiv.2411.15610. arXiv: 2411.15610 [math].

[CL26] Danny Calegari and Ino Loukidou. ***CaTherine Wheels***. https://arxiv.org/abs/2604.24619v1. Apr. 2026.

[CT07] James W Cannon and William P Thurston. "Group Invariant Peano Curves". In: ***Geometry & Topology*** 11.3 (July 2007), pp. 1315–1355. DOI: 10.2140/gt.2007.11.1315.

[DKL15] Spencer Dowdall, Ilya Kapovich, and Christopher J. Leininger. "Dynamics on Free-by-Cyclic Groups". In: ***Geometry & Topology*** 19.5 (Oct. 2015), pp. 2801–2899. DOI: 10.2140/gt.2015.19.2801. arXiv: 1301.7739 [math].

[DKL17] Spencer Dowdall, Ilya Kapovich, and Christopher J. Leininger. ***Endomorphisms, Train Track Maps, and Fully Irreducible Monodromies***. Aug. 2017. DOI: 10.48550/arXiv.1507.03028. arXiv: 1507.03028 [math.GR].

[Fen12] Sergio R. Fenley. "Ideal Boundaries of Pseudo-Anosov Flows and Uniform Convergence Groups, with Connections and Applications to Large Scale Geometry". In: ***Geometry & Topology*** 16.1 (Jan. 2012), pp. 1–110. DOI: 10.2140/gt.2012.16.1. arXiv: math/0507153.

[FK26] Steven Frankel and Lucas Kerbs. "Pinched Jordan Curves, Cannon Thurston Maps, and Their Decompositions". In: (2026).

[FL25] Steven Frankel and Michael Landry. ***From Quasigeodesic to Pseudo-Anosov Flows***. Oct. 2025. DOI: 10.48550/arXiv.2510.02217. arXiv: 2510.02217 [math].

[FM01] Sérgio Fenley and Lee Mosher. "Quasigeodesic Flows in Hyperbolic 3-Manifolds". In: ***Topology*** 40.3 (May 2001), pp. 503–537. DOI: 10.1016/S0040-9383(99)00072-5.

[Fra15] Steven Frankel. "Quasigeodesic Flows and Sphere-Filling Curves". In: ***Geometry & Topology*** 19.3 (May 2015), pp. 1249–1262. DOI: 10.2140/gt.2015.19.1249. arXiv: 1210.7050 [math].

[Fre95] Eric M. Freden. "Negatively Curved Groups Have the Convergence Property I". In: ***Annales Fennici Mathematici*** 20.2 (Aug. 1995), pp. 333–348.

[Gab92] David Gabai. "Convergence Groups Are Fuchsian Groups". In: ***The Annals of Mathematics*** 136.3 (Nov. 1992), p. 447. DOI: 10.2307/2946597. JSTOR: 2946597.

[GMSW01] Ross Geoghegan, Michael L. Mihalik, Mark Sapir, and Daniel T. Wise. "ASCENDING HNN EXTENSIONS OF FINITELY GENERATED FREE GROUPS ARE HOPFIAN". In: ***Bulletin of the London Mathematical Society*** 33.3 (May 2001), pp. 292–298. DOI: 10.1017/S0024609301007986.

[KK00] Michael Kapovich and Bruce Kleiner. "Hyperbolic Groups with Low-Dimensional Boundary". In: ***Annales Scientifiques de l'École Normale Supérieure*** 33.5 (Sept. 2000), pp. 647–669. DOI: 10.1016/S0012-9593(00)01049-1.

[KP26] Lucas Kerbs and Maxwell Plummer. "Universal Cantor Sets and Cannon Thurston Maps for Free by Cyclic Groups". In: (2026).

[LMT24] Michael P. Landry, Yair N. Minsky, and Samuel J. Taylor. ***Simultaneous Universal Circles***. Dec. 2024. DOI: 10.48550/arXiv.2412.06986. arXiv: 2412.06986 [math.GT].

[Mit98] Mahan Mitra. "Cannon-Thurston Maps for Hyperbolic Group Extensions". In: ***Topology*** 37.3 (May 1998), pp. 527–538. DOI: 10.1016/S0040-9383(97)00036-0.

[Mut21] Jean Pierre Mutanguha. "Irreducibility of a Free Group Endomorphism Is a Mapping Torus Invariant". In: ***Commentarii Mathematici Helvetici*** 96.1 (Mar. 2021), pp. 47–63. DOI: 10.4171/cmh/506. arXiv: 1910.04285 [math.GR].

[Ope26] OpenAI. ***A Modulus Proof of Cannon's Conjecture***. 2026.

[Thu88] William P. Thurston. "On the Geometry and Dynamics of Diffeomorphisms of Surfaces". In: ***Bulletin (New Series) of the American Mathematical Society*** 19.2 (Oct. 1988), pp. 417–431.

[Tuk86] Pekka Tukia. "On Quasiconformal Groups". In: ***Journal d'Analyse Mathématique*** 46.1 (Dec. 1986), pp. 318–346. DOI: 10.1007/BF02796595.

Department of Mathematics, Washington University in St. Louis, St. Louis, MO

Email address: kerbs@wustl.edu